



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines


Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>



LANE

MEDICAL



LIBRARY

LEVI COOPER LANE FUND



AN INTRODUCTION
TO
PATHOLOGY AND MORBID ANATOMY.

CORNIL & RANVIER'S PATHOLOGICAL HISTOLOGY—Nearly Ready.

A MANUAL OF PATHOLOGICAL HISTOLOGY. By V. CORNIL, Prof. in the Faculty of Medicine, Paris, and L. RANVIER, Prof. in the College of France. Translated, with Notes and Additions, by E. O. SHAKESPEARE, M.D., Pathologist and Ophthalmic Surgeon to the Philada. Hospital. In one handsome octavo volume of about 600 pages, with about 300 illustrations.

So much has been done of late years in the elucidation of pathology by means of the microscope, and this subject now occupies so prominent a position as one of the most important branches of medical science, that the American profession cannot fail to welcome a translation of the present work, which, through its own merits and through the well-known reputation of its distinguished authors, is regarded in Europe as the standard text-book and work of reference in its department. Such investigations and discoveries as have been made since its appearance will be introduced by the translator, and the work is confidently expected to assume in this country the same position which has been so universally accorded to it abroad.

FINLAYSON'S CLINICAL DIAGNOSIS.—Nearly Ready.

CLINICAL MANUAL FOR THE STUDY OF MEDICAL CASES. For the use of Students and Practitioners of Medicine. Edited by JAMES FINLAYSON, M.D., Physician and Lecturer on Clinical Medicine in the Glasgow Medical Infirmary, etc. etc. With eighty-five illustrations on wood. In one very handsome volume, royal 12mo., of about 500 pages.

The scope of this useful and somewhat novel work can best be understood by the subjoined condensed

SUMMARY OF CONTENTS.

CHAPTER I. The Physiognomy of Disease. II. Examination and Reporting of Medical Cases. III. Temperature—Pulse—General Signs of Pyæmia. IV. Skin—Hair—Nails—Glands—Joints. V. Examination of the Organs of Special Sense. Subjective Disorders of the Special Senses—Testing of Cranial Nerves. VI. Symptoms of Disorder in the Nervous System. VII. The Use of Electrical Instruments. VIII. Insanity. IX. Disorders of the Respiratory and Circulatory Systems. X. Examination of the Fauces, Larynx, and Nares. XI. Disorders of the Digestive System. XII. Jaundice and Dropsy. XIII. Examination of the Urine and Significance of Urinary Symptoms. XIV. Symptoms connected with the Male Generative Organs. XV. Disorders of the Female Organs and their Relations to the General Health. XVI. Physical Examination of Chest and Abdomen—Lungs—Heart—Abdomen. XVII. Method of performing Post-Mortem Examinations.

The concurrence of gentlemen specially familiar with the several subjects being requisite to the satisfactory development of a plan so extensive, Dr. Finlayson has secured the coöperation of Professor Gairdner, who has contributed the chapter on the Physiognomy of Disease; Professor Wm. Stephenson that on Disorders of the Female Organs; Dr. Alex. Robertson that on Insanity; Prof. Samson Gemmell those on the Sphygmograph and Physical Diagnosis; and Dr. Joseph Coates those on the Fauces, Larynx, and Nares and on the method of performing *post mortem* examinations. Other chapters have enjoyed the advantage of revision by gentlemen specially versed in their several subjects; and the volume is presented as thoroughly on a level with the most advanced condition of knowledge in a department which has made such rapid strides of advancement within the last few years.

HENRY C. LEA—Philadelphia.

AN
INTRODUCTION
TO
PATHOLOGY AND MORBID ANATOMY.

T. HENRY GREEN, M.D., LOND.,
FELLOW OF THE ROYAL COLLEGE OF PHYSICIANS, LOND., PHYSICIAN TO CHARING
CROSS HOSPITAL AND LECTURER ON PATHOLOGY AND MORBID ANATOMY
AT CHARING CROSS HOSPITAL MEDICAL SCHOOL, ASSISTANT-
PHYSICIAN TO THE HOSPITAL FOR CONSUMPTION
AND DISEASES OF THE CHEST, BROMPTON.

THIRD AMERICAN
FROM THE
FOURTH REVISED AND ENLARGED ENGLISH EDITION.
WITH ONE HUNDRED AND THIRTY-TWO ILLUSTRATIONS.



PHILADELPHIA:
HENRY C. LEA.
1878.

B

Y9A9B1.3BA1

COLLINS, PRINTER.

379
1878

TO

WILSON FOX, M.D., F.R.S.,

PHYSICIAN EXTRAORDINARY TO HER MAJESTY THE QUEEN, PHYSICIAN IN ORDINARY
TO THEIR R. H.'S THE DUKE AND DUCHESS OF EDINBURGH, HOLMKS PROFESSOR
OF CLINICAL MEDICINE, FORMERLY PROFESSOR OF PATHOLOGICAL
ANATOMY AT UNIVERSITY COLLEGE, LONDON, AND
PHYSICIAN TO UNIVERSITY COLLEGE
HOSPITAL,

THIS SMALL WORK

Is Inscribed

WITH MUCH RESPECT AND GRATITUDE

BY HIS FORMER PUPIL,

THE AUTHOR.

PREFACE.

IN preparing the fourth edition of my text-book of Pathology and Morbid Anatomy, I have again added much new matter, with the object of making the work a more complete guide for the student. All the chapters have been carefully revised, and a large addition has been made to the number of wood-cuts. These new wood-cuts, as in previous editions, have been drawn by Mr. Collings from my own microscopical preparations.

T. HENRY GREEN.

74, WIMPOLE STREET, CAVENDISH SQUARE,

February, 1878.

CONTENTS.

INTRODUCTION.

	PAGE
Definitions—Health—Disease—Nutrition—Function—Blood as cause of Disease—Nervous System	17-19

CHAPTER I.

THE "CELL."

Cell as seat of Nutrition and Function—Constitution of Cells—Protoplasm—Nucleus—Cell-wall—Physiology of Cells—Genesis of Cells—Vacuolation	20-25
--	-------

CHAPTER II.

NUTRITION ARRESTED.

<i>Gangrene or Necrosis</i> —Results of Arrested Nutrition—Dry Gangrene—Moist Gangrene—Characters of Dead Part—Post-mortem Staining— <i>Rigor Mortis</i> —Nature of Changes in Muscle—Terminations of Gangrenous Process—Causes of Gangrene— <i>Senile Gangrene</i>	26-32
---	-------

CHAPTER III.

NUTRITION IMPAIRED.

<i>Atrophy</i> —Simple—Numerical—Necrobiosis—Causes—Physical Characters— <i>Atrophy of Bone</i> — <i>Pulmonary Emphysema</i> — <i>Degeneration</i> —Causes—The Metamorphoses—The Infiltrations	33-42
--	-------

CHAPTER IV.

FATTY DEGENERATION.

General Pathology—Sources of Fat—Causes of Accumulation—Phosphorus Poisoning—Incomplete Oxidation—Histology of Process— <i>Fatty Infiltration</i> — <i>Fatty Infiltration of Muscle, of Heart, of Liver</i>	43-51
---	-------

CHAPTER V.

FATTY DEGENERATION (*continued*).

PAGE

<i>Fatty Metamorphosis</i> —Histology of Process—Secretion of Milk—Absorption of Fat—Caseation—Calcification—Softening— <i>Fatty Degeneration of Arteries, of Capillaries</i> — <i>Fatty Degeneration of Brain</i> — <i>Cerebral Softenings</i> —White Softening—Yellow Softening—Red Softening— <i>Fatty Degeneration of Muscle</i> — <i>Fatty Degeneration of Heart</i> — <i>Brown Atrophy of Heart</i> — <i>Fatty Degeneration of Kidneys</i>	52-63
--	-------

CHAPTER VI.

MUCOID AND COLLOID DEGENERATION.

<i>Mucoid Degeneration</i> —Mucin— <i>Colloid Degeneration</i> —Colloid Degeneration of Tumors— <i>Muscular Changes in Typhoid Fever</i>	63-66
--	-------

CHAPTER VII.

AMYLOID DEGENERATION.

Circumstances under which it occurs—Nature of Amyloid Substance—Reaction with Iodine—Seat of Change—Physical Characters of affected Organs—Consequent Impairment of Nutrition—Source of Amyloid Substance— <i>Amyloid Degeneration of Liver</i> — <i>Amyloid Degeneration of Kidneys</i> — <i>Amyloid Degeneration of Spleen</i> — <i>Amyloid Degeneration of Lymphatic Glands</i> — <i>Amyloid Degeneration of Alimentary Canal</i> — <i>The Corpora Amylacea</i>	67-79
--	-------

CHAPTER VIII.

CALCAREOUS DEGENERATION.

Circumstances under which it occurs—Causes—Nature of Calcareous Particles—Effect on Tissue— <i>Calcification of Arteries</i>	80-83
--	-------

CHAPTER IX.

PIGMENTARY DEGENERATION.

Source of Pigment—Nature of Process—Hæmatoidin—Pathological Significance—False Pigmentation— <i>Pigmentation of Lungs</i> —Normal—Colliers' Lungs—Pigmentation in Disease	84-90
---	-------

CHAPTER X.

TISSUE-CHANGES IN PYREXIA.

Diseases in which they occur—Physical Characters of Organs—Histology—Nature of Change	91-93
---	-------

CONTENTS.

xi

CHAPTER XI.

NUTRITION INCREASED.

	PAGE
<i>Hypertrophy</i> —Simple Hypertrophy—Numerical Hypertrophy—Functional Hypertrophies	93-94

CHAPTER XII.

NEW FORMATIONS.

Inflammatory Formations— <i>Tumors</i> —Etiology—Constitutional Predisposing Causes—Dyscrasia—Cachexia—Hereditary Predisposition—Direct Exciting Causes—Irritants—Emigration and Transmission of Cellular Elements—Spermatic Influence—Development—Indifferent Cells—Homology—Heterology—Relation of Growth to surrounding Tissues—Retrogressive Changes— <i>Malignancy</i> —Classification	95-109
---	--------

CHAPTER XIII.

THE FIBROMATA.

Structure—Development—Secondary Changes—Varieties—Soft Fibromata—Firm Fibromata—Epulis—Clinical Characters	109-111
--	---------

CHAPTER XIV.

THE LIPOMATA.

Structure—Development—Secondary Changes—Physical Characters—Clinical Characters	112-113
---	---------

CHAPTER XV.

THE ENCHONDROMATA.

Structure—Development—Secondary Changes—Varieties—Physical Characters—Clinical Characters	113-116
---	---------

CHAPTER XVI.

THE OSTEOMATA.

Structure—Development—Exostoses—Osteophytes—Clinical Characters	117-118
---	---------

CHAPTER XVII.

THE MYXOMATA.

Structure—Development—Secondary Changes—Varieties—Physical Characters—Clinical Characters	119-120
---	---------

CHAPTER XVIII.

THE LYMPHOMATA.

	PAGE
Lymphadenoma—Structure—Development—Clinical Characters— <i>Hodg-kin's Disease</i>	121-125

CHAPTER XIX.

THE MYOMATA, NEUROMATA, AND ANGIOMATA.

Structure, etc.— <i>Myoma of Uterus</i>	126-128
---	---------

CHAPTER XX.

THE PAPILLOMATA.

Structure—Development—Secondary Changes—Varieties—Clinical Characters	129-131
---	---------

CHAPTER XXI.

THE ADENOMATA.

Structure—Development—Secondary Changes— <i>Adenoma of Mamma—Adeno-Fibroma, and Adeno-Sarcoma of Mamma—Adenoma of Mucous Membrane</i> —Clinical Characters	131-134
--	---------

CHAPTER XXII.

THE SARCOMATA.

Structure—Development—Secondary Changes—Varieties— <i>Spindle-celled Sarcoma—Melanotic Sarcoma—Osteoid Sarcoma—Round-celled Sarcoma—Glioma—Alveolar Sarcoma—Myeloid Sarcoma—Psammoma—Blood-cysts</i> —Clinical Characters	135-147
---	---------

CHAPTER XXIII.

THE CARCINOMATA.

Definition—Structure—Development—Secondary Changes—Varieties— <i>Scirrhus—Encephaloid—Colloid—Epithelioma—Cylindrical Epithelioma</i> —Clinical Characters	147-161
--	---------

CHAPTER XXIV.

CYSTS.

Origin—Structure—Secondary Changes—Classification	161-164
---	---------

CHAPTER XXV.

INFLAMMATION.

PAGE

Definition—Changes in Bloodvessels and Circulation—Stasis—Migration of Blood-Corpuscles—Exudation of Liquor Sanguinis—Alterations in Nutrition of Inflamed Tissue—Conclusions—*Suppuration*—Origin of Pus—Varieties of Inflammation—Acute Inflammations—Chronic Inflammations—Infective Inflammations—Specific Inflammations . . . 164–178

CHAPTER XXVI.

SCROFULOUS INFLAMMATION.

Scrofula—Scrofulous Lesions—Peculiarities of Scrofulous Inflammation 179–181

CHAPTER XXVII.

TUBERCLE AND ACUTE TUBERCULOSIS.

Definition—General Pathology—Older Doctrines—Gray and Yellow Tubercle—Infective Nature of Acute Tuberculosis—Artificial Production of Tuberculosis in Lower Animals—Source of Infection—Histology—The Giant Cells—Differences in Histological Characters of Lesions—Secondary Changes—Local Tuberculosis—Conclusions—*Tuberculosis of Pia Mater*—*Tuberculous Masses in Brain*—*Tuberculosis of Lymphatic Glands*—*Tuberculosis of Mucous Membranes*—*Tuberculosis of Lungs* . . . 181–199

CHAPTER XXVIII.

PYÆMIA AND SEPTICÆMIA.

Traumatic or Surgical Fever—*Septicæmia*—Nature of Poison—Bacteria—Absorption of Poison—Changes produced by Poison in Blood—*Pyæmia*—Relation of *Pyæmia* to *Septicæmia*—Suppurative Arthritis—Pathology of Metastatic Abscesses—Multiplication of Poison . . . 199–206

CHAPTER XXIX.

SYPHILIS.

Nature of Syphilitic Lesions—Fibroid Changes—*The Gummata*—Their Structure—Development—Secondary Changes—*Changes in Arteries*—*Syphilitic Disease of Liver* . . . 206–212

CHAPTER XXX.

INFLAMMATION OF NON-VASCULAR TISSUES.

Inflammation of Cartilage—*Inflammation of Cornea* . . . 212–214

CHAPTER XXXI.

INFLAMMATION OF COMMON CONNECTIVE TISSUE.

	PAGE
Histological Changes—Resolution—Organization—Cicatrical Tissue—Sup- puration—Granulation	214–218

CHAPTER XXXII.

INFLAMMATION OF BONE.

Histological Changes—Osteo-Myelitis—Sclerosis— <i>Periostitis</i> — <i>Necrosis</i> — <i>Caries</i> — <i>Mollities Ossium</i> — <i>Rickets</i>	218–223
---	---------

CHAPTER XXXIII.

INFLAMMATION OF BLOODVESSELS.

<i>Inflammation of Arteries</i> —Acute Inflammations— <i>Atheroma</i> — <i>Inflammation</i> <i>of Veins</i>	223–226
--	---------

CHAPTER XXXIV.

INFLAMMATION OF THE HEART.

<i>Endocarditis</i> —Acute—Chronic— <i>Myocarditis</i> —Acute—Chronic— <i>Fibroid</i> <i>Induration of Heart</i>	226–232
---	---------

CHAPTER XXXV.

INFLAMMATION OF LYMPHATIC STRUCTURES.

Acute Inflammation—Chronic Inflammation—Scrofulous Glands— <i>Typhoid</i> <i>Fever</i>	232–236
---	---------

CHAPTER XXXVI.

INFLAMMATION OF MUCOUS MEMBRANES.

Catarrhal Inflammation—Fibrinous and Croupous Inflammations— <i>Croup</i> — <i>Diphtheria</i> — <i>Dysentery</i>	237–242
---	---------

CHAPTER XXXVII.

INFLAMMATION OF SEROUS MEMBRANES.

Adhesive Inflammation—Effusion—Suppuration	242–244
--	---------

CHAPTER XXXVIII.

INFLAMMATION OF THE LIVER.

<i>Hepatic Abscess</i> — <i>Cirrhosis</i> — <i>Acute Yellow Atrophy</i>	245–248
---	---------

CONTENTS.

xv

CHAPTER XXXIX.

INFLAMMATION OF THE KIDNEY.

	PAGE
<i>Suppurative Nephritis—Surgical Kidney—Tubal Nephritis—Scarlatinal Nephritis—Interstitial Nephritis</i>	248-259

CHAPTER XL.

INFLAMMATION OF BRAIN AND SPINAL CORD.

<i>Acute Inflammation—Abscess—Inflammatory Softening—Sclerosis</i>	259-262
--	---------

CHAPTER XLI.

INFLAMMATION OF THE LUNGS.

<i>Croupous Pneumonia—Engorgement—Red Hepatization—Gray Hepatization—Terminations of Process—Broncho- or Catarrhal Pneumonia—Hypostatic Pneumonia—Interstitial or Chronic Pneumonia</i>	262-277
---	---------

CHAPTER XLII.

PULMONARY PHTHISIS.

<i>Definition—Older Doctrines—Tubercle as a Cause of Phthisis—Histology of Phthisis—Epithelial Proliferation—Fibrinous Exudation—Cellular Infiltration of Alveolar Walls—Increase of Interlobular Connective Tissue—Changes in Bronchi—Pathology of Phthisis—Nature of Morbid Processes—Causes of Differences in Histological Changes—Resolution—Fibroid Development—Retrograde Metamorphosis—Etiology—Predisposition—Inherent Pulmonary Weakness—Modes in which Injuries may be inflicted on Lungs—Infection—General Health—Varieties of Phthisis</i>	277-290
--	---------

CHAPTER XLIII.

CHANGES IN THE BLOOD AND CIRCULATION.

<i>Active Hyperæmia—Causes—Results—Mechanical Hyperæmia—Causes—Results—Nutmeg Liver—Brown Induration of Lung—Post-mortem Appearance of Hyperæmia</i>	291-298
--	---------

CHAPTER XLIV.

THROMBOSIS.

<i>Ante-mortem Clots—Post-mortem Clots—Organization of Thrombus—Softening—Suppuration—Causes—Results</i>	298-307
--	---------

CHAPTER XLV.

EMBOLISM.

	PAGE
Source of Emboli—Seat of Arrest—Secondary Thrombosis—Results—	
Changes in Vessels—Changes in Organ—Hemorrhagic Infarcts— <i>Embo-</i>	
<i>lism and Thrombosis in Brain</i>	307-314

CHAPTER XLVI.

LEUKÆMIA.

Description—Varieties—Changes in Spleen—In Lymphatic Glands—In	
Liver, etc.	315-317

CHAPTER XLVII.

THE PREPARATION AND MOUNTING OF SPECIMENS.

Examination of Specimens whilst Fresh—Hardening of Specimens—The	
Making of Sections—Imbedding—Staining—Preservation and Mount-	
ing	317-323

LIST OF WOOD ENGRAVINGS.

FIG.	PAGE
1. A Simple Cell	21
2. A Multinucleated Cell	24
3. A Vacuolated Cell. (<i>Klein</i>)	25
4. Atrophy of Adipose Tissue. (<i>Virchow</i>)	34
5. Fatty Infiltration of Connective Tissue. (<i>Rindfleisch</i>)	44
6. Fatty Infiltration of Liver Cells. (<i>Rindfleisch</i>)	47
7. Fatty Infiltration of Heart	48
8. Fatty Liver	51
9. Fatty Metamorphosis of Cells	52
10. Fatty Degeneration of Artery	55
11. Chronic White Softening of Brain	57
12. Fatty Degeneration of Muscle	59
13. Fatty Degeneration of Heart	61
14. Brown Atrophy of Heart	62
15. Colloid Degeneration of Cells. (<i>Rindfleisch.</i>)	65
16. Muscular Change in Typhoid Fever	66
17. Amyloid Degeneration of Cells. (<i>Rindfleisch.</i>)	69
18. Amyloid Liver	72
19. Amyloid Kidney	74
20. Amyloid Spleen	76
21. Corpora Amylacea. (<i>Virchow.</i>)	78
22. Calcareous Degeneration	82
23. Pigmented cells	85
24. Hæmatoidin Crystals. (<i>Virchow.</i>)	85
25. Pigmentation of Lung	89
26. Pigment Granules in Sputum	90
27. Liver Changes in Pyrexia	92
28. Heart Changes in Pyrexia	92
29. Embryonic Tissue	101
30. A Tumor Invading	105
31. Fibrous Tumor	110
32. Lipoma	112
33. Fibrous Enchondroma	113
34. Hyaline Enchondroma	114
35. Myxoma	119
36. Lymphoma	121
37. Cells from a Lymphoma	122
38. A hard Lymphoma	122

FIG.	PAGE
39. Papilloma. (<i>Rindfleisch.</i>)	129
40. Adenoma of Mamma. (<i>Rindfleisch.</i>)	132
41. Adeno-Fibroma of Mamma	133
42. Cells from Spindle-Celled Sarcoma	136
43. Small Spindle-Celled Sarcoma	138
44. Large Spindle-Celled Sarcoma. (<i>Virchow.</i>)	139
45. Melanotic Sarcoma.	140
46. Osteoid Sarcoma	141
47. Round-Celled Sarcoma	142
48. Glioma	143
49. Alveolar Sarcoma	143
50. Myeloid Sarcoma. (<i>Virchow.</i>)	144
51. Cells from Cancer	148
52. Stroma of Cancer	149
53. Scirrhus of Mamma	152
54. Ditto	153
55. Ditto	153
56. Encephaloid Cancer	154
57. Colloid Cancer. (<i>Rindfleisch.</i>)	155
58. Cells from Epithelioma	156
59. Epithelioma of Lip	157
60. Epithelioma of Tongue	158
61. Amœboid Leucocytes. (<i>v. Recklinghausen.</i>)	170
62. Pus-Corpuscles	175
63. Scrofulous Inflammation of a Bronchus	180
64. A Giant Cell.	185
65. Ditto	185
66. Giant-cell Reticulum	186
67. Adenoid-like Tissue of Tubercle	187
68. A Degenerated Tubercle	188
69. Tubercle of Pia Mater. (<i>Cornil and Ranvier.</i>)	191
70. Tuberculosis of Lymphatic Gland	192
71. Tubercular Ulcer of Intestine	194
72. Tubercle from Lung	195
73. Ditto	196
74. Ditto	197
75. Ditto	197
76. Ditto	198
77. Ditto	198
78. Bacteria from Fœtid Pus	201
79. Gumma in Liver. (<i>Cornil and Ranvier.</i>)	208
80. Gumma in Kidney	208
81. Ditto	208
82. Syphilitic Disease of Cerebral Arteries	210
83. Inflammation of Cartilage. (<i>Cornil and Ranvier.</i>)	212
84. Chronic Inflammation of Connective Tissue	216
85. A Granulating Wound. (<i>Rindfleisch.</i>)	217
86. Atheroma of Aorta	224
87. Inflammation of Aorta Valve	227

LIST OF WOOD ENGRAVINGS.

xix

FIG.	PAGE
88. Inflammation of Mitral Valve	227
89. Endocarditis due to Friction	227
90. An Inflammatory Granulation from Mitral Valve. (<i>Rindfleisch.</i>)	228
91. Acute Myocarditis	230
92. Fibroid Induration of Heart	231
93. Ditto	231
94. Chronic Inflammation of a Lymphatic Gland	233
95. Typhoid Swelling of Peyer's Patches	235
96. Typhoid Ulcer of Intestine	235
97. Ditto	236
98. Catarrhal Inflammation of Conjunctiva. (<i>Rindfleisch.</i>)	237
99. Inflamed Epiploon of Rabbit. (<i>Cornil and Ranvier.</i>)	242
100. Inflammation of Pleura. (<i>Rindfleisch.</i>)	243
101. Cirrhosis of Liver	246
102. Ditto	246
103. Surgical Kidney	250
104. Tubal Nephritis	252
105. Ditto	253
106. Interstitial Nephritis	256
107. Ditto	257
108. Ditto	258
109. Arteries of Kidney in Chronic Bright's Disease	259
110. Sclerosis of Spinal Cord	262
111. Croupous Pneumonia—Red Hepatization	263
112. Croupous Pneumonia—Gray Hepatization	265
113. Broncho-Pneumonia—Inhaled Secretion	270
114. Catarrhal Pneumonia	270
115. Pulmonary Fibrosis from Chronic Bronchitis	272
116. Interstitial Pneumonia	275
117. Ditto	276
118. Ditto	276
119. Acute Phthisis	279
120. Ditto	279
121. Ditto	280
122. Chronic Phthisis	281
123. Ditto	283
124. Acute Phthisis	289
125. Nutmeg Liver	296
126. Brown Induration of Lung	297
127. Organization of Thrombus. (<i>Rindfleisch.</i>)	301
128. Ditto (<i>O. Weber</i>)	301
129. Thrombus in Vein. (<i>Virchow.</i>)	308
130. Impacted Embolus. (<i>Virchow.</i>)	309
131. Hemorrhage Infarct. (<i>O. Weber.</i>)	312
132. Leukæmic Blood	316

PATHOLOGY

AND

MORBID ANATOMY.

INTRODUCTION.

PATHOLOGY treats of the origin, nature, course, and causes of those changes in the body which constitute disease ; *Morbid Anatomy* or *Morbid Histology*, of the actual alterations in the tissues which the disease has produced. The former is therefore comparable with Physiology, the latter with the Histology of the normal tissues.

By *disease* is understood some deviation from the state of health ; a deviation consisting for the most part in an alteration in the functions, properties, or structure of some tissue or organ, owing to which its office in the economy is no longer performed in accordance with the normal standard.

Diseases are commonly divided into two classes—*organic* and *functional*. The former comprise all those which are attended by structural change ; the latter those in which no such change is discoverable. Whether it be possible for the function of an organ or tissue to be abnormally performed quite irrespective of any alteration in its structure admits of some doubt. At all events, as our methods of minute investigation improve, and our knowledge of morbid histology increases, the class of functional diseases grows less ; and although there still remain a large number of diseases in which we are unable to recognize any alteration of structure, and which must therefore still be described as functional, it is probable that all disease will ultimately be found to be attended by more or less material change.

As *health* is itself merely a relative term and implies no definite performance of the processes of life, so *disease* is equally indefinite ;

it cannot be separated from health by any well-defined boundary, the one passing by insensible gradations into the other.

Disease is thus, in most cases, an abnormal performance of those processes which constitute life, and a knowledge of these processes must therefore necessarily precede the study of pathology. Life comprises the formation and maintenance of the tissues and the exhibition of their various functions. Such formation and maintenance, which may be included under the general term of *Nutrition*, consist in the continuous supply of new material, the separation of this from the blood and its appropriation by the tissues, together with the removal of the products of their waste. *Function* is the special manifestation of the life of the part, as distinct from its growth and the maintenance of its structure; in the secreting cell, consisting in the alteration of the substances abstracted from the blood to fulfil some special purpose in the economy; in nerve, in the transmission of impulses of motion and sensation, etc. The performance of function is obviously dependent upon the state of nutrition. When both of these are normal the condition is one of *health*, when abnormal one of *disease*.

As in health the nutrition is principally dependent upon the solid tissues, these abstracting, appropriating, and altering the material which is supplied to them by the blood; so in most forms of disease it is these tissues which play the more important part, and alterations in them are amongst the most frequent of the morbid processes.

The supply and composition of the blood must obviously at the same time constitute most important factors in disease. Owing to the intimate relation which subsists between the blood and the solid tissues, any alteration in the composition of the former must exercise more or less influence upon the nutritive processes. In regarding the blood, however, as an element in the causation of disease, it must be borne in mind that this fluid is in a state of constant dependence upon other parts; its component elements are derived from external sources and undergo continual change, and although possibly under certain circumstances it may become altered by virtue of changes in the nutritive activity of its cellular elements, alterations in its constitution must in most cases result either from some change in the process of its formation, as from the ingestion of improper or insufficient food, mal-assimilation, or disease of the lymphatic structures; from changes in the secretory or excretory processes; or from the introduction of foreign substances, derived from extraneous sources. Whilst, therefore, alter-

ations in the composition of the blood may be important agents in the production of diseased processes in other tissues, such alterations are in most cases secondary to some abnormality in its formation or depuration, or to its contamination by substances derived from without. Owing to the facility with which it becomes the recipient of foreign substances, it is often the first tissue to become altered, as, for example, in many of those diseases which owe their origin to the introduction of minute organisms; and it not infrequently constitutes the principal seat of the diseased process.

The blood may also play an important part in disease, owing to an abnormal activity of its white corpuscles and their migration through the walls of the bloodvessels into the surrounding tissues. This occurs in the process of inflammation, and probably also in the development of many of the new formations.

Lastly, the influence of the nervous system must be taken into account in considering abnormal conditions of nutrition and function. This, by regulating the circulation and supply of blood, must to a certain extent control tissue-change. The results of experimental and clinical observation render it exceedingly probable that many nerves also contain fibres which exercise a direct influence upon nutrition, independently of that which is due to their vaso-motor filaments. In support of this view may be adduced the well-known observations of Ludwig and Heidenhain on the influence of the chorda tympani upon the secretion of the submaxillary gland, which prove that the increased secretion which results from the stimulation of this nerve is due to an increased activity of the secreting elements of the gland and not to changes in the bloodvessels. Recent histological researches, also, which are tending to show that nerve-fibres terminate in the ultimate elements of nearly all tissues, and the fact that very rapid atrophy of muscles often follows certain lesions of the nervous centres and many injuries of the nerve-trunks, render it exceedingly probable that nutrition is more or less under the *direct* influence of the nervous system.

The *first* part of this work will be devoted exclusively to the consideration of morbid processes which are characterized mainly by alterations in nutrition; the *second*, to those in which an altered nutrition is associated with certain changes in the bloodvessels and circulation (inflammation); and the *third*, to changes in the blood and circulation alone. In considering the several morbid processes, the general pathology of each process will first be described, and subsequently the same process as it occurs in the different organs and tissues.

CHAPTER I.

THE CELL.

As the most important element in nutrition, both in health and disease, is the activity of the tissues themselves—the supply of nutritive material, although an essential, being merely a *passive* part of the process—it becomes necessary to consider, somewhat minutely, those parts of the tissues in which this activity resides.

Ever since Schwann discovered the cellular nature of animals, and established the analogy between animal and vegetable cells, there has been a gradually increasing conviction amongst physiologists, which has now become an universally accepted physiological and pathological doctrine, that the *cell* is the seat of nutrition and function; and further, that *each individual cell* is itself an independent organism, endowed with those properties, and capable of exhibiting those active changes which are characteristic of life. Every organized part of the body is either cellular or is derived from cells, and the cells themselves originate from pre-existing cells, and under no circumstances do they originate *de novo*.

Whilst therefore the whole body is made up of cells, or of substances derived from cells, and the cell is itself the ultimate morphological element which is capable of exhibiting manifestations of life, it must be borne in mind that in a complex organism, the phenomena of life are the result of the continued activity of innumerable cells, many of which possess distinct and peculiar functions, and that by their combination they become endowed with new powers, and exhibit new forces, so that although each individual unit possesses an independent activity, it is in a state of constant dependence upon others with which it is more or less intimately associated.

CONSTITUTION OF CELLS.—When the analogy was established between the animal and vegetable cell, the former was held to be constructed in all cases upon the same principle as the latter, and to consist of a *cell-wall*, inclosing a cavity in which were contained a *nucleus* and *fluid contents* (Fig. 1). This was the idea of the cell held by Schwann and Remak, and supported especially by Virchow,

who maintained that these three constituents were essential to its vitality and existence.

During recent years, however, this definition has been modified. The existence of a cell-wall was in many cases not evident. In embryonic cells, in those of many rapidly growing new formations, and in the cells of blood, pus, and mucus, no limiting membrane could be demonstrated. This led to a new definition of the cell by Leydig and Max Schultze, who held that a little mass of matter inclosing a nucleus was all that was necessary for its constitution. The latter of these physiologists not only rejected the cell-wall as an essential constituent, but established the identity of the mass of matter (cell-contents) with animal sarcode—a contractile substance existing in the lower animals—and showed that, like it, it was endowed with the power of spontaneous movement. This substance he called *protoplasm*. He further pointed out, that the existence of a distinct cell-wall was the result of a retrograde process taking place in the outer layers of the protoplasm, and that the latter was the real seat of the activity of the cell. These views closely correspond with those held by Dr. Beale in this country.¹

The definition of a cell has been still further modified by Brücke, Stricker, and others, who consider that the existence of a nucleus is not essential to its constitution. This opinion is principally based upon the fact, that in the cryptogamia, and in some of the lowest animal forms, cells occur in which no nucleus is visible.

It would thus appear probable that a simple mass of protoplasm may, in some exceptional cases, be all that is necessary to constitute a cell—*i. e.*, an elementary organism, capable of exhibiting independently all the phenomena of life; but that the nucleus is an exceedingly constant and almost invariable constituent. The cell-wall is much less constant, and being the result of a retrograde change in the outer layers of the protoplasm, it must be regarded, in point of vitality, as inferior to the rest of the cell.

Protoplasm itself is an unstable albuminoid compound, which is insoluble in water and coagulates at death. As usually met with, it

FIG. 1.



Cells from a cancer. Showing cell-wall, cell-contents, nuclei, and nucleoli. The nuclei dividing.

¹ Dr. Beale calls the protoplasm, *germinal matter* or *bioplasm*; the cell-wall, *formed material*.

is a homogeneous structureless material, of a soft and viscid consistence. In consistence, however, it is subject to variations, being sometimes perfectly fluid, at others more or less solid and gelatinous. In old cells it often becomes transformed, by the loss of water, into a more solid albuminoid substance—*keratin*. This occurs in the epidermis and nail. Protoplasm may also become gradually converted into other modifications of the protein group—into mucin, globulin, hæmoglobin, etc. The cell-wall, when it exists, is of much firmer consistence than the protoplasm. In some cells the protoplasm constitutes but a small proportion of the body of the cell, other substances, which are either the result of its metamorphosis, or have been taken up from without, being associated with it. Thus fat is met with in the cells of adipose tissue, and of the liver. (See Fig. 6.) Pigment, calcareous particles, pepsine, etc., are also met with in cells.

The *nucleus* is more constant both in size and form than the cell. It is usually spherical or oval in shape, and often contains one or more minute round or angular bodies, termed *nucleoli*. It offers a greater resistance to chemical reagents than the other constituents of the cell, and in disease often remains after these have been destroyed. It is also stained more deeply by carmine.

The original form of the nucleus is vesicular. In the earliest cells of embryonic tissue it possesses a delicate membrane inclosing a nucleolus and fluid contents, thus resembling in its structure the cell. Subsequently, however, it loses its vesicular character, and as usually met with it is a solid perfectly homogeneous, or faintly granular body, in which the nucleolus is still visible. The recognition of the nucleus is not always possible, owing to its presence being obscured by fat, pigment, or other substances contained within the cell. (See Fig. 27.) In some cells the nuclei gradually disappear. The colored blood-cells and the cells of the superficial layers of the epidermis are examples of cells in which the original nucleus has become lost. Lastly, it must be mentioned, that several nuclei may be contained within the same cell.

PHYSIOLOGY OF CELLS.—The cell, as already stated, is capable of absorbing and transforming matter, or excretion, and of growth. It is also endowed with the power of changing its form, of cohering with other cells, of undergoing more or less active movement, and of reproduction. The question now arises as to what part is played by its respective constituents—whether the cell-wall, the body of the cell (protoplasm), and the nucleus have different offices.

The cell-wall, being the result of a retrogressive change in the protoplasm, cannot be regarded as taking any part in the life of the cell, the activity of which is much diminished by its existence, as is also its power of reproducing itself by simple division. It is in old cells that a cell-wall is most frequently met with, in those newly formed it is entirely wanting.

The nucleus has usually been looked upon as the seat of the nutrition, as distinct from the specific functions of cells, and has been supposed to play an important part in their multiplication and reproduction. The fact that when a cell divides, the division usually commences in the nucleus, and only subsequently takes place in the rest of the cell, would appear to favor this view; as would also the great uniformity of the nucleus both in size and form, whatever be the functional nature of the cell. It must be borne in mind, however, that non-nucleated cells may multiply, and that nucleated cells have been observed to divide, the nucleus itself taking no part in the process.

Whatever be the part played by the nucleus, there can be no doubt that the protoplasm is the most important factor of the cell, and it may itself be the only constituent. The spontaneous movements, alterations in form, and migratory powers characteristic of young cells, are due to the protoplasm. Such movements are observed in the cells of the embryo, in lymphoid and young epithelial cells, in some of the cells of connective tissue, and in white blood and pus cells.

The protoplasm, as already stated, may be the sole seat of the nutritive and formative power of the cell. It would appear, however, probable that it is more especially concerned in the performance of function, and that the specific functional peculiarities of cells are dependent rather upon it than upon their other constituents. The volume and consistence of the protoplasm vary in different cells, and in the same cell, at different times and under different circumstances. It is apparently capable of imbibing and giving up fluids, at the same time undergoing corresponding alterations in volume. These considerations render it probable that it is the seat of the selective power of the cell, and of those other properties which represent its specific functions.

GENESIS OF CELLS.—The proposition of Virchow, that every cell originates directly from a pre-existing cell, forms the basis of the pathology of the present day. To Remak, however, must be ascribed the merit of having first established the cellular origin of the tissues.

The multiplication of cells may take place in three ways—by *simple*

division, by *gemmation*, and by *endogenous growth*. In the first two methods the cell breaks up into fragments; in the last, new cells originate within the parent cell. The process is obviously associated with growth and increase of the protoplasm.

FIG. 2.



A M. Vinucleated Cell. From the lung in a case of chronic phthisis. Showing the large number of nuclei with bright nucleoli. $\times 40$.

The multiplication by simple division is the most frequent method. The cell divides and forms two cells, and each of these again divides and forms two more, and so on. In nucleated cells the nucleus as a rule divides first. The nucleus, however, may divide and multiply within the cell without any division of the cell taking place. If the nuclei multiply within the cell, and the protoplasm continuously increase without subse-


quent division of the cell taking place, large, many-nucleated, irregular-shaped masses of protoplasm are produced. These are the giant or myeloid cells, which are met with in the medulla of young bone, in some new formations, and in certain inflammatory growths. (Fig. 2.) (See also "Myeloid Sarcoma," Fig. 50.) The existence of a dense cell-wall interferes with the process of multiplication by simple division.

By endogenous multiplication is understood the development of cells within pre-existing cells. The multiplication of the cells of cartilage, such as occurs in the growth of bone and in the process of inflammation, has been adduced as an example of this mode of cell formation. Here, however, we have simply the division of the cartilage cell within its capsule, and the process is precisely similar to multiplication by simple division. (See Fig. 83 *d*.) The mode of cell formation which must be regarded as strictly endogenous, is what is now commonly known as *Vacuolation*. This was described some time ago by Virchow in his "Cellular Pathology;" and owing to the more recent researches of Klein and others, it now occupies an important place in the history of cell development. The process consists in the formation of a vacuole in the body of the cell. This vacuole may gradually increase in size until it occupies nearly the whole of the cell, being merely surrounded by a thin layer of protoplasm, in which may often

be seen the displaced nucleus (Fig. 3, *b*). Within this vacuole one or more new elements are formed. These either originate from the nucleus of the cell, or they are produced from the protoplasm which forms the wall of the vacuole. In the latter case, according to Dr. Klein, buds grow out from the protoplasmic wall towards the interior of the vacuole, and these becoming detached form new cells.¹ A vacuolated cell with an endogenous brood is shown in the accompanying drawing (Fig. 3), for which I am indebted to the observer just quoted. It must, however, be remembered, in forming any conclusion as to the origin of small cells within a larger cell, that there exists the possibility that these may have entered from without.

The endogenous mode of cell formation is not that by which tissues regenerate themselves. This is usually effected by simple division. Elements which have an endogenous origin have commonly a destiny different from that of the parent cell.

In multiplication by gemmation, a small portion of the protoplasm projects from the cell and becomes detached by constriction to its base, thus forming a new cell. This is much less frequent than the two former processes.²



A Vacuolated Cell. a. Protoplasm of cell constituting wall of the vacuole. b. Displaced nuclei. c. The vacuole. d. Drop of fat. e. Endogenous brood. (Klein.)

¹ "The Anatomy of the Lymphatic System," vol. i. pp. 55-60; and Klein on "Endothelial vesicles of growing bloodvessels in the embryo chick." "Sitzungsberichte der Wiener k. Academie d. Wissenschaften," Part for March, 1871.

² The author wishes it to be distinctly understood, that in the subsequent pages the term "cell" is employed to designate the most elementary factor of the tissues which is capable of exhibiting the phenomena characteristic of an independent organism, quite irrespectively of the existence of a cell-wall, or even of a nucleus.

CHAPTER II.

NUTRITION ARRESTED.

THE absolute and permanent arrest of nutrition constitutes local or systemic death. Under this head are included the three following processes:—

1. *General or Systemic Death*.—In this, the arrest is general; the nutritive processes cease throughout the whole body.

2. *Gangrene or Necrosis*.—The arrest is local, and confined to a particular part, which, when dead, retains to a greater or less extent its external form and anatomical characters.

3. *Molecular Death or Necrobiosis*.—The arrest is also local, but usually results from a previous gradual impairment of the nutritive processes, which ultimately entirely cease in some of the histological elements. The dead part is a granular débris in which all trace of the original structure is lost. This will be considered under “Atrophy and degeneration.”

Gangrene or Necrosis.

Gangrene or Necrosis is the complete and permanent arrest of nutrition in a part—occurring for the most part more or less suddenly—which, when dead, retains to a greater or less extent its external form and anatomical structure. The series of processes by which this is brought about is called *Mortification*, the dead tissue—a *Sphacelus* or *Slough*. In bone the process is called *Necrosis*, the result—a *Sequestrum*.

The arrest of nutrition is followed by the complete cessation of all the evidence of life both functional and physical; and the part thus removed from the influence of the “vital forces” undergoes those chemical and physical changes which are common to inanimate organic matter. The extent of these changes and the rapidity with which they take place, will vary considerably according to the nature of the surrounding media, and to the amount of blood and fluids contained in the part. When the gangrene is due to, or associated with, venous

obstruction, and the return of blood and absorption of fluids are thus interfered with, the decomposition of the dead tissue, if exposed to the air, proceeds rapidly. In a limb, for example, the coldness and the paralysis of motion and sensation, which are amongst the earliest signs of the cessation of life, are followed by the transudation of the liquor sanguinis from the bloodvessels, and, evaporation being to a great extent hindered by the epidermis, the transuded and accumulated liquids often form large bullæ on the surface. As decomposition proceeds, gases are generated in the part—principally sulphuretted hydrogen, ammonia, nitrogen, and carbonic acid. These give rise to the emphysematous crackling which is so often associated with the gangrenous processes. The tissues at the same time undergo a process of softening or liquefaction, the limb becomes exceedingly offensive, and, owing to alterations in the transuded hæmoglobin, changes from a reddish color to a brownish or greenish-black:—this is *Moist Gangrene*. In those cases, on the other hand, in which the gangrene is associated with obstruction of the main arteries, as in senile gangrene, in the gangrene due to embolism, and in that induced by ergot of rye, there is no accumulation of blood or fluids in the part, so that decomposition proceeds much less rapidly than in the preceding, and the dead issues may gradually dry up and become converted into a black shrunken mass, which undergoes but little further change:—this constitutes *Dry Gangrene* or *Mummification*.

The characters of the dead part vary with its structure, its vascularity, the cause of the gangrene, the acuteness of the process, and the possibility of the access of atmospheric air. The more vascular the tissue, the softer its structure, and the more it is exposed to the atmosphere, the more rapidly and completely does it undergo decomposition. Bone, cartilage, and tendons, which are firm, hard tissues, containing comparatively but few vessels, undergo very little alteration in structure and form; whereas softer parts are much more rapidly and completely destroyed.

The occurrence of decomposition manifests itself in the first place in the blood contained in the part: this fluid undergoes the earliest and most rapid change. The hæmoglobin escapes from the red corpuscles, partly by exudation, and partly by the destruction of the corpuscles themselves, and dissolved in the liquor sanguinis permeates the surrounding tissues. The corpuscles are ultimately completely annihilated, nothing remaining but a few minute granules.

The staining of the tissues with hæmoglobin is commonly known as

post-mortem staining, and the appearances it presents are very characteristic. All the tissues may be more or less affected, but the lining membrane of the heart and large bloodvessels, being in immediate contact with the blood after death, are naturally more so than other parts. The staining is of an uniform pinkish-red color, thus differing from the punctiform and stratiform redness of hyperæmia, from which it must be carefully distinguished. The amount of staining is in proportion to the rapidity with which decomposition has taken place, and to the amount of blood contained in the part at the time of death.

RIGOR MORTIS.—In muscle the arrest of nutrition is accompanied by a state of rigidity, known as the *Rigor Mortis*. This is a peculiar condition of the muscles observed in almost all bodies after death, in which they become firm and somewhat shortened, as though in a state of chronic contraction. It comes on as soon as the muscles have lost their irritability, *i. e.*, their capability of responding to artificial stimulation; in other words, as soon as the nutritive processes have completely ceased. The time of its appearance will therefore depend upon the state of nutrition of the muscles at the time of death; the more healthy and vigorous this is, the longer it is before the nutritive processes completely cease, and consequently the longer it is before the rigor mortis supervenes. The length of its duration and its intensity are in direct proportion to the lateness of its appearance. In people, for example, who are in perfect health, and die suddenly, as from accident, the rigor mortis does not usually come on until from ten to twenty-four hours after death; it is very marked, and often lasts two or three days. In those, on the other hand, who die from some exhausting disease, as from chronic phthisis or the adynamic fevers, in which the nutrition of the muscles becomes much more impaired, the rigor mortis appears very soon, sometimes as early as ten minutes after death; it is very slight and may pass off in less than an hour. It has been said that in cases of death from poisoning by carbonic acid and sulphuretted hydrogen, from lightning, and from some of the severer forms of the adynamic fevers, the rigor mortis is entirely absent. It is doubtful, however, if this is the case, as the rigor mortis has probably escaped observation, owing to its early super-vention and rapid disappearance. As soon as the rigor mortis has passed off, decomposition of the muscular tissue commences.

The rigor mortis occurs not only as the result of systemic and local death, but it may also be induced artificially by temporarily arresting the nutrition of the muscle. If the supply of blood to a

muscle be cut off by the application of a ligature to the artery supplying it, it quickly passes into a condition which is indistinguishable from the rigor mortis. If the ligature be removed sufficiently early, this disappears, and the vitality of the muscle is restored.

With regard to the nature of the change—it was formerly supposed to be a spontaneous contraction, the last act of vitality on the part of the muscle. More recently, however, Kühne and others have shown that it is really owing to the coagulation of the albuminous substance of the muscle—myosin. The myosin, fluid during life, coagulates when nutrition has ceased, the coagulation being attended by the liberation of a free acid. Thus is produced the firmness, hardness, and opacity of the muscle, which disappear as soon as disintegration and decomposition commence. The transverse striation of the fibres then becomes indistinct, and gives place to irregular rows of granules, and fat molecules, the muscle softens, its sarcolemma is destroyed, and ultimately nothing remains but a soft structureless débris.

A similar coagulation takes place in the cells of plain involuntary muscle, and here also a rigor mortis occurs. In the cells of all other tissues, a coagulation of the protoplasm occurs on the cessation of the nutritive processes; the cells become cloudy and granular, and then break up into molecules of various sizes.

Respecting the disintegration of other tissues: The cells of adipose tissue rapidly diminish in size, owing to the escape of the fluid fat, which diffuses itself throughout the surrounding structures. The fibres of connective tissue swell up, become opaque, and ultimately liquefy. In nerve-fibres, the white substance of Schwann coagulates and collects into small drops (myelin) within the neurilemma. Cartilage and bone resist the necrotizing process longer than any of the tissues and are the least altered by it.

The termination of the gangrenous process varies. It may, after involving a greater or less extent of tissue, become arrested, and a “line of demarcation” form between the dead and living parts (*Circumscribed Gangrene*); or the process may continue to extend without any such attempt at recovery (*Diffuse Gangrene*). When the process becomes arrested, the dead tissue—the sphacelus or slough—acts as a foreign body, and as such sets up inflammatory changes in the adjacent living structures, and by this means it is ultimately removed or becomes encapsuled. The tissues immediately surrounding the necrosed parts are thus in a state of inflammation, as is evidenced in external structures by their swelled condition, red color, and high

temperature. As the gangrenous process ceases, the necrosed fragment becomes limited by this line of inflamed tissue, which constitutes the "*line of demarcation*" between the dead and living parts. Along this line a process of ulceration and suppuration takes place, and by means of this the dead mass is gradually separated from the surrounding structures. The ultimate termination of the process depends principally upon the situation of the affected part—if this be superficial, the slough is thrown off, as in external parts, the intestines, the pharynx, etc., an ulcerated surface being left. If the dead mass is deeply seated, its removal becomes possible only by the extension of the necrotizing process to the surface, as is exemplified by the spontaneous removal of necrosed bone through fistulous openings in the soft parts. In other cases the inflammatory process which takes place in the tissues surrounding the dead part is less intense, and the formation of pus is less abundant, and is soon followed by that of connective tissue, a layer of which is ultimately formed around the necrosed mass by which it becomes *encapsuled*. This occurs especially in internal parts. Examples of it are furnished by foreign bodies, hæmorrhagic infarcts, accumulated epithelial products, portions of necrosed bone, and a fœtus in the abdominal cavity, all of which may thus become surrounded by a layer of connective tissue. The part when thus encapsuled is usually rendered inert, and no longer acts as an irritant to the tissues in which it lies; it undergoes a gradual process of absorption and drying up, and often becomes calcified.

CAUSES.—The causes of gangrene may be divided into those which interfere with the supply of nutritive material, and those which directly destroy the vitality of the histological elements, so that they are no longer able to perform their functions.

A. The supply of nutritive material may be interfered with by:—

1. *Obstruction in the Arteries*.—This is a common cause of gangrene. The obstruction may be caused by a ligature, by compression of the vessel, by solution of its continuity, by thrombosis or embolism, and by disease of the arterial coats. If the obstruction be complete and a collateral circulation cannot be established, death of the part quickly ensues. Obstruction to the arteries alone, the return of blood by the veins not being interfered with, usually produces the dry form of gangrene.

2. *Obstruction in the Capillaries*.—Obstruction here is usually the result of pressure upon or stretching of the vessels. This may take place from the accumulation of inflammatory products, hæmorrhage, or

from the pressure exercised by new growths. The resulting obstruction to the capillary circulation causes the death of the immediately adjacent tissues. As examples of gangrene from this cause may be mentioned the necrosis of the superficial layers of the bone which so frequently results from periostitis, owing to the compression of the capillaries between the bone and the periosteum; also the sloughing of the mucous membranes in diphtheria, and of the skin over a quickly-growing abscess.

3. *Obstruction in the Veins.*—Obstruction to the return of blood by the veins must be so complete, in order to arrest nutrition, that it is in itself rarely a cause of gangrene. It is when associated with obstruction in the arteries that it constitutes an important agent in producing this result. This combination of venous and arterial obstruction is seen in a strangulated hernia, in the invagination of a portion of the intestine, in the constriction of a part by a tight bandage, and in contusions and lacerations of the soft parts in which both arteries and veins are injured or become strangulated by the tension of the tissues which subsequently ensues. In all cases in which gangrene is associated with venous obstruction it is of the *moist* variety.

4. *Diminished Cardiac Power.*—This is never independently a cause of gangrene. In cases, however, of excessive general debility or disease of the cardiac substance, the consequent diminution in the contractile power of the heart materially aids the foregoing causes in producing a fatal blood-stasis. The arrest of the circulation in “senile gangrene,” and that which so often occurs in the tissues of the back in adynamic fevers and in chronic exhausting diseases, is in part the result of diminished cardiac power. This arrest in the last-named conditions is usually determined by some injurious stimulation of the tissue—in other words, it is a part of an inflammatory process.

5. *Inflammation.*—The effect of the inflammatory process is to impede or arrest the circulation, and to impair the vitality of the affected part, and the intensity of the process may be so great as to permanently arrest the circulation and cause gangrene. Gangrene from inflammation has been called *inflammatory*, *acute*, or *hot* gangrene. It is always of the moist variety, and is characterized by the heat, swelling, tension, and redness of the affected part, which quickly becomes flaccid, cold, of a greenish-black color, and exhales an offensive odor. Certain inflammations have a special tendency to terminate in gangrene, as diphtheria, erysipelas, carbuncle, and “hospital gangrene.” In these conditions the gangrenous tendency is probably

due to some alteration of the blood, possibly to the presence of minute organisms. In all cases, the more impaired the nutrition of the part which becomes the seat of an inflammatory process, the more likely is this to cause its death.

B. Destruction of the vitality of the histological elements may be caused by:—

1. *Mechanical Agencies*.—Under this head are included external violence, heat, cold, and corrosive substances, all of which, by directly and completely destroying vitality, may be causes of gangrene. In most cases, however, this result is preceded by more or less inflammation.

2. *Poisons introduced into the Circulation*.—Phosphorus and ergot of rye are the most important of these. The necrosis of the jaw which results from phosphorus appears, however, to be owing to the direct contact of the phosphorus with the bone, and the gangrene of the extremities which sometimes follows the long-continued ingestion of ergot, is probably due to that contraction of the small arteries which this substance produces.

SENILE GANGRENE.—This is a form of mortification which affects especially the lower extremities of old people, and is the result of several of those conditions which have already been enumerated as causes of gangrene.

The most important element in the production of senile gangrene consists in the occurrence of atheromatous or calcareous changes in the arteries of the limb, in consequence of which the circulation in it becomes impeded and its vitality impaired. This is evidenced by the coldness of the feet, the cramps, and the other abnormal sensations which are so often experienced by the patient some time before the gangrene sets in. This tendency to local stagnation of the circulation is usually materially increased by the simultaneous atrophy or degeneration of the muscular substance of the heart itself. The combined effect of the diminished *vis à tergo* and of the arterial degeneration may, in some cases, be alone sufficient to cause arrest of the circulation and the formation of thrombi in the vessels of the limb, and thus to cause gangrene. The supervention of the gangrene, however, is usually determined by some injurious stimulation of the tissue, as a slight abrasion of the foot, a bruise, injury to a corn, or excess of heat or cold, which sets up inflammation in the already weakened part, and thus by still further obstructing the circulation in it, and impairing its vitality, causes its death. Senile gangrene, being associated with arterial obstruction, is usually of the *dry* variety.

CHAPTER III.

NUTRITION IMPAIRED.

It has been seen in the preceding chapter that the absolute arrest of nutrition is followed by the complete cessation of all manifestations of vitality and function, constituting local or systemic death. Those conditions must now be considered in which the interference with nutrition, for the most part, falls short of absolute arrest, and in which, although vitality is impaired, death is only an occasional sequence. Such conditions are comprised under "Atrophy," and "Degeneration."

Atrophy.


Atrophy is a diminution in the amount of a tissue, owing either to diminution in the *size*, or diminution in the *number*, of the histological elements of which it is composed. It is attended by loss of weight, and impairment of function.

When the elements are diminished in *size*, it is called *Simple Atrophy*; when the *number* is diminished, it is called *Numerical Atrophy*. These two varieties are often associated, being different stages of the same process: simple atrophy may, however, exist without numerical atrophy, but numerical never exists without simple.

I. SIMPLE ATROPHY.—Simple diminution in the size of the elements of a tissue is by far the most common condition met with in atrophy. It is well exemplified by what takes place in ordinary emaciation, in which the fat gradually disappears from the subcutaneous adipose tissue. Adipose tissue is merely common connective tissue, many cells of which are distended with fat. When a person emaciates, the fat is gradually removed from the cells, so that they diminish in size, and the fat which completely filled the cell may be reduced to a few isolated drops: the cell-wall and nucleus at the same time often become distinctly visible (Fig. 4). Here there is no destruction of the cells, no diminution in their number, but simply the removal of some of their contents. As the fat is removed from the cells it is often partially replaced by a serous fluid.

This diminution in the size of the elements may take place in any tissue. The cells of all granular organs may thus become atrophied, and so produce a diminution in the size of the whole organ; this is constantly met with in the liver, kidneys, mammary gland, spleen,

FIG. 4.



Adipose tissue. A. Normal. B. Atrophic, from a case of phthisis. a. A single fat-cell with cell-wall, nucleus, and drop of fat. $\times 300$ (Virchow).

testicles, lymphatic glands, and other parts. Muscular tissue in the same way atrophies by the diminution in the size of its primitive fasciculi; this is seen in the heart and in the voluntary muscles. In all these cases the elements remain almost unchanged, and hence all that is necessary for the restitution of the tissue is an increase in their nutritive activity, and the assimilation of more material.

II. NUMERICAL ATROPHY.—This is an advanced stage of the former process; the elements are not only diminished in size, but some of them have actually perished and ceased to exist as vital agents.

This destruction of histological elements which occurs both in atrophy and in some forms of degeneration—and is called by Virchow, *Necrobiosis*—must be distinguished from the death of circumscribed portions of tissue which constitutes gangrene and necrosis. The two processes resemble one another in so far as death is common to both of them. In *Necrobiosis*, however, the change is a molecular one; there is a gradual exhaustion of vital power, a molecular disintegration and destruction of elements, so that at the termination of the process all that remains is a granular débris, in which but little or no trace of the former structure of the part can be discovered. The death and desquamation of the superficial layers of the epidermis is a well-known example of this molecular change. *Gangrene*, on the other hand, affects circumscribed tracts of tissue, and is the result of some sudden arrest of nutrition, as distinguished from the gradual exhaustion of vital power. Death is more suddenly induced, and a necrosed mass remains at the termination of the process in such a condition that the

structure of the part can usually be recognized. Numerical atrophy is thus of much graver import than that in which the elements continue to exist as such. In it, restitution is only possible by the production of new elements, whereas in simple atrophy, repair can be effected without new formation.

Atrophy may be *general*—affecting to a greater or less extent all the organs and tissues of the body, or it may be *partial* and limited to particular parts. General atrophy is usually *simple*, and is rarely accompanied by destruction of elements. It affects in the first place the subcutaneous adipose tissue, then the adipose tissue in other situations, as that surrounding the viscera and in the omentum, then the muscles and glandular organs, and lastly the nervous and osseous structures.

Although atrophy in its strict signification consists simply in a diminution in the size or in the number of the component elements of a tissue, it is rarely a perfectly simple process, but is usually associated with more or less *fatty degeneration*. This is owing to the fact that whenever the nutrition of a part is so much interfered with as to cause it to atrophy, it is very prone to undergo fatty changes; and it will be seen, when speaking of “fatty degeneration,” that this process owes its origin to causes similar to those which produce atrophy itself.

CAUSES.—In speaking of the causes of atrophy, it will be necessary to distinguish between those which act upon the tissues generally, and those which have merely a local influence.

General Atrophy may be caused by:—

1. *Deficient supply of Nutritive Material*.—Whatever interferes with the supply of nutritive material to the tissues will be followed by their atrophy. Deficient supply of food; obstruction to the passage of the food into the stomach or intestines, as in stricture of the œsophagus or pylorus; the mal-assimilation which results from the various conditions giving rise to dyspepsia; interference with the absorption of the chyle, from obstruction of the thoracic duct, or disease of the mesenteric glands constituting the so-called “*tabes mesenterica*,” may all in this manner be causes of general atrophy.

2. *Excessive Waste*.—All those conditions which are attended by the loss of large quantities of nutritive material, may be causes of general atrophy. Such conditions are furnished by continuous hemorrhages, profuse and long-continued suppuration such as often occurs in caries and empyema, diarrhœa, and the excretion of large quantities of albumen or sugar as in Bright’s disease or diabetes. The waste

resulting from the increased tissue-change which accompanies acute febrile diseases, must also be included under this head.

3. *Impaired Nutritive Activity*.—This constitutes an important element in the production of the atrophy of old age,—*senile atrophy*. As life advances, the vitality of the elements gradually diminishes, their ability to separate nutritive material from the blood and to assimilate it for their own maintenance becomes less and less, and hence they gradually atrophy, and ultimately all manifestations of their vitality cease.

Although general atrophy may thus be referred to one of the foregoing causes, it is rarely a simple process, but usually depends upon the combined influence of two or more of them. The atrophy associated with pulmonary phthisis, for example, results partly from the loss of nutritive material in the profuse expectoration and diarrhoea, partly from the deficient supply consequent upon the imperfect oxidation of the blood and upon the interference with assimilation which is so often caused by structural changes in the stomach and intestines, and partly from the increased tissue-change. In senile atrophy, again, in addition to the general diminution of nutritive activity, there is frequently some condition of the digestive organs interfering with assimilation which materially aids in producing the ultimate result. The atrophy which accompanies the acute febrile diseases is by no means a simple process, increased tissue-change, loss of appetite, and interference with assimilation being all component parts of it.

Partial Atrophy may be caused by:—

1. *Imperfect supply of Blood*.—The effect of interfering with the supply of blood to a part will depend upon the extent of the interference. If it be entirely cut off the part will die (see “Gangrene”), if merely diminished it will atrophy.

Diminished supply of arterial blood is a common cause of atrophy, and may be brought about in various ways. The nutrient vessels may be obstructed by pressure exercised upon them within or without the organ to which they are distributed. In cirrhosis of the liver, the pressure exercised by the new growth of inter-lobular tissue causes atrophy of the secreting structures. In other cases the supply of blood is diminished by interference with the circulation at some distance from the part, as by the pressure of a tumor upon the artery leading to it. The atrophy of the proximal end of the shaft of a bone after fracture above the point of entrance of its nutrient artery, is due to the same cause.

The atrophies which result from pressure exercised directly upon the part itself, are also owing to the consequent interference with nutritive supply. Atrophy of the sternum from the pressure of an aneurism, atrophy of the kidney from the pressure of retained secretion—as in stricture of the urethra, enlarged prostate, etc., and atrophy of the skull in chronic hydrocephalus, are well-known examples of these atrophies from pressure.

Mechanical congestion in the same way is not an uncommon cause of atrophy. The circulation is impeded, the blood is not returned normally by the veins, hence there is deficient arterial supply, and atrophy results.

2. *Diminished Functional Activity.*—This is the most common cause of atrophy, many examples of which are furnished both by physiological and pathological processes. After birth those parts which are no longer required to serve any purpose in the economy gradually atrophy and waste. The ductus arteriosus, the umbilical arteries and vein, the Wolffian bodies, and later—the thymus glands, all in this manner disappear. The involution of the uterus after delivery, the wasting of the spleen and lymphatic glands in advanced life, and of the lower jaw after the loss of the teeth, are other physiological examples of atrophy from this cause.

Muscles which from any cause have long remained inactive atrophy. This is seen in the various forms of paralysis, especially in the so-called essential paralysis of children; also in limbs which have become incapacitated, either on account of ankylosis, or of chronic diseases of the bones or joints.

After the establishment of an artificial anus, the lower part of the intestine atrophies, and becomes converted into a fibro-cellular cord.

Bones in the same manner atrophy from want of use. After the amputation of a limb, the cut end of the bone atrophies; and atrophy of the orbit follows extirpation of the eyeball.

Interference with the function of nerves is also followed by their atrophy. This is seen in the atrophy of the optic nerve, which often follows destruction of the eye, and also in the atrophy of the nerves of a limb which has long been incapable of action.

3. *Increased Functional Activity.*—This may in quite exceptional cases be a cause of atrophy; much more commonly, however, it is a cause of hypertrophy. Certain glands sometimes atrophy from excessive use, especially the testicle.

4. *The Action of Special Substances.*—Certain substances administered internally appear to be capable of producing atrophy. Iodine, bromine, mercury, lead, and the alkalies, may be enumerated amongst the most important of these. Iodine and mercury exercise a special influence upon the lymphatic system, and bromine upon the organs of generation.

5. *Nervous Influence.*—Respecting the influence of the nervous system as a *direct* cause of atrophy, little is certainly known. That atrophy is a frequent sequence of changes in the nerve-trunks and nervous centres, there can be no doubt; but it is probably in most cases to be attributed to an *indirect* influence. The atrophy of muscles, for example, which have become paralyzed from lesions in the brain or spinal cord, is usually rather the result of the consequent interference with their function, than of any direct influence upon their nutrition. The very rapid atrophy of muscles, however, which follows certain alterations in the nervous centres and injuries of the nerve-trunks, renders it exceedingly probable that some nerves contain fibres which exercise a direct influence upon nutrition, and that nervous influence may also be a *direct* cause of atrophy. (See “Introduction.”) In some cases also, changes in the nervous system may cause atrophy by affecting the size of the bloodvessels, and so interfering with the supply of blood.

6. *Inflammation.*—This is sometimes described as a common cause of atrophy. The atrophy, however, which is such a frequent sequence of an inflammatory process, is in most cases due to that new growth of fibroid tissue which is developed in chronic inflammations. The new tissue, by the pressure which it exercises, causes atrophy of the proper structures of the organ. This is seen, for example, in cirrhosis of the liver. It is probable, also, that the impairment of vitality which results from an inflammatory process may lead to atrophy of the histological elements, independently of the existence of new fibroid growth.

PHYSICAL CHARACTERS.—The estimation of atrophy is often a matter of considerable difficulty. The great criterion is diminution in absolute weight. The weight of an organ, however, varies considerably in health; it varies with the weight of the body as a whole, and it may be less than natural from incomplete development. The same is true also of the muscular and osseous systems. An accumulation of blood and serosity in an organ may again increase its weight, and thus constitute a source of fallacy. This is often the case in organs which have been for some time mechanically congested, in which,

although their size and weight may be increased, their tissue is considerably diminished in amount.

Organs which are atrophied are usually diminished not only in weight, but also in size. In most cases they contain less blood, they are drier, firmer, and more fibrous in consistence than in health. Their functional powers are invariably diminished.

The whole of the textures of which an organ is composed may suffer; some, however, do so more than others. The fibrous constituents are the last to atrophy, hence the firmness, toughness, and loss of elasticity so commonly met with in the atrophied parts. In glandular organs, the secreting cells are usually the first to show signs of atrophy; they become smaller, and are often finely granular, from the presence of molecular fat; the vessels and nerves also share in the wasting process. In the subcutaneous cellular tissue, the fat is gradually removed from the cells, which thus diminish in size. In muscles the primitive fasciculi become smaller, and their transverse striæ gradually disappear; ultimately the whole of the contents of the sarcolemma may be entirely removed, and nothing remain but the connective tissue. This process is usually accompanied by more or less fatty degeneration of the muscular fibres, and in some cases by the development of fat between the fasciculi. (See "Fatty Infiltration of Muscle.")

Atrophy of Bone.

Atrophy of bone is always attended by a diminution in weight, but not always by a diminution in size. It is met with in two forms. In one, the compact and cancellous tissue gradually becomes absorbed, the medullary canal diminishes in size, and the whole bone thus becomes smaller. This is known as *concentric atrophy*. It is met with especially in the long bones, in cases of long-standing ankylosis, dislocations, or paralysis.

In the other variety of atrophy there is no diminution in the size of the bone, but merely a gradual conversion of compact into cancellous tissue. The whole bone thus becomes rarefied, and it is exceedingly light and brittle, so that it fractures with great facility. This, in contradistinction to the former variety, is known as *eccentric atrophy*. It is usually met with as a senile change, and is in most cases accompanied by more or less fatty degeneration.

Pulmonary Vesicular Emphysema.

This appears to be the proper place to describe the changes met with in the lungs in emphysema, inasmuch as these changes are characterized mainly by atrophy of the walls of the air-vesicles.

Emphysema consists essentially in a permanent enlargement of the infundibula and air-vesicles in larger or smaller areas of the lungs. The dilatation appears usually to commence in the infundibulum, and to extend from this to the air-vesicles which open into it, so that ultimately the whole may be thrown into one large cavity. As the process proceeds, communications are established between adjacent groups of air-vesicles, and thus cavities of still larger area are produced.

Atrophous Emphysema.—The more minute histological changes which accompany emphysema vary somewhat in different varieties of the disease. In that form of emphysema which occurs in old people, and which is essentially a senile change, the alterations in the walls of the air-vesicles consist simply of atrophy of the several structures of which they are composed:—hence the term *atrophous* or *small-lunged* emphysema, which is applied by Sir W. Jenner to this variety of the disease. The air-vesicles may not be much increased in size, but several of them are thrown into one, their walls are considerably thinner than natural, the connective tissue, elastic tissue, and blood-vessels all having apparently shared in the wasting process. There is usually also an abnormal amount of pigmentation. Such lungs are smaller than natural, and quickly collapse when the thorax is opened.

Hypertrophous Emphysema.—In the other important variety of emphysema the lungs are increased in size, so that they often bulge forwards when the thorax is opened, and in contradistinction to the former variety, certain constituents of the lung-tissue appear to be increased in amount, inasmuch as the lungs are less crepitant, and feel somewhat denser and tougher than natural. This is described by Sir W. Jenner as *hypertrophous* or *large-lunged* emphysema.

When such lungs are examined microscopically, it will be found that the dilatation of the air-vesicles is more marked than in atrophous emphysema, although less general in its distribution. The atrophic changes also do not affect equally the various tissues which make up the alveolar walls. The elastic fibres appear to be more especially wasted, whilst, according to some observers, the connective tissue is increased. I have been unable to discover any marked increase of the connective tissue, in the alveolar walls, although an increase of

this tissue is often to be seen around the smaller interlobular blood-vessels and bronchi. The capillary bloodvessels which are distributed on the walls of the air-vesicles are atrophied and diminished in calibre, owing to the stretching and pressure which result from the vesicular dilatation, whilst the larger interlobular vessels are often found thickened and distended with blood. In some cases there is more or less fatty degeneration of the epithelium, and usually an abnormal pigmentation of the lung.

Etiology.—It would be beyond the scope of the present work to discuss the various theories which have been propounded to account for the development of emphysema. It is, however, obvious that all conditions which increase the pressure on the inside of the air-vesicles, or damage the resisting powers of their walls, may be causes of permanent vesicular dilatation.

Increased pressure on the inside of the air-vesicles may result from—

1st. Violent expiratory efforts with closed glottis, such as occur during the act of coughing, blowing wind instruments, violent muscular exertion, etc. Those parts of the lungs which are least supported will be over-distended. This is the expiratory theory of Jenner.

2d. Certain portions of the lungs being incapable of expansion, owing to collapse, consolidation, asthmatic spasm, etc. There will be excessive tension in those parts into which the air can enter.

Impairment of the resisting power of the air-vesicles may result from—

1st. The loss of elasticity and atrophy which is a concomitant of old age. This is the most important element in the causation of atrophous emphysema.

2d. The atrophy of the air-vesicles resulting from that stretching of their walls and obliteration of their bloodvessels which is caused by their over-distension from increased pressure exercised upon their inner surface.

3d. Damage to the walls of the air-vesicles, which probably in certain cases results from previous attacks of pulmonary inflammation, or from some interference with their nutrition due to mode of life or to other causes.

Degeneration.

The “Degenerations” include a class of morbid processes which are characterized by an alteration in the *quality* of the tissues, and which,

like atrophy, are attended by impairment of function, and often by annihilation of histological elements.

The alteration in the quality of the tissue results either from its direct metamorphosis into a new material, or from infiltration with some substance which has been conveyed to it from without.

Atrophy and degeneration thus so far resemble one another, that in both processes nutrition is impaired and function interfered with. In atrophy, however, as pointed out by Virchow, nutrition is simply altered in *quantity*, the waste of the tissue is in excess of the assimilation of new material, and consequently, there is a diminution in the amount of the tissue and an impairment of its functional powers. In degeneration, on the other hand, nutrition is altered in *quality*, a new substance exists in the tissues, which either originates in the tissue itself, or infiltrates it from without; this is attended by impairment of the vitality and functions of the elements of which the tissue is composed, resulting either from the presence of the new material, or dependent upon the same conditions as those which give rise to its formation.

CAUSES.—Of the causes of the Degenerations as a class, but little can be said, the various forms depending for the most part upon different conditions. These will be described under their respective heads.

The Degenerations may be divided into two classes—the *Metamorphoses* and the *Infiltrations*.

1. THE METAMORPHOSES.—These are characterized by the direct metamorphosis of the albuminoid constituents of the tissues into a new material. This is usually followed by the destruction of the histological elements and the softening of the intercellular substance, so that ultimately all trace of structure may be lost, and function be completely arrested. The Metamorphoses include Fatty, Muroid, and Colloid Degeneration.

2. THE INFILTRATIONS.—These differ from the Metamorphoses inasmuch as the new material which exists in the tissues is not derived from their albuminoid constituents, but is deposited in them from the blood: there is an infiltration and deposition of a new substance. This is rarely followed by destruction of the histological elements, or by softening of the intercellular substance; hence the structure of the tissue is much less altered than in the Metamorphoses, and function is usually much less interfered with. The infiltrations include Fatty, Amyloid, Calcareous, and Pigmentary Infiltration.

CHAPTER IV.

FATTY DEGENERATION.

By Fatty Degeneration is usually understood the replacement of the normal constituents of the tissues by molecules, or drops of fat. An accumulation of fat in the tissues occurs, however, under very different circumstances, and under the general term of "fatty degeneration" are included different pathological processes. Before proceeding to describe these processes and the histological changes which they produce, it will be well to consider, in the first place, the sources from which the fat met with in the body is derived; and secondly, the circumstances under which it may accumulate so as to constitute a morbid process.

General Pathology of Fatty Degeneration.—The chief source of the fat met with in the body is the oleaginous constituents of the food. A portion of these are stored up in the cells of certain tissues, to be utilized as producers of force and heat when the requirements of the system may demand it. The cells of adipose tissue, and, to a less extent, those of the liver, thus serve as physiological reservoirs for fat.

The other sources from which fat may be derived are from saccharine and albuminous principles. The albuminous principles in the process of nutrition undergo decomposition, and the products of their decomposition contain a certain amount of fat. This is usually completely removed by oxidation, but under certain circumstances the oxidation is incomplete, and the fat accumulates in the cells of the tissue.


In considering the circumstances under which an accumulation of fat in the tissues may constitute a morbid process, it is to be remarked that it often becomes exceedingly difficult here to draw any sharp line of demarcation between health and disease. This is especially the case when the accumulation of fat is excessive in situations where fat is normally met with. When it occurs in abnormal situations, the morbid nature of the process is evident.

An accumulation of fat in the tissues may occur so as to constitute a morbid process under the four following conditions:—

1st. When the food contains an excess of fat, or of substances capable of becoming converted into fat. Under such circumstances the oxygen taken into the body is insufficient to oxidize the excess, and it consequently accumulates in the cells.

An accumulation of fat from this cause occurs as a physiological process in the growth of adipose tissue. Adipose tissue is a connective tissue containing numerous cells which are distended with fat. The growth of this tissue thus consists simply in the fatty infiltration of more of these cells (Fig. 5). If this be excessive it constitutes obe-

FIG. 5.



Fatty Infiltration of Connective Tissue. Showing the accumulation of fat within the cells.
 X 300 (Rindfleisch).

sity. The temporary accumulation of fat in the liver during the digestion of an aliment rich in fatty substances is another example of this kind of deposition. This will be described when speaking of the "fatty liver." If the amount of fat be very great it may accumulate, not only in normal situations, but also in tissues where fat is not usually met with, and in both cases the accumulation will thus constitute a morbid process.

2d. When there is no such excess of fatty substances in the food, but the processes of oxidation are so imperfectly performed, either locally or generally, that the fat contained in a natural diet is incompletely oxidized.

3d. When the fat which is liberated from the nitrogenous constituents of the food during the process of nutrition does not undergo the complete oxidation which it should, and so remains within the cells.

4th. When the fat which is liberated from the protoplasm of cells during the process of nutrition is incompletely oxidized, and so accu-

mulates in, and takes the place of, the protoplasm. Here there is a gradual transformation of the protoplasm into fat, so that the cell is ultimately completely destroyed.

Fatty degeneration in which there is this destruction of histological elements is one of the most common forms of the disease, and it will hereafter be more fully described as *fatty metamorphosis*. Its nature was first pointed out by Dr. Quain in his well-known researches on fatty degeneration of the heart.¹ Dr. Quain there stated that the fat met with in the muscular fibres in this condition was the result of a metamorphosis of the fibres themselves, and was not derived from without. The truth of Dr. Quain's teaching has since been confirmed by the experimental investigations of Drs. Voit and Bauer.

Voit and Bauer's investigations were made with the object of determining the source of the fat in that acute form of fatty degeneration which is produced by poisoning with phosphorus, in which the degeneration is due to the destruction of the red blood cells, and the consequent diminution in the oxidizing power of the blood.² They gave phosphorus to dogs which had for some days previously been starved, so that any fat which might be present in the tissues after death could not have been derived either from the food or from the adipose tissue of the animals. The phosphorus produced very extensive and general fatty degeneration, and the fat must obviously have arisen from the protoplasm of the cells. Voit concludes from these investigations—1st. That the transformation of albumen which takes place in cells is independent of the supply of oxygen, but that if the oxygen be deficient, the fat and other products of the transformation, being incompletely oxidized, accumulate in the cell. 2d. That the presence of fat in the cells may thus be due to increased transformation of the albumen, or to diminished oxidation of the products of its decomposition. 3d. That the fatty degeneration in poisoning by phosphorus is due both to an increased transformation of the albumen of the cells, and to diminished oxidation of the fat and other products of the transformation.

It will thus be seen that of the four conditions enumerated as causes of fatty degeneration, in all, with the exception of the first, the accumulation of the fat is principally due to incomplete oxidation, whilst in the first there is no imperfection in the oxidizing processes, but the

¹ "Medico-Chirurgical Trans.," Lond., vol. xxxiii.

² Voit and Bauer, "Zeitschrift f. Biologie," vii. pp. 63-85; and Voit, "Neues Repertorium für Pharmacie," xx. pp. 340-349.

oxidizable materials are in excess. These two conditions are frequently associated.

An incompleteness of oxidation, and a consequent tendency to the production of fat, occur under various circumstances. The red blood cells being the carriers of oxygen, all those conditions in which the supply of blood is interfered with, the red blood cells diminished in number or defective in quality, or the oxygenation of the blood imperfectly performed, may lead to fatty degeneration. The interference with the supply of blood to the brain, for example, by thrombosis or embolism, causes fatty degeneration of the brain-tissue. (See "Cerebral Softening.") The destruction of the red blood cells by phosphorus and other poisons, and that general condition of anæmia which sometimes exists in young people and is also produced by many chronic and acute diseases, are also followed by a similar result. The senile forms of fatty degeneration which are seen in the cornea and cartilage, and are due to that diminution in the activity of the circulation which exists in old age; and the fatty degeneration of organs and tissues which have been long disused, and in which consequently the quantity of blood circulating through them and the oxidation processes become diminished (see "Fatty Infiltration of Muscle"), are other examples of degeneration from this cause. The cancers and other rapidly growing tumors in which the rapidity of growth is out of proportion to the vascular supply, also undergo extensive fatty changes. Lastly, the imperfect oxygenation of the blood which results from certain chronic diseases of the lungs constitutes one element in the causation of the fatty degeneration which so frequently exists in these diseases.

In proceeding to consider the histological changes which are produced in the tissues by an accumulation of fat, it must be stated, in the first place, that these vary very considerably according to the cause of the degeneration. In those cases in which the fat is derived from the metamorphosis of the nitrogenous constituents of the tissues, the tissues are destroyed in the process; whilst in other cases no such destruction usually takes place. Although these two conditions may occasionally be associated, yet, owing to the marked difference in the results which they respectively produce, it will be well to speak of them separately; that in which the fat is derived from the metamorphosis of the tissues being termed *Fatty Metamorphosis*, that in which it is derived from the oleaginous, saccharine, or nitrogenous principles of the food—*Fatty Infiltration*.

Fatty Infiltration.

In Fatty Infiltration, the fat which is deposited within the cells usually occurs as distinct drops of oil. In the earliest stages of the process these are very small, but as the deposition proceeds they gradually accumulate and run together, displacing and obscuring the nucleus and protoplasm, until the cell is completely filled and distended with oil. (Fig. 6.) The vitality and functions of the cells are but little impaired by the accumulation, and the protoplasm—although rendered almost invisible when this is excessive—remains unaltered. The cells within which the fat accumulates not being destroyed, the removal of the fat is all that is necessary to restore them to their original condition. As already stated, fatty infiltration occurs as a physiological process in the growth of adipose tissue, and also in the liver during the digestion of an aliment rich in fatty substances.





FIG. 6.
Liver Cells in various stages of Fatty Infiltration. $\times 300$ (Rindfleisch).

Fatty Infiltration of Muscle.

In muscle, fatty infiltration is frequently met with as a morbid process. The cells in the connective tissue which surrounds the fasciculi of the muscle become filled with fat; and this development of fat between the primitive muscular fasciculi has often been confounded with degeneration of the fibres themselves. In this latter process, however, which will subsequently be described as *fatty metamorphosis* of muscle, there is a direct metamorphosis of the muscular fibres into fat; whereas in the condition now under consideration, there is a deposition of fat *between* the fasciculi, which remain—during the early stages at all events—unaffected. The interstitial fat varies in amount. In some cases single rows of fat cells alternate with rows of muscular fasciculi, at other times the accumulation is less regular, more existing between some fibres than between others; in all cases, however, the muscular elements may be discovered lying amongst the fat. (Fig. 7.) If the latter be very considerable in amount, the muscle may appear to the naked eye to be entirely converted into fat; but the microscope will always reveal the muscular structure in which it is imbedded.

This condition is frequently met with in animals which have been fattened, the fat increasing not only in the usual situations, but also accumulating between the fasciculi of the muscles. In muscles also which from any cause have for some time been incapacitated, and in

FIG. 7.



Fatty Infiltration of Heart. A section from the more external portion of the left ventricle of the heart, showing the growth of fat *between* the muscular fibres. The fibres are in some places atrophied and commencing to undergo fatty metamorphosis. $\times 200$.

which consequently the circulation of blood and the oxidation processes are reduced to a minimum, this interstitial growth is extremely liable to occur; *ex. gr.*, in the extensors of the wrist-joint in cases of lead-poisoning, and in long-standing paralysis from lesions of the brain or cord, also in muscles which have been rendered useless by ankylosis of a joint. In progressive muscular atrophy, as Virchow has shown, the affected muscles exhibit this change, together with true fatty metamorphosis.

Fatty Infiltration of the Heart.—In the heart fatty infiltration is not unfrequently met with; and here it is especially important to distinguish it from the much more grave condition in which the fibres themselves are primarily affected. In health, there is a varying amount of fat covering the surface of the heart beneath the visceral layer of the pericardium, which is always most abundant in the grooves

between the auricles and ventricles, where it surrounds the blood-vessels. This may increase so as to completely envelop the organ, and at the same time gradually insinuate itself between the muscular fibres, so that to the naked eye all appearance of muscular structure may be lost, the walls looking like a mass of fat. In hearts less affected, striæ of fat will be seen lying amongst the muscles. (See Fig. 7.) The fat is always most abundant near the surface, the muscular structure becoming more evident towards the endocardium.

The immediate effect of the interstitial growth is to displace and compress the muscular fibres between which it insinuates itself, and in doing so it diminishes the contractile power of the muscle. This is especially important when occurring in the heart. The pressure, however, which it exercises upon the fibres and the accompanying blood-vessels, ultimately causes atrophic and degenerative changes. Thus the fasciculi gradually atrophy, the transverse striation becomes indistinct and is replaced by molecular fat; in fine, true metamorphosis of the muscle is established. These two processes, indeed, not uncommonly go hand in hand together, the interstitial infiltration preceding the intrastitial metamorphosis.

Fatty Infiltration of the Liver.

It is in the liver that fatty infiltration assumes its most important aspect, and in this organ it is exceedingly frequent, constituting what is commonly known as the "fatty liver." The hepatic cells always contain a small quantity of fat, which is temporarily increased after the ingestion of fatty substances. It will be well to describe this physiological infiltration before proceeding to the morbid process.

The ingestion of an aliment rich in fatty substances is followed by a temporary excess of fat in the portal blood, and by the deposition and temporary accumulation of a portion of this within the hepatic cells. This fat is first deposited in the cells which are in immediate contact with the capillaries of the portal vein, and thus is produced an excess of fat in the cells at the circumference of the hepatic lobules. This gradually passes from the cells at the circumference to those in the interior, whence it is ultimately conveyed again into the circulation. This process goes on until the excess of fat is removed from the blood, when the hepatic cells again acquire their former character. There is thus a transitory accumulation of fat within the hepatic cells


which is gradually removed, the vitality of the cells not being thereby impaired.

The morbidly fatty liver is one which contains an abnormal quantity of fat, and here also, as the fat is usually deposited from the blood in the portal capillaries, the increase is first observable in the external zone of the hepatic lobules. (Fig. 8.) It accumulates here within the cells as minute globules, which as they increase coalesce and form large drops of fat. These ultimately completely fill and distend the cells, which at the same time become larger and more globular in shape. (See Fig. 6.) As the process proceeds, the accumulation advances from the periphery towards the centre of the lobule, until its whole mass may be involved, and the cells universally become distended with fat. The vitality of the cells is not materially impaired by the infiltration; they continue to perform their functions, as is shown by the presence of bile in the stools and in the gall-bladder. In many exceptional cases the accumulation of fat is most marked around the hepatic vein. This, according to Virchow, is probably to be explained by supposing that the fat is becoming excreted, and that only the last cells retain a little of it.

The fatty liver is somewhat increased in size, in advanced stages often considerably so. The surface is smooth, the edges are thickened and rounded, the specific gravity is diminished, although the absolute weight may be increased. If the infiltration be slight, involving merely the portal zone of the lobules, the cut surface will present a mottled appearance, the external fatty zone being of an opaque yellowish-white color, whilst the central portion remains unaltered, or is perhaps somewhat hyperæmic. The more extensive the infiltration the larger is the pale zone, and ultimately, when the whole lobule is involved, there may be left in the centre merely a reddish-brown point, which corresponds with the commencement of the hepatic vein; and in many cases even this point is lost. The organ is then of an almost uniform opaque yellowish-white color, and the boundary between the individual lobules may be completely obscured. In very exceptional cases the accumulation of fat is much more abundant in some portions of the liver than in others, so that on section, yellowish points and streaks are seen scattered over its surface. The consistence of the organ is much diminished, it feels doughy, and pits on pressure with the finger, and the knife used to cut it becomes coated with oil. The pressure exercised by the infiltrated fat produces considerable anæmia of the

organ, but the interference with the circulation is never sufficient to cause ascites, hemorrhage, or other evidence of portal congestion.

FIG. 8.



Fatty Liver. Showing the accumulation of fat in those cells more especially which are situated in the external zone of the lobule. There is also an increase in the interlobular connective tissue (cirrhosis). V. Hepatic vein. I. Interlobular connective tissue. $\times 50$.

The liver is especially liable to become the seat of fatty accumulation. This, according to the late Dr. Bence Jones, is owing—firstly, to the excess of non-nitrogenous oxidizable matter which is always present in it; secondly, to the deoxidized condition of the portal blood; and thirdly, to the low pressure and slowness of circulation in the portal vessels—conditions the least favorable to oxidation.¹

An accumulation of fat in the liver occurs under two opposite conditions—one in which there is general obesity, and the fat accumulates in the liver in common with other parts; and another, in which there is general emaciation, and a consequent impairment of the oxygenating power of the blood. The fatty infiltration of the liver which is so constantly associated with certain chronic diseases of the lungs, is probably also partly due to imperfect oxygenation of the blood from destruction of lung-tissue. Fatty liver caused by phosphorus and other poisons has been already alluded to.

The other variety of fatty degeneration—fatty metamorphosis, will be described in the following chapter.

¹ "Lectures on Pathology and Therapeutics." Dr. Bence Jones, p. 179.

CHAPTER V.

FATTY DEGENERATION (*continued*).**Fatty Metamorphosis.**

THIS differs from fatty infiltration, inasmuch as the fat is derived from the albuminous constituents of the tissues themselves, and not from the fatty, saccharine, or nitrogenous principles of the food.

The process consists in the transformation of the albuminous constituents of the tissues into fat, in the manner described in the preceding chapter, the cells being the parts which are most frequently affected. This fat makes its appearance as minute granules and molecules within the cells, usually first in the protoplasm, and subsequently in the nucleus. The granules—which are characterized by their dark color, sharp contour, strong refractive power, and solubility in ether—gradually increase in number, until the whole of the protoplasm has undergone the transformation. As they increase some of them may coalesce, and so form distinct drops of fat. As the process proceeds the cells undergo an increase in size and become more globular in shape. Ultimately the nucleus becomes involved, the cell-wall,

when this exists, is destroyed, and the cell is thus transformed into a mass of granular fat (Fig. 9).

These granules of fat may remain in a coherent form for some time; they then constitute what were formerly known as the “inflammatory” or “exudation corpuscles,” or “corpuscles of Gluge,” which are so common in chronic cerebral softening, and in other forms of fatty degeneration (Fig. 9, *a*). Ultimately

the corpuscles break up, the albuminous matter between the granules of fat liquefies, and the fat becomes distributed in the tissue (Fig. 9, *b*).

FIG. 9.



Fatty Metamorphosis of Cells. *a.* From a cancer. *b.* From the brain in chronic softening. The latter show the large “granular corpuscles,” and also the manner in which these become disintegrated. $\times 200$.

Types of this pathological process are furnished by many well-known physiological ones, one of the most characteristic of which is perhaps the secretion of milk. The mammary gland is a large racemose gland, consisting of numerous groups of lobules lined with epithelial cells. The secretion of milk takes place in the following manner: The cells lining the lobules of the gland multiply abundantly, and the new cells as they are produced gradually become converted into fat; the cells break up, and the fatty matters in a more or less coherent form constitute the milk-corpuscles. In the earliest stages of the process the granules of fat cohere and form the colostrum-corpuscles, which are precisely analogous to the large granular corpuscles met with in chronic cerebral softening, etc. (Fig. 9, *b*); but as the secretion becomes fully established, and the multiplication and disintegration of the cells take place more rapidly, the fatty molecules become at once distributed in the liquid in which they are suspended, giving to the secretion its characteristic white color. The milk-corpuscles thus formed are replaced by the continuous formation of new cells from below, which in their turn undergo fatty metamorphosis, and in this manner a continuous formation and destruction of cells take place.

Other examples of fatty metamorphosis are afforded by the formation of the sebaceous matter of the skin, the cerumen of the ears, and the corpus luteum in the ovary; all of which take place in the same way by the fatty metamorphosis and destruction of newly formed cells.

The immediate effect of fatty metamorphosis is to produce more or less softening of the affected part, the cellular elements are completely destroyed, the intercellular substance also undergoes fatty changes, and thus all trace of the original structure may be ultimately lost. This destruction of the cells is the essential feature of the change, and distinguishes it from fatty *infiltration*, in which the cells within which the fat accumulates usually remain intact.

If large tracts of tissue are affected, the change is readily recognizable by the diminution in consistence and elasticity which is produced, and in many cases also by the opaque yellowish-white color. If, however, the change is limited to minute and isolated portions of the tissue, its existence can only be discovered with the aid of the microscope.

The fatty particles into which the cells have been transformed under favorable circumstances are readily absorbed. This is seen in the fatty degeneration and absorption of inflammatory products, such as occurs in croupous pneumonia. In order for such absorption to

1st. When the food contains an excess of fat, or of substances capable of becoming converted into fat. Under such circumstances the oxygen taken into the body is insufficient to oxidize the excess, and it consequently accumulates in the cells.

An accumulation of fat from this cause occurs as a physiological process in the growth of adipose tissue. Adipose tissue is a connective tissue containing numerous cells which are distended with fat. The growth of this tissue thus consists simply in the fatty infiltration of more of these cells (Fig. 5). If this be excessive it constitutes obe-

FIG. 5.



Fatty Infiltration of Connective Tissue. Showing the accumulation of fat within the cells.
X 300 (Rindfleisch).

sity. The temporary accumulation of fat in the liver during the digestion of an aliment rich in fatty substances is another example of this kind of deposition. This will be described when speaking of the "fatty liver." If the amount of fat be very great it may accumulate, not only in normal situations, but also in tissues where fat is not usually met with, and in both cases the accumulation will thus constitute a morbid process.

2d. When there is no such excess of fatty substances in the food, but the processes of oxidation are so imperfectly performed, either locally or generally, that the fat contained in a natural diet is incompletely oxidized.

3d. When the fat which is liberated from the nitrogenous constituents of the food during the process of nutrition does not undergo the complete oxidation which it should, and so remains within the cells.

4th. When the fat which is liberated from the protoplasm of cells during the process of nutrition is incompletely oxidized, and so accu-

mulates in, and takes the place of, the protoplasm. Here there is a gradual transformation of the protoplasm into fat, so that the cell is ultimately completely destroyed.

Fatty degeneration in which there is this destruction of histological elements is one of the most common forms of the disease, and it will hereafter be more fully described as *fatty metamorphosis*. Its nature was first pointed out by Dr. Quain in his well-known researches on fatty degeneration of the heart.¹ Dr. Quain there stated that the fat met with in the muscular fibres in this condition was the result of a metamorphosis of the fibres themselves, and was not derived from without. The truth of Dr. Quain's teaching has since been confirmed by the experimental investigations of Drs. Voit and Bauer.

Voit and Bauer's investigations were made with the object of determining the source of the fat in that acute form of fatty degeneration which is produced by poisoning with phosphorus, in which the degeneration is due to the destruction of the red blood cells, and the consequent diminution in the oxidizing power of the blood.² They gave phosphorus to dogs which had for some days previously been starved, so that any fat which might be present in the tissues after death could not have been derived either from the food or from the adipose tissue of the animals. The phosphorus produced very extensive and general fatty degeneration, and the fat must obviously have arisen from the protoplasm of the cells. Voit concludes from these investigations—1st. That the transformation of albumen which takes place in cells is independent of the supply of oxygen, but that if the oxygen be deficient, the fat and other products of the transformation, being incompletely oxidized, accumulate in the cell. 2d. That the presence of fat in the cells may thus be due to increased transformation of the albumen, or to diminished oxidation of the products of its decomposition. 3d. That the fatty degeneration in poisoning by phosphorus is due both to an increased transformation of the albumen of the cells, and to diminished oxidation of the fat and other products of the transformation.

It will thus be seen that of the four conditions enumerated as causes of fatty degeneration, in all, with the exception of the first, the accumulation of the fat is principally due to incomplete oxidation, whilst in the first there is no imperfection in the oxidizing processes, but the

¹ "Medico-Chirurgical Trans.," Lond., vol. xxxiii.

² Voit and Bauer, "Zeitschrift f. Biologie," vii. pp. 63-85; and Voit, "Neues Repertorium für Pharmacie," xx. pp. 340-349.


oxidizable materials are in excess. These two conditions are frequently associated.

An incompleteness of oxidation, and a consequent tendency to the production of fat, occur under various circumstances. The red blood cells being the carriers of oxygen, all those conditions in which the supply of blood is interfered with, the red blood cells diminished in number or defective in quality, or the oxygenation of the blood imperfectly performed, may lead to fatty degeneration. The interference with the supply of blood to the brain, for example, by thrombosis or embolism, causes fatty degeneration of the brain-tissue. (See "Cerebral Softening.") The destruction of the red blood cells by phosphorus and other poisons, and that general condition of anaemia which sometimes exists in young people and is also produced by many chronic and acute diseases, are also followed by a similar result. The senile forms of fatty degeneration which are seen in the cornea and cartilage, and are due to that diminution in the activity of the circulation which exists in old age; and the fatty degeneration of organs and tissues which have been long disused, and in which consequently the quantity of blood circulating through them and the oxidation processes become diminished (see "Fatty Infiltration of Muscle"), are other examples of degeneration from this cause. The cancers and other rapidly growing tumors in which the rapidity of growth is out of proportion to the vascular supply, also undergo extensive fatty changes. Lastly, the imperfect oxygenation of the blood which results from certain chronic diseases of the lungs constitutes one element in the causation of the fatty degeneration which so frequently exists in these diseases.

In proceeding to consider the histological changes which are produced in the tissues by an accumulation of fat, it must be stated, in the first place, that these vary very considerably according to the cause of the degeneration. In those cases in which the fat is derived from the metamorphosis of the nitrogenous constituents of the tissues, the tissues are destroyed in the process; whilst in other cases no such destruction usually takes place. Although these two conditions may occasionally be associated, yet, owing to the marked difference in the results which they respectively produce, it will be well to speak of them separately; that in which the fat is derived from the metamorphosis of the tissues being termed *Fatty Metamorphosis*, that in which it is derived from the oleaginous, saccharine, or nitrogenous principles of the food—*Fatty Infiltration*.

Fatty Infiltration.

In Fatty Infiltration, the fat which is deposited within the cells usually occurs as distinct drops of oil. In the earliest stages of the process these are very small, but as the deposition proceeds they gradually accumulate and run together, displacing and obscuring the nucleus and protoplasm, until the cell is completely filled and distended with oil. (Fig. 6.) The vitality and functions of the cells are but little impaired by the accumulation, and the protoplasm—although rendered almost invisible when this is excessive—remains unaltered. The cells within which the fat accumulates not being destroyed, the removal of the fat is all that is necessary to restore them to their original condition. As already stated, fatty infiltration occurs as a physiological process in the growth of adipose tissue, and also in the liver during the digestion of an aliment rich in fatty substances.




Fatty Infiltration of Muscle.

In muscle, fatty infiltration is frequently met with as a morbid process. The cells in the connective tissue which surrounds the fasciculi of the muscle become filled with fat; and this development of fat between the primitive muscular fasciculi has often been confounded with degeneration of the fibres themselves. In this latter process, however, which will subsequently be described as *fatty metamorphosis* of muscle, there is a direct metamorphosis of the muscular fibres into fat; whereas in the condition now under consideration, there is a deposition of fat *between* the fasciculi, which remain—during the early stages at all events—unaffected. The interstitial fat varies in amount. In some cases single rows of fat cells alternate with rows of muscular fasciculi, at other times the accumulation is less regular, more existing between some fibres than between others; in all cases, however, the muscular elements may be discovered lying amongst the fat. (Fig. 7.) If the latter be very considerable in amount, the muscle may appear to the naked eye to be entirely converted into fat; but the microscope will always reveal the muscular structure in which it is imbedded.

This condition is frequently met with in animals which have been fattened, the fat increasing not only in the usual situations, but also accumulating between the fasciculi of the muscles. In muscles also which from any cause have for some time been incapacitated, and in

FIG. 7.



Fatty Infiltration of Heart. A section from the more external portion of the left ventricle of the heart, showing the growth of fat *between* the muscular fibres. The fibres are in some places atrophied and commencing to undergo fatty metamorphosis. $\times 200$.

which consequently the circulation of blood and the oxidation processes are reduced to a minimum, this interstitial growth is extremely liable to occur; *ex. gr.*, in the extensors of the wrist-joint in cases of lead-poisoning, and in long-standing paralysis from lesions of the brain or cord, also in muscles which have been rendered useless by ankylosis of a joint. In progressive muscular atrophy, as Virchow has shown, the affected muscles exhibit this change, together with true fatty metamorphosis.

Fatty Infiltration of the Heart.—In the heart fatty infiltration is not unfrequently met with; and here it is especially important to distinguish it from the much more grave condition in which the fibres themselves are primarily affected. In health, there is a varying amount of fat covering the surface of the heart beneath the visceral layer of the pericardium, which is always most abundant in the grooves

between the auricles and ventricles, where it surrounds the blood-vessels. This may increase so as to completely envelop the organ, and at the same time gradually insinuate itself between the muscular fibres, so that to the naked eye all appearance of muscular structure may be lost, the walls looking like a mass of fat. In hearts less affected, striæ of fat will be seen lying amongst the muscles. (See Fig. 7.) The fat is always most abundant near the surface, the muscular structure becoming more evident towards the endocardium.

The immediate effect of the interstitial growth is to displace and compress the muscular fibres between which it insinuates itself, and in doing so it diminishes the contractile power of the muscle. This is especially important when occurring in the heart. The pressure, however, which it exercises upon the fibres and the accompanying blood-vessels, ultimately causes atrophic and degenerative changes. Thus the fasciculi gradually atrophy, the transverse striation becomes indistinct and is replaced by molecular fat; in fine, true metamorphosis of the muscle is established. These two processes, indeed, not uncommonly go hand in hand together, the interstitial infiltration preceding the intrastitial metamorphosis.

Fatty Infiltration of the Liver.

It is in the liver that fatty infiltration assumes its most important aspect, and in this organ it is exceedingly frequent, constituting what is commonly known as the "fatty liver." The hepatic cells always contain a small quantity of fat, which is temporarily increased after the ingestion of fatty substances. It will be well to describe this physiological infiltration before proceeding to the morbid process.

The ingestion of an aliment rich in fatty substances is followed by a temporary excess of fat in the portal blood, and by the deposition and temporary accumulation of a portion of this within the hepatic cells. This fat is first deposited in the cells which are in immediate contact with the capillaries of the portal vein, and thus is produced an excess of fat in the cells at the circumference of the hepatic lobules. This gradually passes from the cells at the circumference to those in the interior, whence it is ultimately conveyed again into the circulation. This process goes on until the excess of fat is removed from the blood, when the hepatic cells again acquire their former character. There is thus a transitory accumulation of fat within the hepatic cells

.....


which is gradually removed, the vitality of the cells not being thereby impaired.

The morbidly fatty liver is one which contains an abnormal quantity of fat, and here also, as the fat is usually deposited from the blood in the portal capillaries, the increase is first observable in the external zone of the hepatic lobules. (Fig. 8.) It accumulates here within the cells as minute globules, which as they increase coalesce and form large drops of fat. These ultimately completely fill and distend the cells, which at the same time become larger and more globular in shape. (See Fig. 6.) As the process proceeds, the accumulation advances from the periphery towards the centre of the lobule, until its whole mass may be involved, and the cells universally become distended with fat. The vitality of the cells is not materially impaired by the infiltration; they continue to perform their functions, as is shown by the presence of bile in the stools and in the gall-bladder. In many exceptional cases the accumulation of fat is most marked around the hepatic vein. This, according to Virchow, is probably to be explained by supposing that the fat is becoming excreted, and that only the last cells retain a little of it.

The fatty liver is somewhat increased in size, in advanced stages often considerably so. The surface is smooth, the edges are thickened and rounded, the specific gravity is diminished, although the absolute weight may be increased. If the infiltration be slight, involving merely the portal zone of the lobules, the cut surface will present a mottled appearance, the external fatty zone being of an opaque yellowish-white color, whilst the central portion remains unaltered, or is perhaps somewhat hyperæmic. The more extensive the infiltration the larger is the pale zone, and ultimately, when the whole lobule is involved, there may be left in the centre merely a reddish-brown point, which corresponds with the commencement of the hepatic vein; and in many cases even this point is lost. The organ is then of an almost uniform opaque yellowish-white color, and the boundary between the individual lobules may be completely obscured. In very exceptional cases the accumulation of fat is much more abundant in some portions of the liver than in others, so that on section, yellowish points and streaks are seen scattered over its surface. The consistence of the organ is much diminished, it feels doughy, and pits on pressure with the finger, and the knife used to cut it becomes coated with oil. The pressure exercised by the infiltrated fat produces considerable anæmia of the

organ, but the interference with the circulation is never sufficient to cause ascites, hemorrhage, or other evidence of portal congestion.

FIG. 8.



Fatty Liver. Showing the accumulation of fat in those cells more especially which are situated in the external zone of the lobule. There is also an increase in the interlobular connective tissue (cirrhosis). V. Hepatic vein. I. Interlobular connective tissue. $\times 50$.

The liver is especially liable to become the seat of fatty accumulation. This, according to the late Dr. Bence Jones, is owing—firstly, to the excess of non-nitrogenous oxidizable matter which is always present in it; secondly, to the deoxidized condition of the portal blood; and thirdly, to the low pressure and slowness of circulation in the portal vessels—conditions the least favorable to oxidation.¹

An accumulation of fat in the liver occurs under two opposite conditions—one in which there is general obesity, and the fat accumulates in the liver in common with other parts; and another, in which there is general emaciation, and a consequent impairment of the oxygenating power of the blood. The fatty infiltration of the liver which is so constantly associated with certain chronic diseases of the lungs, is probably also partly due to imperfect oxygenation of the blood from destruction of lung-tissue. Fatty liver caused by phosphorus and other poisons has been already alluded to.

The other variety of fatty degeneration—fatty metamorphosis, will be described in the following chapter.

¹ "Lectures on Pathology and Therapeutics." Dr. Bence Jones, p. 179.

CHAPTER V.

FATTY DEGENERATION (*continued*).**Fatty Metamorphosis.**

THIS differs from fatty infiltration, inasmuch as the fat is derived from the albuminous constituents of the tissues themselves, and not from the fatty, saccharine, or nitrogenous principles of the food.

The process consists in the transformation of the albuminous constituents of the tissues into fat, in the manner described in the preceding chapter, the cells being the parts which are most frequently affected. This fat makes its appearance as minute granules and molecules within the cells, usually first in the protoplasm, and subsequently in the nucleus. The granules—which are characterized by their dark color, sharp contour, strong refractive power, and solubility in ether—gradually increase in number, until the whole of the protoplasm has undergone the transformation. As they increase some of them may coalesce, and so form distinct drops of fat. As the process proceeds the cells undergo an increase in size and become more globular in shape. Ultimately the nucleus becomes involved, the cell-wall,

when this exists, is destroyed, and the cell is thus transformed into a mass of granular fat (Fig. 9).

These granules of fat may remain in a coherent form for some time; they then constitute what were formerly known as the “inflammatory” or “exudation corpuscles,” or “corpuscles of Gluge,” which are so common in chronic cerebral softening, and in other forms of fatty degeneration (Fig. 9, *a*). Ultimately

the corpuscles break up, the albuminous matter between the granules of fat liquefies, and the fat becomes distributed in the tissue (Fig. 9, *b*).

FIG. 9.



Fatty Metamorphosis of Cells. a. From a cancer. b. From the brain in chronic softening. The latter show the large “granular corpuscles,” and also the manner in which these become disintegrated. × 200.

Types of this pathological process are furnished by many well-known physiological ones, one of the most characteristic of which is perhaps the secretion of milk. The mammary gland is a large racemose gland, consisting of numerous groups of lobules lined with epithelial cells. The secretion of milk takes place in the following manner: The cells lining the lobules of the gland multiply abundantly, and the new cells as they are produced gradually become converted into fat; the cells break up, and the fatty matters in a more or less coherent form constitute the milk-corpuscles. In the earliest stages of the process the granules of fat cohere and form the colostrum-corpuscles, which are precisely analogous to the large granular corpuscles met with in chronic cerebral softening, etc. (Fig. 9, *b*); but as the secretion becomes fully established, and the multiplication and disintegration of the cells take place more rapidly, the fatty molecules become at once distributed in the liquid in which they are suspended, giving to the secretion its characteristic white color. The milk-corpuscles thus formed are replaced by the continuous formation of new cells from below, which in their turn undergo fatty metamorphosis, and in this manner a continuous formation and destruction of cells take place.

Other examples of fatty metamorphosis are afforded by the formation of the sebaceous matter of the skin, the cerumen of the ears, and the corpus luteum in the ovary; all of which take place in the same way by the fatty metamorphosis and destruction of newly formed cells.

The immediate effect of fatty metamorphosis is to produce more or less softening of the affected part, the cellular elements are completely destroyed, the intercellular substance also undergoes fatty changes, and thus all trace of the original structure may be ultimately lost. This destruction of the cells is the essential feature of the change, and distinguishes it from fatty *infiltration*, in which the cells within which the fat accumulates usually remain intact.

If large tracts of tissue are affected, the change is readily recognizable by the diminution in consistence and elasticity which is produced, and in many cases also by the opaque yellowish-white color. If, however, the change is limited to minute and isolated portions of the tissue, its existence can only be discovered with the aid of the microscope.

The fatty particles into which the cells have been transformed under favorable circumstances are readily absorbed. This is seen in the fatty degeneration and absorption of inflammatory products, such as occurs in croupous pneumonia. In order for such absorption to

take place, it is necessary that the tissue should be freely supplied with bloodvessels. If this is not the case, the degenerated products are liable to undergo certain changes whereby they become converted into a pultaceous crumbling material somewhat resembling cheese:—this is known as *caseation*.

CASEATION.—This is a modification of the degenerative process in which the fatty products gradually dry up into a yellowish friable material, which has been compared to soft cheese. This change appears to be owing to a natural dryness of the degenerated tissue, resulting from deficient vascular supply. It is most frequent in parts which contain but few vessels, or in those in which these become obliterated by inflammatory products or by some new growth. Growths composed of closely-crowded cells—as epithelial accumulations within the pulmonary alveoli, growths in the lymphatic glands, in the brain, and in the osseous structure, are the most liable to become caseous.

The process consists in a gradual drying up of the degenerated elements; the fluids are absorbed, the cells—which are many of them incompletely degenerated—shrink and atrophy, the fat undergoes partial saponification, cholesterine forms, and the tissue thus becomes converted into a soft, yellowish-white, cheesy substance, composed of atrophied cells, fatty debris, and cholesterine crystals. This material may gradually dry up more and more, and ultimately become encapsuled by a layer of fibrous tissue.

These cheesy matters are constantly met with, especially in the lungs, and considerable confusion has arisen as to their nature and origin in this situation. This has proceeded from its having been formerly the custom to look upon all cheesy masses as essentially tubercular. Tubercle, it is true, often undergoes, to a greater or less extent, fatty degeneration, and it may thus, like all other growths which have undergone this process, become converted into a yellow cheesy substance; but it is by no means true that all cheesy masses are tubercular. The pathological significance of caseation is thus less limited than was formerly supposed, and although this change is undoubtedly most common in tuberculous and scrofulous lesions, its occurrence merely indicates that the elements have undergone fatty metamorphosis, and under no circumstances is it in itself evidence of any one particular form of morbid growth. (See “Scrofulous Inflammation.”)

The caseous mass may subsequently become calcified, or undergo a process of softening and liquefaction.

CALCIFICATION.—This is an advanced stage of the preceding process. It most frequently occurs in those cases in which the caseous mass is completely inclosed and isolated from the external air, as when in the lymphatic glands, in bone, or when encapsuled in the lungs. The mass becomes infiltrated with calcareous particles, and is thus converted into a calcareous concretion. (See “Calcareous Degeneration.”)

SOFTENING.—This process consists in a liquefaction of the caseous substance, which is probably owing to some chemical change in its constituents. It most commonly occurs in parts which come into contact with the external air, especially in those situated in the lungs. The caseous mass liquefies, and is converted into a thin puriform liquid, containing curd-like cheesy matter, which to the naked eye looks much like pus, but under the microscope is seen to consist simply of granular debris, fat, and cholesterine crystals. This, if not discharged, may, like the caseous masses, ultimately dry up and become calcified.


Fatty Degeneration of Arteries.

Fatty degeneration of arteries may be a primary or secondary affection. As a secondary process it is met with in atheroma and other inflammatory conditions of the vessels in which the fatty change is preceded by a cellular infiltration of the sub-endothelial connective tissue (see “Atheroma”).

Primary fatty degeneration is a passive process, not being preceded by any increased nutritive activity of the parts affected by it. It may affect both the internal, middle, and external coats of the artery, but it is most common in the first-named situation. The change usually commences

in the endothelium and the connective tissue cells in the most internal layers of the inner coat, small groups of cells becoming affected in various parts of the vessel; and it may gradually extend from within outwards, the intercellular substance softening, until, in exceptional cases, the whole thickness of the intima is destroyed. (Fig. 10.)

FIG. 10.



Fatty Degeneration of the Internal Coat of the Aorta. Minute yellowish white patches scattered over the lining membrane of the vessel. A very thin layer peeled off and $\times 200$, showing the groups of fat molecules, and the distribution of fat in the intima.

In the earlier stages of the process, the condition is recognized by the existence of small, irregular-shaped patches of an opaque yellowish-white color, projecting very slightly above the surface of the intima. These, which are so constantly met with on the lining membrane of the aorta, may at first be mistaken for atheroma. They are in most cases, however, readily distinguishable by their superficiality, and by the facility with which they can be stripped off from the subjacent layers, which present a natural appearance. In atheroma, on the other hand—which affects the deeper structures—if the superficial layer be removed, the opacity and thickening are seen to exist beneath it. In many cases the change is limited entirely to the innermost layers of the vessel; the more the subjacent tissues are involved, the greater is the irregularity in the shape of the patches, and less readily can they be separated with the forceps. The opaque patches occasionally break down, the cells are destroyed, the intercellular substance softens, and the granular debris is carried away by the circulation, leaving small, irregular, superficial erosions upon the lining membrane of the vessel. These erosions are not ulcers in the true sense of that term, not being the result of an active process. They resemble the superficial erosions so common upon the mucous membrane of the stomach, as described by Dr. Wilson Fox.

Simple fatty degeneration may occur in any of the arteries, but it is in the smaller ones that its injurious influence is most marked, and in these it is more especially liable to affect the external coat. Here, by diminishing the elasticity and contractility of the vessels, it causes degenerative changes in the parts which they supply, and often leads to rupture. This is exemplified by many cases of chronic cerebral softening and cerebral hemorrhage, although here atheromatous are often associated with the fatty changes. In the larger arteries, as the aorta—where it is exceedingly common—it is of less importance, the inflammatory process, atheroma, having here a far more deleterious effect.

Fatty Degeneration of Capillaries.—The capillaries may also be the seat of fatty changes. Here they are most common in the nervous centres, and in the kidneys in Bright's disease. The process commences in the endothelial cells, and may involve considerable areas of the capillary wall, so that rupture is often the ultimate result. This is common in the smallest cerebral bloodvessels, where it is sometimes a cause of cerebral (capillary) hemorrhage.

Primary fatty degeneration of bloodvessels is in most cases a senile

change; it is an expression of that general impairment of vitality which exists in advanced life, and is usually associated with similar changes in other parts. When, however, it is limited to the lining membrane of the largest arteries, it is often met with in early life and in persons who are otherwise perfectly healthy. Fatty degeneration secondary to inflammatory conditions will be considered subsequently. (See "Inflammation of Bloodvessels.")

Fatty Degeneration of the Brain.—Cerebral Softening.

Fatty degeneration of the brain is met with in those morbid conditions comprised under the common term of "Cerebral Softenings." Whatever interferes with the supply of blood to the cerebral substance will tend to produce fatty degeneration, and hence softening. The portions of the brain which are the seat of this change may be merely rather softer than the surrounding healthy tissue—breaking down more readily under a stream of water which is allowed to fall upon them—or they may be completely diffuent. They are never distinctly circumscribed, but pass by insensible gradations into the neighboring tissue.

Under the microscope the change is seen to consist in a disintegration of the nerve-tissue. The white substance of the fibres first coagulates, then breaks up into large masses (myeline), and these subsequently undergo fatty metamorphosis. The cells of the neuroglia, the small bloodvessels, and when the gray matter is implicated, the large nerve-cells, are also involved in the fatty change. The tissue is thus converted into broken-down fibres, a large amount of molecular fat, and numerous large granular corpuscles, the so-called "exudation corpuscles" or "compound inflammatory globules of Gluge." (Fig. 11.) These corpuscles, as already stated, were formerly looked upon in all cases as the result of inflammation, hence their name; they are, however, simply conglomerations of fat granules formed by the degeneration of the cellular elements. (See Fig. 9.) The cells from which they originate are, according to Virchow and Robin, the cells of the neuroglia—the

FIG. 11.



Chronic White Softening of the Brain. Showing the granular corpuscles, broken-down nerve-fibres, and fat granules, of which the softened substance is composed. One or two nucleated cells (probably nerve-cells) are also visible. $\times 250$.

connective tissue of the brain:—these cells share in the fatty change, and in doing so, appear to undergo considerable enlargement before they are destroyed and the fatty matter breaks up. They vary in size from $\frac{1}{800}$ to $\frac{1}{2000}$ inch in diameter, the average being $\frac{1}{1000}$. When the softening affects the gray matter the nerve-cells also contain molecular fat. The small arteries and capillaries running through the softened part are many of them filled with fat granules and granular cells. These latter probably originate in the white blood-corpuscles which have accumulated in the part and undergone fatty changes. Molecular fat and the large granular corpuscles will also be seen adhering to the external surface of the vessels; and here care is required to distinguish these from fatty degeneration of the vessels themselves, with which the cerebral softening is so frequently associated. As the process proceeds the cerebral substance is completely destroyed and all trace of nerve-structure is ultimately lost.

The color of the softened portion varies considerably. It may resemble that of the surrounding healthy tissue; in other cases it is altered to a yellowish or deep red tint. According to these variations in color, cerebral softenings have been classified into *white*, *yellow*, and *red*. The color depends in great measure upon the amount of blood contained in the part, and on this account is important, as indicating the manner in which the softening has been brought about.

White Softening.—This is, in the great majority of cases, a chronic condition. It occurs especially in old people, and is usually due to that disease of the smaller cerebral bloodvessels and consequent interference with the circulation which is common as a result of age. The impairment of the contractile power of the heart must also constitute an auxiliary in the causation of the imperfect vascular supply. It is the gradual manner in which the supply of blood is diminished which accounts for the absence of hyperæmia or hemorrhage, so that the color of the softened portion either resembles that of healthy brain-tissue, or is an opaque dirty white. White softening is sometimes acute, in which case it is usually due to the sudden obstruction of the circulation by the impaction of an embolus in one of the *larger* arteries. (See “Embolism in the Brain.”)

Yellow Softening.—This is, in most cases, simply a variety of the former process, in which, from the fine state of division and close aggregation of the fatty particles, a dead yellowish-white color is imparted to the softened tissue. This color is probably often partly owing to the presence of altered blood pigments, the result of some

previous slight extravasation. The pigment may sometimes be seen as fine dark granules, scattered through the cells of the neuroglia and the nerve-cells of the gray matter, where at first sight they look like fatty particles; they are distinguished, however, by their dark black color. A softening of the brain more rapidly induced, as by embolism to thrombosis, may also occasionally be of a yellow color. This, however, is only the case when the softened portion has attained a certain age, and much of the extravasated blood has been removed by absorption. Lastly, a condition of gelatinous œdema of a yellow color, which has been described by Rokitansky as often being present in the immediate vicinity of cerebral tumors, has been regarded as a variety of yellow softening.

Red Softening.—This is commonly a more acute affection, most frequently dependent upon vascular obstruction, either from embolism or thrombosis. There is collateral hyperæmia, rupture of capillaries, and extravasation of blood; the softened tissue is consequently of a deep red color. These forms of softening will be described in the chapter on “Embolism.” Red softening is also sometimes associated with the chronic white variety, some of the diseased vessels giving way, and thus extravasation of blood taking place into the already softened tissue. Lastly, red softening may be inflammatory. (See “Inflammation of the Brain.”)

Fatty Degeneration of Muscle.

Both striated and non-striated muscle may be the seat of fatty degeneration. In the latter, the muscular fibre-cells are the seat of the change; they become filled with fat granules and are ultimately destroyed. This condition is frequently met with in the middle coat of arteries which are undergoing fatty degeneration.

In striated muscle—both in the voluntary and in the involuntary of the heart—the fibres themselves are the seat of the morbid process, which consists in the conversion of the albuminous matter of which the fibre is composed into fat. The earliest stage of the affection is characterized by an indistinctness in the transverse markings of the fibres, which in many parts become studded with minute particles of fat. (Fig. 12.) These gradually increase in number and size, and are usually dis-

FIG. 12.



Fatty Degeneration of Muscular Fibres of Heart. a. Earlier stage. b. More advanced. $\times 400$.

tributed somewhat irregularly within the sarcolemma. In some parts single rows of granules are found running along the length of the fibre; in others, they are grouped around the nuclei or arranged in transverse lines corresponding with the striæ of the muscle. The fibres become extremely friable, and are readily broken up into short fragments. As the process proceeds the transverse markings entirely disappear, and nothing but molecular fat and oil globules are seen within the sarcolemma. The sarcolemma itself may ultimately be destroyed, and nothing remain of the original fibre but the fatty débris into which its albuminous constituents have been converted. This is true fatty degeneration of muscle; in it the muscular elements are destroyed, and it thus differs essentially from fatty *infiltration*, in which there is simply a development of fat between the fasciculi, the fasciculi themselves not being primarily affected. (See "Fatty Infiltration of Muscle.")

Fatty Degeneration of the Heart.

It is in the heart that fatty degeneration of muscle is most frequently met with, and here it assumes a most important aspect from the deleterious influence which it exercises upon the motor power of the organ. The muscular substance may be affected throughout, or the degeneration may be confined to certain portions of it. It is more common in some situations than in others. The order of frequency with which it occurs in different parts is, according to Dr. Quain, firstly, the left ventricle; secondly, the right ventricle; thirdly, the right auricle; and fourthly, the left auricle.


The wider the extent of tissue that is affected, the less advanced, as a rule, is the degree of the degeneration. It is in those cases in which small tracts of tissue only are involved that the process is met with in its most advanced stage.

The consistence of the degenerated part is always diminished, and its color altered. When the change is slight, and more or less general, the muscle is somewhat softer and more flabby than natural; it is more friable, and often breaks with a soft granular fracture. The color is uniformly rather paler, and more opaque than that of healthy cardiac tissue. Under the microscope the muscular fibres are seen to have lost their striated appearance, and to contain granules of fat.

This diffuse form of degeneration may occur in the course of those diseases in which the oxidation processes are reduced to a minimum;

in all those diseases, in short, which are attended by marked anæmia, whether this anæmia be gradually or rapidly induced. In the case from which the accompanying drawing was taken the degeneration was acute. (Fig. 13.) This was the case of a weakly young girl

FIG. 13.



Acute Fatty Degeneration of Heart and of other Muscles. a. Heart. b. Rectus abdominis. The whole of the heart-tissue was affected, and also the muscles in other parts of the body. X 400.

who was under my care, suffering from slight valvular disease. She quickly succumbed with acute fatty degeneration of the heart and other muscles, which was induced by a profuse loss of blood during a menstrual period, and by inability to retain food.¹ In its slightest degrees a diffuse fatty degeneration of the heart sometimes occurs in the course of acute febrile diseases. This will be again referred to in the chapter treating of the histological changes produced by pyrexia.

Less frequent than this diffuse and comparatively slight degree of degeneration, affecting the whole or the greater part of the organ, is a condition in which, although perhaps the change may be more or less general, it is much more advanced in some parts than in others. In such cases the heart presents a mottled appearance; numerous opaque pale yellowish or brownish patches are seen irregularly distributed throughout its substance. These patches vary considerably in size

¹ This case is reported in "Trans. Clinical Society, Lond.," vol. viii. 1875.

and form; they are met with especially in the papillary muscles, the columnæ carneæ, and in the layers of fibres immediately beneath the endocardium. They may also occur beneath the pericardium, and in the deeper portions of the organ. They correspond with the most degenerated portions of the tissue. They are soft and flabby, and have a rotten consistence, tearing readily under the finger. Under the microscope, the fibres are often seen to be in an advanced stage of fatty degeneration; their sarcolemma contains large molecules of fat and oil globules, which in many parts have escaped and lie free amongst the surrounding less degenerated tissue. (Fig. 12, *b*.) These more localized degenerations are most common in old people, and usually result from imperfect supply of blood due to disease of the coronary bloodvessels, and not from conditions of general anæmia. The peripheral layers of muscular tissue also frequently undergo extensive fatty degeneration as the result of pericarditis. The connection between these localized degenerations and rupture and aneurism of the heart is well known.

Brown Atrophy of the Heart.

Closely allied to and occasionally associated with fatty degeneration of the heart, is the condition known as brown atrophy. This consists in a gradual atrophy of the muscular fibres, together with the formation of granules of brownish-yellow or blackish pigment. These granules of pigment are either grouped in clusters around the nuclei, or more generally distributed within the fibre. The fibres are frequently, at the same time, the seat of more or less fatty degeneration. (Fig. 14.)

FIG. 14.



Brown Atrophy of the Heart. Showing the granules of pigment and the atrophy of the fibres. The latter have in some parts undergone slight fatty metamorphosis. $\times 400$.

Brown atrophy of the heart is rare. It usually occurs as a senile change, or as a part of general marasmus from other causes. It is also met with in some cases of cardiac hypertrophy. It is more chronic in its course than simple fatty degeneration, and its recognition is in most cases impossible without the aid of the microscope.

Fatty Degeneration of the Kidneys.

Fatty degeneration of the kidneys frequently occurs as a result of inflammation of the organs. This *secondary* degeneration will be

alluded to when treating of renal inflammations. Primary fatty degeneration is much less frequent. It must be borne in mind that the renal epithelium very commonly contains more or less fat; but it is only when this is excessive that it can be regarded as a diseased condition. This excessive formation of fat in the kidney is, I think, less common than is generally supposed. It is, however, occasionally met with in chronic diseases, especially in pulmonary phthisis. It is also a result of poisoning by phosphorus.

In simple fatty degeneration the change is usually confined to the epithelium of the cortex. The cortex presents on section a somewhat yellowish-white surface, often slightly mottled, and this, in most cases, is most marked near the bases of the pyramids. There is no adhesion of the capsule or granulation of the surface. This change appears to interfere but little, if any, with the functions of the organs, and in this respect it resembles the analogous change in the liver. It is not usually accompanied by albuminuria.

CHAPTER VI.

MUCOID AND COLLOID DEGENERATION.

UNDER this head is included a class of morbid changes which are characterized by a peculiar softening of the tissues. Colloid and mucoid degeneration have frequently been described under the common term of "colloid softenings," but, although they are very closely allied and sometimes associated, they appear to constitute two distinct processes: the former affecting more especially the cells, the latter the intercellular substance.

MUCOID DEGENERATION.—This consists in the transformation of the albuminoid constituents of the tissues into *mucin*, owing to which they become converted into a material of a soft, mucilaginous, jelly-like consistence. This is the condition of nearly all tissues in their immature or foetal state: the connective tissues in the foetus consist almost entirely of this soft mucin-yielding substance. Some tissues retain

these characters after birth. The umbilical cord, and the vitreous humor of the eye, are both composed of this substance.

A mucoid change occurs as a physiological process in the secretion of mucus. The newly-formed cells undergo mucoid transformation, which results in their destruction, and the mucus is thus liberated. It is considered probable by some that the cells may also evacuate the mucus without being destroyed. This process occurs much more rapidly in all catarrhal conditions of the mucous membrane.

As a pathological process mucoid degeneration affects especially the intercellular substance. The intercellular substance of the connective tissues in their fully developed state consists of gelatin and chondrin, and the mucoid change is thus a reversion of this substance to its foetal condition.

Mucin is closely allied to albumen, more so than to either gelatin or chondrin: it differs from it in not containing sulphur. Like albumen, it is only met with in alkaline fluids—being held in solution by the free alkali—from which it is precipitated by dilute acetic acid. It differs from albumen in being insoluble in an excess of the acid, and also in not being precipitated by boiling, by tannin, or by bichloride of mercury. Its behavior with these two reagents will also distinguish it from gelatin and chondrin, which are both precipitated by them.

The mucoid change is by no means a common one. It is most frequently met with in cartilage, especially in the inter-vertebral and costal cartilages of old people. It also occurs in serous membranes, in bone, and in many of the new formations, especially in those of the connective tissue class. Wherever it occurs it produces softening of the affected parts; which are transformed into a homogeneous, colorless material, of a soft mucilaginous jelly-like consistence. If the change is limited to isolated portions of the tissue, the softened parts surrounded by those which are unaltered, often present the appearance of cysts. These cyst-like formations containing mucoid substance are not uncommonly met with in the costal cartilages and in new growths.

As to the cause of the mucoid change, nothing is known.

COLLOID DEGENERATION.—This differs from the former, inasmuch as it is the cells which are more especially involved in the process.

The change consists in the metamorphosis of the albuminoid constituents of the cells into a substance known as *colloid* material. Colloid closely resembles mucin, but it differs from it chemically, in con-


taining sulphur, and in not being precipitated by acetic acid. It is a colorless, transparent, glistening material, of the consistence of jelly or half-set glue. It makes its appearance within the cells as small lumps, which gradually increase in size, pushing the nucleus to one side, until they completely fill the cell. (Fig. 15.) The cells are thus destroyed, and converted into colloid masses. The small colloid masses subsequently swell up, coalesce, and so form larger masses of firm, transparent, yellowish, jelly-like material, which are readily to be recognized by the naked eye. As the colloid matter increases, and the cells are destroyed, the intercellular substance atrophies or softens, and in this way cyst-like cavities are formed, within which is contained the gelatinous substance. Here it may subsequently undergo a process of liquefaction. (See Fig. 57.)

The colloid change is most common in enlargements of the thyroid gland (goitre), in the lymphatic glands, in the choroid plexus, and in many of the new formations. (See "Colloid Cancer.") Its causes and nature are as obscure as those of the allied mucoid softening.

It is when occurring in new formations that these two forms of degeneration assume their most important aspects. Certain varieties of tumors may originate as mucoid or colloid growths, or may subsequently undergo these morbid transformations. The mucous tumors (myxomata), which resemble in structure the umbilical cord, consist entirely of a gelatinous mucin-yielding substance. The sarcomata, lipomata, enchondromata, and the cancers may also become the seats of these forms of softening.

Such growths have sometimes been described as gelatiniform or colloid *cancers*, this term having been applied to them without any regard to their structure or real nature. Cancers, it is true, may undergo a colloid change (see "Colloid Cancer"); but it is by no means true that all tumors possessing these soft gelatiniform characters are cancers. The terms "mucoid" or "colloid," applied to a new growth, merely imply certain physical and chemical characters, and convey but little information as to its real nature.

FIG. 15.



*Colloid Cells, from
a colloid cancer.
(Rindfleisch.)*

and form; they are met with especially in the papillary muscles, the columnæ carneæ, and in the layers of fibres immediately beneath the endocardium. They may also occur beneath the pericardium, and in the deeper portions of the organ. They correspond with the most degenerated portions of the tissue. They are soft and flabby, and have a rotten consistence, tearing readily under the finger. Under the microscope, the fibres are often seen to be in an advanced stage of fatty degeneration; their sarcolemma contains large molecules of fat and oil globules, which in many parts have escaped and lie free amongst the surrounding less degenerated tissue. (Fig. 12, *b*.) These more localized degenerations are most common in old people, and usually result from imperfect supply of blood due to disease of the coronary bloodvessels, and not from conditions of general anæmia. The peripheral layers of muscular tissue also frequently undergo extensive fatty degeneration as the result of pericarditis. The connection between these localized degenerations and rupture and aneurism of the heart is well known.

Brown Atrophy of the Heart.

Closely allied to and occasionally associated with fatty degeneration of the heart, is the condition known as brown atrophy. This consists in a gradual atrophy of the muscular fibres, together with the formation of granules of brownish-yellow or blackish pigment.

FIG. 14.



Brown Atrophy of the Heart. Showing the granules of pigment and the atrophy of the fibres. The latter have in some parts undergone slight fatty metamorphosis. $\times 400$.

These granules of pigment are either grouped in clusters around the nuclei, or more generally distributed within the fibre. The fibres are frequently, at the same time, the seat of more or less fatty degeneration. (Fig. 14.)

Brown atrophy of the heart is rare. It usually occurs as a senile change, or as a part of general marasmus from other causes. It is also met with in some cases of cardiac hypertrophy. It is more chronic in its course than simple fatty degeneration, and its recognition is in most cases impossible without the aid of the microscope.

Fatty Degeneration of the Kidneys.

Fatty degeneration of the kidneys frequently occurs as a result of inflammation of the organs. This *secondary* degeneration will be

alluded to when treating of renal inflammations. Primary fatty degeneration is much less frequent. It must be borne in mind that the renal epithelium very commonly contains more or less fat; but it is only when this is excessive that it can be regarded as a diseased condition. This excessive formation of fat in the kidney is, I think, less common than is generally supposed. It is, however, occasionally met with in chronic diseases, especially in pulmonary phthisis. It is also a result of poisoning by phosphorus.

In simple fatty degeneration the change is usually confined to the epithelium of the cortex. The cortex presents on section a somewhat yellowish-white surface, often slightly mottled, and this, in most cases, is most marked near the bases of the pyramids. There is no adhesion of the capsule or granulation of the surface. This change appears to interfere but little, if any, with the functions of the organs, and in this respect it resembles the analogous change in the liver. It is not usually accompanied by albuminuria.

CHAPTER VI.

MUCOID AND COLLOID DEGENERATION.

UNDER this head is included a class of morbid changes which are characterized by a peculiar softening of the tissues. Colloid and mucoid degeneration have frequently been described under the common term of "colloid softenings," but, although they are very closely allied and sometimes associated, they appear to constitute two distinct processes: the former affecting more especially the cells, the latter the intercellular substance.

MUCOID DEGENERATION.—This consists in the transformation of the albuminoid constituents of the tissues into *mucin*, owing to which they become converted into a material of a soft, mucilaginous, jelly-like consistence. This is the condition of nearly all tissues in their immature or foetal state: the connective tissues in the foetus consist almost entirely of this soft mucin-yielding substance. Some tissues retain

these characters after birth. The umbilical cord, and the vitreous humor of the eye, are both composed of this substance.

A mucoid change occurs as a physiological process in the secretion of mucus. The newly-formed cells undergo mucoid transformation, which results in their destruction, and the mucus is thus liberated. It is considered probable by some that the cells may also evacuate the mucus without being destroyed. This process occurs much more rapidly in all catarrhal conditions of the mucous membrane.

As a pathological process mucoid degeneration affects especially the intercellular substance. The intercellular substance of the connective tissues in their fully developed state consists of gelatin and chondrin, and the mucoid change is thus a reversion of this substance to its foetal condition.

Mucin is closely allied to albumen, more so than to either gelatin or chondrin: it differs from it in not containing sulphur. Like albumen, it is only met with in alkaline fluids—being held in solution by the free alkali—from which it is precipitated by dilute acetic acid. It differs from albumen in being insoluble in an excess of the acid, and also in not being precipitated by boiling, by tannin, or by bichloride of mercury. Its behavior with these two reagents will also distinguish it from gelatin and chondrin, which are both precipitated by them.

The mucoid change is by no means a common one. It is most frequently met with in cartilage, especially in the inter-vertebral and costal cartilages of old people. It also occurs in serous membranes, in bone, and in many of the new formations, especially in those of the connective tissue class. Wherever it occurs it produces softening of the affected parts; which are transformed into a homogeneous, colorless material, of a soft mucilaginous jelly-like consistence. If the change is limited to isolated portions of the tissue, the softened parts surrounded by those which are unaltered, often present the appearance of cysts. These cyst-like formations containing mucoid substance are not uncommonly met with in the costal cartilages and in new growths.

As to the cause of the mucoid change, nothing is known.

COLLOID DEGENERATION.—This differs from the former, inasmuch as it is the cells which are more especially involved in the process.

The change consists in the metamorphosis of the albuminoid constituents of the cells into a substance known as *colloid* material. Colloid closely resembles mucin, but it differs from it chemically, in con-


taining sulphur, and in not being precipitated by acetic acid. It is a colorless, transparent, glistening material, of the consistence of jelly or half-set glue. It makes its appearance within the cells as small lumps, which gradually increase in size, pushing the nucleus to one side, until they completely fill the cell. (Fig. 15.) The cells are thus destroyed, and converted into colloid masses. The small colloid masses subsequently swell up, coalesce, and so form larger masses of firm, transparent, yellowish, jelly-like material, which are readily to be recognized by the naked eye. As the colloid matter increases, and the cells are destroyed, the intercellular substance atrophies or softens, and in this way cyst-like cavities are formed, within which is contained the gelatinous substance. Here it may subsequently undergo a process of liquefaction. (See Fig. 57.)

The colloid change is most common in enlargements of the thyroid gland (goitre), in the lymphatic glands, in the choroid plexus, and in many of the new formations. (See "Colloid Cancer.") Its causes and nature are as obscure as those of the allied mucoid softening.

It is when occurring in new formations that these two forms of degeneration assume their most important aspects. Certain varieties of tumors may originate as mucoid or colloid growths, or may subsequently undergo these morbid transformations. The mucous tumors (myxomata), which resemble in structure the umbilical cord, consist entirely of a gelatinous mucin-yielding substance. The sarcomata, lipomata, enchondromata, and the cancers may also become the seats of these forms of softening.

Such growths have sometimes been described as gelatiniform or colloid *cancers*, this term having been applied to them without any regard to their structure or real nature. Cancers, it is true, may undergo a colloid change (see "Colloid Cancer"); but it is by no means true that all tumors possessing these soft gelatiniform characters are cancers. The terms "mucoid" or "colloid," applied to a new growth, merely imply certain physical and chemical characters, and convey but little information as to its real nature.

FIG. 15.



Colloid Cells, from a colloid cancer. (Rindfleisch.)

Muscular Changes in Typhoid Fever.

Allusion must be made here to that change met with in the muscles in typhoid fever, which was first described by Zenker, and which appears to be somewhat allied to colloid degeneration. This change is most marked in the recti muscles of the abdomen, the abductors of the thigh, and in the diaphragm. The portions of muscle affected are paler than natural, of a reddish-gray or brownish-yellow color, and abnormally friable. Under the microscope, the altered fibres are much swollen, the transverse striation is lost, and the sarcolemma is

occupied by a homogeneous, structureless material. This material is exceedingly brittle, and, as usually seen, presents a wrinkled appearance, or is broken up transversely into several irregular fragments. (Fig. 16.) The fibres are never universally affected, but many normal are associated with the altered elements. This change necessarily impairs the contractile power of the muscle, and it often leads to rupture of some of the fasciculi and hemorrhage. The new material appears to be readily absorbed, and the lost fibres to be quickly regenerated.




FIG. 16.
A Portion of the Soleus Muscle from a Case of Typhoid Fever. Preparation teased after treatment with Müller's Fluid. $\times 200$. Reduced $\frac{1}{2}$.

With regard to the nature of the change but little is known. Although most frequent in typhoid, it occurs also occasionally in other severe febrile diseases. The transformed muscle, as already stated, appears to be allied to colloid. A similar change is described by Cornil and Ranvier as sometimes occurring in muscles in the neighborhood of abscesses, of inflamed bone, and of tumors.

CHAPTER VII.

AMYLOID DEGENERATION.

AMYLOID degeneration, which is one of the most important of the degenerative processes, consists in the infiltration of the tissues with a peculiar homogeneous translucent substance, closely resembling albumen, by which their vitality becomes diminished and their functions impaired.

It is often known as the *lardaceous*, or *waxy* change, the organs affected by it having somewhat the appearance of lard or wax. The term "*amyloid*"—which was applied to it by Virchow, from the supposed resemblance of the new material to cellulose or starch—as being that by which it is perhaps most generally known, is here adopted.

This form of degeneration is very rarely a primary affection, but almost invariably occurs as the sequel of some other disease. It is in those diseases which are attended by profuse and long-continued suppuration, such as chronic diseases of bone, empyema, chronic disintegrative diseases of the lungs, chronic pyelitis, and chronic intestinal ulceration, that the amyloid change is most frequently met with. It also occurs in the advanced stages of constitutional syphilis, but especially in those cases in which there is chronic bone disease or chronic ulceration.

Every organ and tissue may be the seat of the change; those, however, in which it is especially prone to occur are the *kidneys*, the *liver*, the *spleen*, the *lymphatic glands*, and the *intestines*. It is met with less frequently in the stomach, in the supra-renal capsules, in the pharynx, and œsophagus, in the bladder, prostate, and generative organs, in serous membranes, in the membranes of the brain and cord, and in muscle. It is rarely limited to one organ, but several organs are almost invariably simultaneously affected by it.

Respecting the nature of the new material which infiltrates the tissues, the analyses of Kekulé and Schmidt show that it is a nitrogenous substance closely allied to albumen. The conclusions arrived at by these observers are, however, not satisfactory, as they were unable completely to separate the substance from the tissues in which it was

deposited. More recently, Kühne succeeded in more completely isolating it. He submitted the affected organs to a process of artificial digestion, and inasmuch as the amyloid substance is not dissolved by digestion with pepsin, it was thus obtained free from the tissues in which it was contained. The results of Kühne's analyses are very similar to those of Kekulé and Schmidt. Dr. Dickinson regards the new substance as fibrin deprived of its alkaline salts. The investigations of Dr. Marcet¹ show that the infiltrated organs are considerably deficient in potash and phosphoric acid, whilst they contain an excess of soda and chlorine. In conclusion it may be stated that, although the precise composition of the amyloid substance has not yet been determined, the results of the several analyses appear to justify the opinion that it is some modification of fibrin.

The most characteristic feature of the amyloid substance is the peculiar reaction which it gives with iodine, and with iodine and sulphuric acid. If an aqueous solution of iodine—made with the help of potassium iodide—be applied to an amyloid organ, the affected portion changes to a deep reddish-brown color. This is not permanent, but gradually passes off, and the part regains its former appearance. If the application of the iodine be followed by the cautious addition of sulphuric acid, a blackish-blue or violet tint is produced. This latter reaction, however, is not easily obtained, considerable nicety being required in the application of the reagents. The following is the method for obtaining it, recommended by Professor Virchow: A dilute aqueous solution of iodine must be allowed to soak well into the tissue, the excess must be poured off, and a single drop of concentrated sulphuric acid gradually added, when a blue or violet color will be produced, either at once or after some time. In the hands of English pathologists this latter reaction has certainly met with but little success; and if the color be obtained, it is by no means satisfactory, and more nearly resembles a black than the blue which has been described. Fortunately, however, the reaction with iodine alone is perfectly characteristic, and the attempt to obtain the blue by the subsequent addition of sulphuric acid is therefore quite unnecessary. If the change is at all advanced, the reddish-brown color will be produced by merely pouring the aqueous solution of iodine over the cut surface of the organ; but in slighter degrees of the affection, thin sections must be

¹ See "Report of Committee on Lardaceous Disease," Trans. Path. Soc. Lond. 1871.

made with a Valentin's knife, and well washed with water to remove the blood, before the coloration with iodine can be obtained.

Upon considering the above reactions, it will be seen that they resemble, in some respects, those exhibited by some well-known organic substances. Cellulose—the substance which forms the external membrane of vegetable cells—yields a blue when treated with iodine and sulphuric acid, and so far resembles the amyloid material; but it differs in not being colored by iodine alone. The same is true of cholesterine; this also gives a reddish-brown with sulphuric acid, but is not altered by iodine. To starch the resemblance is not so marked as to the two preceding substances; with this, iodine produces a blue color without the addition of the acid.

The amyloid substance almost invariably makes its appearance first in the small arteries and capillaries, and subsequently extends from them to the surrounding tissues. The cells of the intima and of the muscular coat are the first to become infiltrated, then the remaining structures of the artery. When the vessels have become infiltrated, the amyloid matter extends to the immediately surrounding parts; it invades both the cells and the intercellular substance, and may gradually extend until the whole organ is completely impregnated.

The changes produced in the tissues by this infiltration are very characteristic. The cells as they become filled with the new material gradually increase in size; they lose any irregularities in their contour, and become rounder and more regular in shape; their nuclei disappear, and the whole cell is converted into a structureless homogeneous body, which has a peculiar translucent glistening appearance. (Fig. 17, *a*.) If the cells are in close contact many of them may coalesce, and their distinctive boundaries thus become obliterated. (Fig. 17, *b*.) The intercellular substance in the same way acquires a homogeneous glistening appearance. The walls of the small arteries—in which, as already stated, the change commences—become considerably thickened, the cells of the muscular coat enlarge and ultimately coalesce, the calibre of the vessel becomes diminished, and the circulation through it is consequently impeded. (See Fig. 19.)

Organs in which this change is at all advanced, present features so




FIG. 17.
Liver Cells infiltrated with the Amyloid Substance. a. Single cells. b. Cells which have coalesced. X 300. (Rindfleisch.)

characteristic that its nature can be readily recognized by the naked eye. They are usually considerably increased in size; their absolute weight is increased, and also their specific gravity; their surface is smooth, and the capsule tense and stretched; their consistence is firm and somewhat elastic. On section they exhibit a peculiar homogeneous, glistening, translucent appearance, somewhat resembling wax or glue. Owing to the diminished calibre of their bloodvessels and to the pressure exercised by the new material, they contain but little blood, and hence are always pale in color. The change may involve the whole organ, or it may be limited to certain portions. In the spleen, for example, it is frequently limited to the Malpighian corpuscles; and in the liver to the cells in the more immediate vicinity of the hepatic artery.

Although the above characters are often sufficiently marked, they should always be confirmed by the application of iodine to the cut surface of the organ. In slighter degrees of the affection—when the physical characters are but little altered—the application of this reagent becomes necessary in order to discover the presence of the new substance. In these cases, merely pouring the solution over the organ will often fail to produce the characteristic staining; and it will be necessary to make thin sections with a Valentin's knife, and wash them thoroughly with water to remove the blood, before the reaction can be obtained. In the earliest stages of the change the use of the microscope may be necessary for its recognition.

The effect of amyloid degeneration is to impair or even to completely destroy the nutrition and function of those organs which are affected by it. This is owing to two causes—the obstruction offered to the circulation and the direct influence of the new material upon the parenchyma of the organ. The obstruction to the circulation, which results partly from the diminution in the calibre of the small arteries, and partly from the general pressure exercised by the infiltrated substance, causes an insufficiency in the supply of arterial blood. As a consequence of this, secondary atrophic changes are induced in various parts; the cells may atrophy and undergo fatty metamorphosis, which indeed is frequently associated with the amyloid change. The vitality of the cellular elements is still further impaired by their infiltration with the new material, and thus all manifestation of their functions may ultimately cease.

As this form of degeneration is almost invariably secondary to some grave constitutional state, it can rarely be looked upon as in

itself a cause of death, although it may materially hasten, and even determine, the fatal termination.

Having thus described the nature of the amyloid substance, and the way in which it makes its appearance in the several tissues of an organ, it remains to consider the source from which it is derived. The disease, as already stated, appears to be an infiltration, and to consist in the deposition of a new material from the blood in the various tissues and organs. The existence, however, of any albuminoid substance in the blood which resembles amyloid in its reaction with iodine, has never been made out, even in the most marked cases of the disease. This fact must therefore negative the supposition that it is a simple infiltration. If the new material be derived from the blood at all, it must undergo some chemical change subsequently to its deposition in the tissues. Dr. Dickinson considers that it is dealkalized fibrin, which is deposited in consequence of the loss of the alkali which it normally contains. This loss of alkali he attributes to the chronic suppuration which usually precedes the disease, pus containing large quantities of potassium and sodium salts. He consequently terms the disease "*depurative infiltration*."¹ Whether this be so or no, it is probable that some abnormal condition of the blood is the cause of the disease, and that the change itself consists in the retention and coagulation in the tissues of some albuminoid substance which is derived from the nutritive fluids. The way in which the several tissues of an organ are affected, the change almost invariably commencing in the small nutrient bloodvessels, and extending from them to the surrounding parts; the general character of the affection, several organs being simultaneously involved; together with the fact that it is almost invariably secondary to chronic suppurative diseases or to syphilis—point to some alteration in the composition of the blood as the cause of the degeneration.

Amyloid Degeneration of the Liver.


The liver is one of the most frequent seats of the amyloid change, and here, as in other parts, it probably commences in the small nutrient bloodvessels, although the alterations are much the most marked in the hepatic cells. The minute branches of the hepatic artery are first affected, and from these the infiltration gradually extends to the

¹ "Medico-Chirurgical Transactions," vol. i.

hepatic cells immediately adjacent, until ultimately the whole organ may become involved.

If a liver be examined in the earlier stages of the affection, and the iodine solution applied to thin washed sections of the organ, it will be found that the characteristic staining is limited to certain portions of the lobules—viz. to those which are situated between their external and central parts. This intermediate portion corresponds with the distribution of the hepatic artery, and the ramifications of this vessel together with the hepatic cells situated in their vicinity are the first to become affected. (Fig. 18.) As the change advances the whole lobule may ultimately become involved. The alterations in the hepatic cells are very characteristic. They are much enlarged, irregular in outline, their nuclei are imperceptible, and many of them are fused together into irregular-shaped masses. (See Fig. 17.)

FIG. 18.



Amyloid Liver. A portion of one lobule, showing the enlargement and fusion of the hepatic cells, and the greater implication of the intermediate zone of the lobule. At the more external portion of the lobule are seen several fat cells, a certain amount of fatty infiltration being associated with the amyloid change. v. Hepatic vein. $\times 103$.

The earliest seat of the amyloid change thus differs from that of the fatty. In fatty infiltration it is the most external portion of the lobule in which the fat first accumulates—that which corresponds with the distribution of the portal vein. (See Fig. 8.) It differs also from that pigmentation of the hepatic cells resulting from mechanical congestion, which takes place in the most central portion, around the hepatic vein. (See “Nutmeg Liver.”) Thus in each hepatic lobule three zones may be distinguished:—an external one, which is the chief seat of the fatty change; a central one, which is the chief seat of the

pigmentary change ; and an intermediate one, which is the chief seat of the amyloid change. These three zones, indeed, may frequently be recognized by the naked eye, the pale opaque external one contrasting strongly with the intermediate one, which is translucent, and with the darker central one. In the most advanced stages of the disease, however, both the external and central portions of the lobule may become involved, and the cut surface present an almost uniformly homogeneous appearance.

The amyloid liver is increased in size, often very considerably so ; it may be so large as almost completely to fill the abdominal cavity. The enlargement is uniform, and hence the natural configuration of the organ is but little altered. Its weight is increased, and also its specific gravity. Its edge is rounded, the surface is smooth, and the capsule appears tense and stretched. The consistence is firm and elastic. The cut surface is dry, bloodless, smooth, translucent, and waxy-looking, and of a pale reddish-gray or dirty yellow color. If the change is very far advanced, the tissue may be perfectly homogeneous, all distinction between the individual lobules being lost. In other cases the lobules are distinctly mapped out ; they are enlarged, and the external zone may be of an opaque, yellowish-white color, owing to the presence of fat. This association of the fatty and amyloid changes is exceedingly common. Amyloid infiltration does not obstruct the portal circulation, and hence does not cause ascites. It impairs the vitality of the hepatic cells, and thus interferes with the functions of the organ.


Amyloid Degeneration of the Kidneys.

The kidneys are very liable to be involved in the amyloid change, and here it is the smaller bloodvessels which are more especially affected. It may constitute in them the primary lesion, or it may occur subsequently to inflammatory conditions, implicating the secreting and interstitial structures. As a primary change it is an important variety of Bright's disease.

The process commences in the tufts of vessels which form the Malpighian bodies, the walls of which become thickened by the infiltration of the new material, so that the tufts are increased in size. It then proceeds to the small afferent arteries, and ultimately to the vasa efferentia, and to the arteriolæ rectæ which run through the medullary portion of the organ. The changes produced in the vessels

are very characteristic. Their walls are considerably thickened, and their calibre is so much diminished that the smallest ones cannot be artificially injected. This thickening of the walls of the vessels is mainly owing to alterations in their muscular coat, and especially to the cells of the circular muscular layer. These cells are much increased in size, they are more or less globular in shape, and many of them have lost their distinctive outlines. The longitudinal muscular fibres and the most internal coat of the vessel are often seen as one homogeneous, glistening, structureless layer. (Fig. 19.) After the

FIG. 19.



Amyloid Degeneration of a Malpighian Tuft and small Artery of the Kidney. Showing the thickening of the walls of the vessel, the enlargement of the cells of the circular muscular coat, and the homogeneous layer formed by the intima and longitudinal muscular fibres. $\times 200$, reduced $\frac{1}{2}$.

vessels have become infiltrated, and before those in the pyramids are affected, the deposition extends to the intertubular tissues of the cortex, into which the amyloid substance is poured, imbedding the tubuli uriniferi in its course. In some parts it appears that the exudation takes place into the tubes themselves, as the casts which they contain are occasionally deeply stained by the iodine solution: this, however, is far from being frequently the case. In the earlier stages of the process—if the organ is not the seat of any other morbid change—the tubes and their lining epithelium present a perfectly natural appearance. Many of them contain pale hyaline casts, which also appear in the urine. These, however, are probably simply exudation products; although from the reaction they occasionally exhibit, it appears that they sometimes consist of the same material as that which permeates the vessels and intertubular structures. As the change proceeds, and the new material increases in amount, the tubes become compressed,

and in many places completely obstructed. If the compression is not uniform, they may dilate and form small cysts. The epithelium, which was at first normal, owing to the interference with its nutrition, ultimately atrophies and undergoes fatty changes. In some cases it appears to be the seat of a catarrhal process, and the tubes are found blocked with the epithelial products. In the later stages of the process there is almost invariably an increase in the intertubular connective tissue.

The first effect of this change is to obstruct the circulation in the cortex. The bloodvessels, diminished in calibre, allow little but the liquor sanguinis to pass through them, the passage of the blood-corpuscles being to a great extent prevented: hence the pallor of this portion of the organ. The arterial walls are so altered that fluids and albumen readily permeate them; and thus is produced the large quantity of urine, loaded with albumen, which characterizes the earlier stages of this affection. As the infiltration increases, and the tubes become obstructed, the urine diminishes in quantity. The excretion of urea is less interfered with than in other forms of Bright's disease, and hence symptoms due to its retention seldom occur. Tube casts are rarely numerous; they are for the most part hyaline or finely granular, though sometimes they are covered with fatty epithelium.


In the earlier stages of the affection, the cortex of the kidney is merely rather paler than natural, and perhaps somewhat firmer in consistence; but otherwise it presents no abnormal appearance. It is only upon the application of iodine to the cut surface, or to thin washed sections of the organ, that its diseased condition becomes evident. When this test is employed, the Malpighian bodies at once become apparent as minute red points scattered through the cortex. As the disease advances, the size of the organ increases; the enlargement, however, is principally confined to the cortex. The surface is smooth, and the capsule separates readily. The enlarged cortex is remarkably pale and anæmic, and has a peculiar translucent, homogeneous, wax-like appearance. Its consistence is hard and firm. A few scattered vessels may be seen on the surface, and the bases of the pyramids sometimes exhibit an increased amount of vascularity. If iodine be poured over the cut surface, the Malpighian bodies and the arteries of the cortex become mapped out almost as clearly as in an artificial injection. The enlarged Malpighian bodies may indeed usually be seen as glistening points before the iodine is applied. Frequently, the homogeneous appearance of the cortex is interrupted by minute,

opaque, yellowish-white lines and markings; these are produced by the fatty changes in the epithelium of the tubes, which so commonly occur in the later stages of the affection. Ultimately the capsule becomes more or less adherent, and slight irregular depressions make their appearance upon the surface of the organ: the latter are due to atrophic changes in some of the tubes. If, as is usually the case, the infiltration is associated with an increase in the intertubular connective tissue, the atrophy of the organ will be more marked. (See "Interstitial Nephritis.")

Amyloid Degeneration of the Spleen.

Amyloid degeneration of the spleen is met with in two forms—one in which the disease is limited to the Malpighian corpuscles ("Sago Spleen"), and the other in which the pulp appears to be chiefly implicated. The former is much the more common condition. In it the Malpighian corpuscles become infiltrated with the amyloid substance and converted into translucent wax-like bodies, much like boiled sago; hence its name. The process commences in the small arteries of the corpuscle, and it is only after these have become infiltrated that it extends to the corpuscles themselves. The small lymphatic cells of which the corpuscle is made up are then involved in the process, they increase in size, become more irregular in outline, many of them coalesce, and ultimately the whole is converted into a pale, firm, translucent glistening mass. (Fig. 20.)

FIG. 20.



Amyloid Degeneration of the Spleen—"Sago Spleen." A portion of one of the infiltrated Malpighian corpuscles *a*, with the adjacent normal splenic tissue *b*. Showing the increase in size and, in many parts, the coalescence of the cells, of which the corpuscle is composed. $\times 200$.

The sago spleen is more or less enlarged; its weight and density are also increased. The cut surface is smooth, dry, and studded all over with small glistening sago-like bodies, varying in size from a

millet to a hemp-seed, which are stained a reddish-brown color by the iodine solution. These may become so large as to occupy a large portion of the organ, although in earlier stages of the affection they are so minute that they can only be seen in thin sections of the tissue.

In the other variety of amyloid spleen, the pulpy parenchyma between the corpuscles is infiltrated with the new material. This is probably merely an advanced stage of the former condition, in which the disease extends from the corpuscles to the surrounding pulp; the whole organ being ultimately involved. Under these circumstances the organ often attains a considerable size, much larger than is met with in the sago spleen. It is remarkably hard and firm, and the capsule is tense and transparent. On section it presents a dry, homogeneous, translucent, bloodless surface, of a uniform dark reddish-brown color. Thin sections can be readily made with a knife, the organ cutting like soft wax. The corpuscles are not visible as in the former variety, being probably obscured by the surrounding pulp, which has become so densely infiltrated with the amyloid substance.

Amyloid Degeneration of Lymphatic Glands.

In the lymphatic glands the process much resembles that in the spleen. The small arteries in connection with the follicles of the gland are the earliest seats of the change; and from these it extends to the lymphoid cells. The follicle thus becomes ultimately converted into a small homogeneous mass.

The glands are enlarged, and on section the minute wax-like bodies can often be seen scattered through the cortex. The cut surface is smooth, pale, and translucent.

As these glands are largely concerned in the formation of the blood-corpuscles, their implication in the amyloid change must to a large extent aid in producing the emaciation and anæmia which characterize this affection. The same is true of the spleen, which is usually simultaneously involved.

Amyloid Degeneration of the Alimentary Canal.


The whole of the alimentary tract may be the seat of the amyloid change, and here it assumes an important aspect from the deleterious influence which it exercises upon the absorbent and secreting processes, and from the consequent impairment of the general nutrition which

results. The disease, however, in this situation is very apt to escape observation, as it produces but little alteration in the appearance of the parts. The mucous membrane may look somewhat pale, translucent, and œdematous, but otherwise to the naked eye nothing is discoverable. It is only upon the application of iodine to the washed mucous surfaces that the nature of the change becomes apparent. In the small intestine—which is perhaps the part most commonly affected—the effect of the application of iodine is very characteristic. A number of small reddish-brown points appear over the whole surface of the membrane; these correspond to the intestinal villi, the arteries and capillaries of which are infiltrated with the amyloid substance. In the stomach and œsophagus the vessels are mapped out in a similar manner by the iodine solution. The change in the intestine gives rise to serous diarrhœa, this being probably due to an increased permeability of the infiltrated walls of the vessels.

The Corpora Amylacea.

The corpora amylacea or “amyloid bodies,” so frequently met with in the nervous system, in the prostate, and in other parts, have usually been looked upon as more or less allied to the amyloid substance; there appears, however, with the exception of a certain similarity in their behavior with iodine and sulphuric acid, to be no connection between them.

They are round or oval bodies, formed of a succession of concentric layers, and are often changed to a deep-blue color by iodine, thus bearing both in their structure and chemical properties the strongest resemblance to granules of vegetable starch. (Fig. 21.) Sometimes, however, the blue is only exhibited after the subsequent addition of sulphuric acid, and thus a resemblance is shown to the amyloid substance. They vary in size from microscopic granules, to bodies which are distinctly visible to the naked eye; sometimes being as much as one or two lines in diameter. The larger ones are usually formed by the conglomeration of the smaller granules, which are often inclosed by a common envelope.



They occur especially in conditions of atrophy or softening of the nervous system; the ependyma of the ventricles, the white substance

of the brain, the choroid plexus, the optic nerve and retina, and the spinal cord, being their favorite seats. The larger forms are met with most frequently in the prostate. The prostate of nearly every adult contains some of these bodies; and they may accumulate here to such an extent as to form large concretions. They are occasionally met with in the lungs and in mucous and serous membranes.

As has been said, they usually exhibit a bright-blue color upon the application of iodine alone, although in some cases not until the subsequent addition of sulphuric acid. Many of them, however, are colored green, or even brown by these reagents. The green is due to their admixture with nitrogenous matters, which give a yellow color with iodine, and hence the combination yields a green. The greater the amount of nitrogenous matter the more brown does the color become.

From the laminated structure of these bodies they would appear to be formed by the gradual precipitation of some material, layer by layer, upon the surface of pre-existing particles. The nature of the material, however, does not appear to resemble that of the substance which infiltrates the organs in amyloid degeneration. The two processes are so essentially different, both in the circumstances under which they occur and in the characters and seat of the morbid products, that they cannot be looked upon as in any way analogous. Amyloid degeneration is a general change, in which numerous organs become infiltrated with an albuminoid substance, whereas the formation of the corpora amylacea is evidently of a local nature. The latter is often preceded by those local atrophic changes associated with advanced life, and appears to consist in the deposition of some starch-like material, probably liberated in the tissues themselves, upon any free body which may exist in its vicinity.

The corpora amylacea, especially those occurring in the choroid plexus and in the lateral ventricles, are very liable to become calcified, and they then constitute one form of "brain-sand," which is so often met with in these situations.

CHAPTER VIII.

CALCAREOUS DEGENERATION.

CALCAREOUS Degeneration—or, as it is more commonly called *Calcification*—consists in the infiltration of the tissues with calcareous particles. Physiologically, an infiltration of calcareous particles takes place in the formation of bone, in which lime and magnesian salts are deposited in the fibrous or cartilaginous matrix. This physiological is precisely similar to the pathological process. It is important, however, to distinguish simple calcification from ossification. In the latter there is not only a deposition of lime salts, but an *active* change in the tissue itself—a proliferation of the cellular elements, an intimate union of the calcareous matters with the tissue, and the formation of a true osseous structure in which the calcareous particles are not visible. Calcification, on the other hand, is a purely *passive* process, there is no increased nutritive activity of the part, no multiplication of elements, no alteration of the structure, but merely an infiltration with calcareous particles.

An infiltration and deposition of calcareous substances occur under two opposite conditions; one—in which there is an absolute increase in the amount of these constituents in the blood, and a portion of the excess becomes deposited in the tissues; the other—in which there is no such increase; but the deposition takes place owing to some alteration in the tissue itself.

An absolute increase of the saline constituents in the blood, and the deposition of the excess in the tissues, is much the less frequent form of calcification. It occurs in some forms of softening of bone, especially in extensive caries and osteomalacia. In these diseases the lime salts are removed from the bone, returned into the blood, and some of them deposited in other tissues. In such cases the calcification is usually more or less general—many organs being simultaneously involved. In osteomalacia, it is not uncommon to find the kidneys, the lungs, the stomach, the intestines, and even the dura mater and liver, infiltrated to a greater or less extent with lime salts. The deposition commences in the tissue immediately surrounding the bloodvessels from

which the calcareous matters are derived ; thus, in the lungs, the seat of the change is the interlobular tissue ; in the stomach, the stroma between the glands ; and in the kidney, the tubuli uriniferi and the intertubular tissue. Precisely analogous to this form of calcification is the deposition of the excess of urate of soda which takes place in gout.


In the great majority of cases, however, calcification is a *local* change, depending not upon any alteration in the composition of the blood, but upon changes in the tissues themselves, owing to which some of the saline matters which are normally held in solution in the blood are deposited in them. The alteration in the tissues consists in some impairment of their nutrition, associated with a diminution in their amount of blood, and a retardation of its circulation. All those conditions which tend to produce atrophic and retrogressive changes in a part, and at the same time to interfere with the circulation in it, are liable to be followed by its calcification. Inflammation, diminished nutritive supply, and general impairment of vitality, may all of them give rise to this process. This is seen in the calcification of atheromatous arteries, of caseous masses in the lungs and lymphatic glands, of many new formations, and of the bloodvessels and cartilages in old people. In its morbid antecedents calcification thus somewhat resembles fatty degeneration, and its pathological signification is in many cases equally difficult to determine. Fatty degeneration indeed very frequently precedes the calcification, which is merely a later stage of the retrograde process.

Respecting the cause of the deposition of the calcareous substances in the atrophied structures—it is probably partly owing, as stated by Rindfleisch, to the stagnation of the nutritive fluids in the part, owing to which the free carbonic acid which appears to hold the salts in solution, escapes, and they are consequently precipitated ; and partly to the non-assimilation of these fluids by the degenerated elements of the tissue.

The calcareous particles make their appearance both within the cells and in the intercellular substance ; they are much more frequent, however, in the latter situation. They are seen at first as fine molecules scattered irregularly through the intercellular substance. (Fig. 22.) They are characterized, when viewed by transmitted light, by their opacity, dark black color, and irregular outline, and also by their solubility in dilute mineral acids. They gradually increase in number until ultimately large tracts of tissue may be converted into an opaque calcareous mass, in which the cells are inclosed and can

no longer be recognized. These larger masses have a sharp black irregular outline, and as the calcification becomes complete, acquire a homogeneous, glistening, semi-transparent appearance. The cells them-

FIG. 22.



A Calcified Sarcomatous Tumor. Showing the minute calcareous particles scattered through the intercellular substance. To the left of the figure they are so abundant as to almost completely obscure the cells. $\times 203$

selves are much less frequently infiltrated, being usually merely enclosed and obscured by the calcified intercellular substance. Calcareous particles may, however, make their appearance in the protoplasm, and gradually increasing convert the cell into a homogeneous calcareous body.

The calcareous matters consist for the most part of lime and magnesian salts, especially the phosphates and carbonates. If the latter are present, the addition of a little dilute hydrochloric acid is followed by the appearance of numerous minute air bubbles in the tissue, owing to the liberation of carbonic acid. In those cases in which calcification is associated with retained gland secretions, the calcareous matters will consist of the specific gland salts.

A part which has become calcified undergoes no further change, its vitality is completely destroyed, and it remains as an inert mass. In this respect calcareous differs from fatty degeneration. In the latter, subsequent changes invariably take place; the part either softening, caseating, or becoming the seat of calcification itself. It differs also in its effect upon the tissue. The structure of the affected part is not destroyed, and there is no annihilation of histological elements, such as occurs in fatty degeneration. The tissue is simply impregnated with calcareous matters, which have no other effect upon it than to render it inert; its vitality is destroyed, but its structure—in so far as the calcification is concerned—remains unaltered. If the saline matters are dissolved out with a little dilute mineral acid, the structure of the part may be again recognized, unless, indeed—as is so often the case—it has been destroyed by any antecedent change.

Calcification must thus be looked upon in many cases as a salutary lesion, the impregnation with calcareous matters preventing subsequent changes in the part. This is especially the case when it is secondary to other forms of degeneration. It is often the most favorable termination of the large class of fatty changes, as is exemplified by the calcification of caseous products in the lungs, and of many new formations. It may, on the other hand, under certain circumstances, be attended with most deleterious consequences, as in the case when it affects the arterial system.

Calcification of Arteries.

Calcification of arteries, like fatty degeneration, may be a *primary* or *secondary* affection. As a secondary change it constitutes one of the terminations of the atheromatous process, and as such is constantly met with in the aorta and its branches, and in many other situations. (See "Atheroma.")

Primary calcification is much less frequent than that which occurs in atheromatous vessels. It is essentially a senile change, and is the result of that general impairment of vitality which exists in advanced life. It is associated with atrophy of the arterial tissues, and in some cases with fatty degeneration. As it is greatly dependent upon general malnutrition, the change is a more or less general one, and when occurring in one part is met with in others. It usually occurs in vessels of medium size, the arteries of the upper and lower extremities and of the brain being those most commonly affected. Its most common seat is the middle coat, where it commences in the muscular-fibre cells. The calcareous particles make their appearance at first around and within the nucleus, and gradually increase until they fill the cell, which becomes converted into a small calcareous flake. The process may go on until the muscular coat is completely calcified, or it may be limited to isolated portions of the coat giving rise to numerous calcareous rings and plates which are irregularly distributed throughout it. From the muscular, it may extend to the external and internal coats until ultimately the vessel becomes calcified throughout.

The vessel thus calcified, loses its elasticity and contractility; its lumen is diminished, and it is transformed into a hard, rigid, brittle tube. This condition is common in the external iliac and in the vessels of the lower extremity, where it is a frequent cause of senile gangrene. (See "Senile Gangrene.")

CHAPTER IX.

PIGMENTARY DEGENERATION.

PIGMENTARY Degeneration, or Pigmentation, consists in an abnormal formation of pigment in the tissues. All true pigments are derived from the coloring matter of the blood. Physiologically, many of them are eliminated by the kidneys and liver; others are deposited in the tissues and there remain permanent. The choroid coat of the eye and the skin of the negro, are well-known examples of tissues in which there is this permanent accumulation of pigment. The cells in these situations appear to be endued with a special power to abstract the coloring matters from the blood, and to store them up in their interior, where they undergo certain chemical changes and become converted into pigment.

In the pathological process also, the pigment is derived from the same source, although its presence in the tissues is rarely dependent upon any abnormal secreting powers in their cellular elements, but is usually the result of certain changes in the circulation or in the blood-vessels, owing to which the coloring matter of the blood escapes and infiltrates the surrounding parts. This escape of hæmoglobin may be owing to rupture of the vessels themselves, or to conditions of congestion or stasis in which the blood-corpuscles and liquor sanguinis pass through their walls. In either case the hæmoglobin will permeate the tissues and ultimately be converted into pigment. Rupture of the vessels and the direct extravasation of blood, is, however, the most common antecedent of the pigmentary change. Soon after the extravasation has taken place, the hæmoglobin escapes from the red blood-corpuscles, either by exudation or by destruction of the corpuscle, and mixed with the liquor sanguinis, infiltrates the surrounding tissues. The red blood-corpuscles themselves may also penetrate the adjacent cells (Langhans). In other cases the process takes place without any solution of continuity in the walls of the vessel. This frequently occurs in conditions of inflammatory stasis and mechanical congestion, in which the red corpuscles pass through the walls of the capillaries, and some of the hæmoglobin is also liberated from the corpuscles

within the vessels, from which it transudes, dissolved in the liquor sanguinis, without rupture having taken place. In whichever of these ways the hæmoglobin is derived, it infiltrates the tissues, staining both the cells and the intercellular substance a yellowish or brownish-red color. It is taken up, however, more readily by the cells than by the intercellular substance or by membranous or fibrous structures.

After the hæmoglobin has remained in the tissue for some length of time, it undergoes certain changes: It becomes darker and more or less granular, minute reddish-brown or black granules and crystals make their appearance both in the cells and in the intercellular substance, and these may gradually increase and form larger masses. This change in the extravasated hæmoglobin is a chemical one, and the substance into which it is converted is *hæmatoidin*. Hæmatoidin appears to be closely allied to the coloring matter of the bile, *cholepyrrhin*, which is also a derivative of hæmoglobin. It exhibits similar reactions when treated with concentrated mineral acids, displaying the same variations of green, blue, rose, and yellow colors. It is insoluble in water, alcohol, ether, and in dilute mineral acids and alkalis; it is soluble in the caustic alkalies giving a red color. It contains more carbon than hæmoglobin; and it also contains iron.


The granules of hæmatoidin vary in size from the smallest particles to masses as large as a red blood-corpuscle. (Fig. 23.) The larger ones are round, or more commonly irregular in shape, and have a sharp defined border. Their color varies from yellow, red, and brown, to black. These variations appear to depend upon the age of the granules and the tissue in which they are formed; the older they are

FIG. 23.



Cells containing pigment. From a melanotic sarcoma of the liver. $\times 350$.

FIG. 24.



Hæmatoidin crystals. (Virchow.)

the blacker they become. The smaller granules are usually dull and opaque; the larger ones, however, often present a more or less glistening appearance. The crystals of hæmatoidin are opaque rhombic prisms, usually of a beautiful yellowish-red or ruby-red color, sometimes approaching to brown or black. They may also occur as little

plates and fine needles, but these are less common forms. (Fig. 24.) They are in most cases so small that considerable care is required to recognize their crystalline nature under the microscope, and they may easily be overlooked as merely irregular granular masses. In some cases, however, they attain a larger size. They are more or less transparent, and present a shining strongly refracting surface.

Whether the hæmoglobin is converted into granular or crystalline hæmatoidin appears partly to depend upon the tissue in which it is situated, the crystals being exceedingly common in some situations, as in the brain and ovaries, whereas in others, as mucous membranes, only the granules are met with. Both the granules and crystals are characterized by their durability and by their great powers of resistance; when once formed they undergo no further change.

Those forms of pigment—both granular and crystalline—which are of an intensely black color, have been supposed to consist of a substance, which differs in chemical composition from hæmatoidin, and which has been called *melanin*. There appears, however, to be no foundation for such a distinction. Melanin is probably merely hæmatoidin, which has become more or less altered by age. It is endued with greater powers of resistance, being less readily soluble in reagents than the more recently formed hæmatoidin, and it contains more carbon.

Pigmentation, although one of the most common and universal forms of degeneration, is of comparatively little importance as a morbid process. The mere existence of pigment within and between the histological elements of the tissues, has in itself but little influence upon their vitality and functions. The atrophy and impairment of function which so frequently accompany it, must rather be looked upon as the result of those conditions upon which the formation of the pigment depends, than as in any way owing to the presence of the pigment itself.

As evidence of other antecedent conditions, pigmentation assumes a more important aspect. The pigment being derived from extravasated hæmoglobin, in whatever situation it occurs, it is usually to be looked upon as the result of some alteration in the circulation or in the bloodvessels, owing to which the escape of the coloring matter is permitted. Exceptions to this exist, however, in the case of certain pigmented new formations, in which the presence of the pigment appears to be mainly owing to the selective power of the cells; these, like those of the choroid, separating the coloring matter from the blood. It is those growths which originate in tissues normally containing pigment,

as the choroid and rete mucosum, which are most frequently melanotic. (See "Melanotic Sarcoma.") In Melanæmia, again, the large quantities of pigment which exist in the blood are the result of a local formation—for the most part by the spleen. Lastly, in Addison's disease, the pathology of the pigmentation of the skin is at present involved in obscurity.

Pigment is often the only evidence of a former extravasation. This is frequently the case in cerebral hemorrhage, where the crystals of hæmatoidin may be all that remains to indicate that rupture of the capillaries has taken place. In the ovaries, also, the slight hemorrhage which follows the escape of the ovum at each menstrual period, is marked by the formation of pigment which constitutes the "corpus luteum." In mechanical congestion and inflammation, again, the consequent pigmentation may be the principal evidence of the former existence of these conditions: this is especially seen in pigmentation of the mucous membrane of the stomach and intestines. The formation of pigment is thus, with the few exceptions above named, the result of some antecedent change in the bloodvessels or circulation; and its presence in the tissues appears to be little more than a testimony to the existence of those processes upon which its formation depends.

FALSE PIGMENTATION.—There are certain forms of discoloration of the tissues which are not due to the presence of hæmatoidin: these must be distinguished from true pigmentation. The most important of them, and that which is most closely allied to the process already described, is the staining of the tissues with the coloring matter of the bile, which is itself a derivative of hæmoglobin, and is, as before stated, very analogous to hæmatoidin. This yellow staining may affect nearly all the tissues, constituting "jaundice;" or it may occur in the liver alone, from local obstructions to small bile-ducts, as is often seen in cirrhosis of that organ. In these cases, however, there is merely the staining of the tissues with the coloring matter of the bile, and no subsequent conversion of this into pigment.

The discoloration caused by the long-continued use of the salts of silver must also be distinguished from true pigmentation: the color here is due to the deposition of the silver in the tissues. The black color of gangrenous parts, and that sometimes produced by the effusion of large quantities of blood into the tissues, must again not be confounded with pigmentation. The discoloration in these cases is the result of the action of the sulphuretted hydrogen upon the color-

ing matter of the blood. The greenish-black discoloration so often seen on the surface of the liver, kidneys, and other abdominal organs after death, is in the same manner due to the intestinal gases. Lastly, the minute particles of inhaled carbon which are always met with in the lungs, must be distinguished from true pigment.

Pigmentation of the Lungs.

In no organs is pigment met with so frequently and in such large quantities as in the lungs, and here much discussion has arisen as to its nature and origin. The lungs normally contain more or less black pigment, the amount of which gradually increases with advancing age—the lungs of infants and young children being almost free from it, whereas those of adults invariably contain it in considerable quantities.


This normal pigmentation of the lungs is principally due to the presence of carbon, and not to that of true hæmatoidin-pigment. The carbon—which is derived from the incomplete combustion of wood, coal, and other substances, and is always present in varying quantities in the atmosphere—is inhaled, and the minute particles pass into the finest bronchial tubes. Having entered the bronchi, many of them are taken up by the mucus-corpuscles, where they may be seen as small black granules within the cells. These may readily be observed in the cells of the grayish-black sputum which is so frequently expectorated in the early morning. Much of the carbon thus inhaled is eliminated by expectoration; many of the particles, however, pass into the air-vesicles, and here their removal by this means being less readily effected, they gradually penetrate the pulmonary substance, and make their way into the alveolar walls and interlobular tissue. It is in these situations that most of the pulmonary pigment is found, and there it may be seen either within the connective tissue-cells, or lying free amongst the fibres. (Fig. 25.)

The means by which the particles of carbon penetrate the walls of the air-vesicles, and make their way into the inter-alveolar tissue, has recently been explained by the researches of Dr. Klein on the histology of the lungs.¹ Dr. Klein finds that the branched connective-tissue cells of the alveolar walls send a process, or a greater or less portion of their body, between the epithelial cells of the alveolus into the

¹ "On the Anatomy of the Lymphatic System of the Lungs," by Dr. E. Klein. "Proceedings Royal Society, No. 149." 1874.

alveolar cavity. As these connective-tissue cells lie in the serous canals which constitute the commencement of the perivascular lymphatics, it is easy to understand how these openings in the alveolar walls (pseudostomata) may become sufficiently distended to allow cells and

FIG. 25.



Pigmentation of the Lungs. From a woman, æt. sixty-five, with slight emphysema. Showing the situation of the pigment in the alveolar walls, and around the bloodvessel v. $\times 75$.

other substances to pass through them from the alveolar cavity into the inter-alveolar tissue.

When once the carbon has made its way into the interlobular tissue, some of it is taken up by the fixed cells in this situation, whilst that which is not thus detained, passes on to the lymphatics, and is deposited in the bronchial lymphatic glands, where the black particles are also visible.

Closely allied to this physiological pigmentation of the lung from the inhalation of carbon, are those morbid conditions which result from the inhalation of particles of coal, stone, iron, and other substances,—of which the lungs of miners, stonemasons, and grinders afford frequent examples. Here also minute particles enter the bronchi, penetrate the walls of the alveoli, and are deposited principally in the interstitial tissue. In the case of miners—in which this is the most common—the particles of coal enter the lungs in such large quantities as to give to them a uniform dark black color. In stonemasons, grinders, etc., the lungs also become deeply pigmented, although to a less extent than those of miners.

The black color of the lungs in these cases, however, is not entirely due to the presence of the inhaled substances, but partly to that of true

hæmatoidin-pigment. The inhalation of the irritating particles sets up inflammatory changes in the bronchi and pulmonary tissue, causing chronic bronchitis, chronic lobular pneumonia, and a large increase in the fibrous tissue of the lungs, which thus ultimately become consolidated, excavated, tough, and fibrous. ("Colliers" and "Knife-grinders' Phthisis," etc.) Owing to these structural changes there is a considerable escape of coloring matter, either from rupture of the capillaries or transudation of serum, and hence a large formation of true pigment; and to this true pigment much of the dark color of these lungs must undoubtedly be ascribed. The lungs of stonemasons and grinders are, like those of miners, deeply pigmented, although to a less degree, but the black color in the former cases cannot be entirely accounted for on the supposition that it is due to the presence of inhaled particles.

Pigmentation of the lungs from the presence of hæmatoidin occurs as the result of many other morbid conditions, many diseases of these organs being attended by the formation of pigment. In chronic phthisis, pigmentation occurs, partly as the result of the inflammatory process, and partly from the obstruction of the vessels caused by the

new growth:—lines of pigment are constantly seen surrounding the nodules of consolidation. In acute croupous pneumonia, the blood which is extravasated into the air-vesicles, and which in the early stages gives to the expectoration a rusty or prune-juice color, subsequently becomes converted into pigment, and the sputum becomes of a grayish-black; the pigment granules being visible in the newly formed cells. The cells met with in the sputum of bronchitis also contain granules of pigment (Fig. 26); and pigmentation plays an important part in the condition of the lungs known as brown induration. (See "Brown Induration of the Lungs.")




FIG. 26.
Cells from the Sputum of Acute Bronchitis. Showing the minute granules of pigment within the cells. Some of the cells also contain a few fatty molecules. $\times 400$.

ment (Fig. 26); and pigmentation plays an important part in the condition of the lungs known as brown induration. (See "Brown Induration of the Lungs.")

Pigment in the lung usually occurs as black irregular granules; it is rarely met with in a crystalline form. In all cases in which it is found in any quantity in the lung it is also found in the bronchial glands. It is taken up by the lymphatics, and, like the inhaled carbon, it becomes arrested in its passage through these glands, where it remains permanently.

CHAPTER X.

TISSUE-CHANGES IN PYREXIA.

It is proposed in the present chapter to allude very briefly to those alterations in the tissues which are met with in certain pyrexial diseases, to which have been applied the terms "parenchymatous" or "granular degeneration," "albuminous infiltration," "acute," or "cloudy swelling."

It is well known that in most diseases which are accompanied by a considerable elevation of the bodily temperature, and especially in those in which the blood has undergone marked changes, the organs and tissues are found much altered after death. The diseases in which such alterations are most frequently met with are pyæmia, erysipelas, typhus, typhoid, and other acute specific fevers, and acute rheumatism. They also occur in other diseases which are attended by considerable pyrexia; and the amount of tissue-alteration usually bears—*cæteris paribus*—a direct relation to the amount of the pyrexia. The organs in which the alterations are more especially marked are the liver, the kidneys, the heart and muscles, and the lungs.

The physical characters of the altered organs vary. It may, however, be stated generally that the organs are more or less swollen and opaque, somewhat diminished in consistence, and abnormally friable. Their vascularity is in some cases diminished, in others slightly increased. When examined microscopically, the cellular elements are found to be increased in size, and their protoplasm markedly granular, so that, in some cases, the nucleus is so much obscured as to be indistinguishable. (Fig. 27.) The granular condition of the protoplasm appears in most cases to be due to albuminous particles, inasmuch as it disappears upon the addition of dilute acetic acid. In other cases, however, in which the change is apparently more advanced, the granules are larger, insoluble in acetic acid, and obviously fatty.

The Liver.—Here the change is usually met with in its most marked degree. The organ is slightly enlarged, abnormally soft and friable, and the cut surface has a dull opaque look, being paler than natural.

FIG. 27.



Liver from a case of Acute Rheumatism with high Temperature. Showing the swollen and granular condition of the liver-cells. In many of the cells the nucleus is so much obscured as to be almost indistinguishable. $\times 200$.

The liver cells are swollen and granular, and in many cases contain fatty particles. (See Fig. 27.)

The Kidneys.—In the kidneys the change affects especially the cortex. This is swollen, opaque, and friable. The Malpighian bodies and the pyramids are usually abnormally vascular, and thus contrast with the pale cortex. The epithelium in the tubes of the cortex presents the appearances above described. These are precisely similar to those met with in the earlier stages of tubal nephritis. (See Fig. 104.)

The Heart.—The alteration produced in the heart consists in slight opacity, pallor, and diminution in the consistence of the muscular tissue.

FIG. 28.



Muscular Tissue of the Heart, from a case of severe Typhoid Fever. Showing the granular condition of the fibres and the loss of their striation. $\times 400$.

Under the microscope the muscular fibres are seen to have lost their distinct striation and to be finely granular. (Fig. 28.) Such a condition must materially interfere with the contractile power of the organ. A similar change is met with less frequently in other muscles.

The Lungs.—The change in the lungs has been described by Buhl as consisting in swelling of the alveolar epithelium. The epithelial elements are markedly granular from the presence of albuminous and fatty particles, and they become loosened from the alveolar walls. The change affects, more or less, the whole of both lungs. The organs are enlarged, œdematous, and abnormally friable.¹

This change occurs not only in pyrexia, but also in the earlier stages of the process of inflammation. Here also a swollen and granular condition of the protoplasm is met with, especially in epithelial and endothelial elements. It is to this that Virchow gave the name of “cloudy swelling.” (See chapter on “Inflammation.”)

¹ Buhl, “Lungenentzündung, Tuberkulose, und Schwindsucht.”

Respecting the nature of the change—nothing is certainly known. Dr. Wickham Legg produced it artificially in animals by submitting them to a high temperature, and he, in common with some other pathologists, is inclined to look upon it simply as a result of the high temperature. An exact knowledge of its pathology, however, must probably await further accurate experimental investigation. In the mean time, when it is borne in mind that the conditions in which it occurs—fever and inflammation, are both attended by marked alterations in the nutritive processes, and that in advanced degrees of the change it is accompanied by more or less fatty metamorphosis, it may be regarded as probable that its occurrence will be found to be due partly to interference with the normal processes of tissue-oxidation, and partly to increased transformation of the protoplasm of the cells. (See “General Pathology of Fatty Degeneration.”)

Whatever be the nature of the change, there can be no doubt that it must very materially interfere with function, and that its supervention in the course of acute disease—especially when affecting the heart—must constitute a most important source of danger.

CHAPTER XI.

NUTRITION INCREASED.

THE morbid changes thus far described, have been attended either by *arrest* or by *impairment* of nutrition;—those remain to be considered in which the nutritive activity is *increased*. They include Hypertrophy and the New Formations.

Hypertrophy.

Hypertrophy is an increase in the amount of a tissue, owing to an increase in the size, or to an increase in the number, of its histological elements. This increase, however, in order to constitute hypertrophy must affect those elements more especially upon which the peculiar functions of the part depend; in muscle, for example, the muscular fibres; in the kidney—the urine-tubes and bloodvessels.

When this increase in the amount of the tissue is owing simply to an increase in the *size* of the elements of which it is composed, it is termed *simple* hypertrophy; when to an increase in their *number*, and to the formation of a new tissue, *numerical* hypertrophy, or *hyperplasia*. The two forms of hypertrophy are thus comparable with the two forms of atrophy:—in simple hypertrophy as in simple atrophy, there is merely an alteration in the *size*; in numerical—an alteration in the *number* of the elements.

Hypertrophy is in most cases the result of an increase in the functional activity of the tissue, this increased activity being induced by a necessity for some increased manifestation of function; in muscle, for example, in order to overcome some obstruction, in a secreting organ—to secrete more fluid. Such hypertrophies are consequently *conservative* in their nature, and they are usually spoken of as *functional* hypertrophies.

It is in muscular tissue that functional hypertrophy is most frequently met with. Examples of it are furnished by the hypertrophy of the muscles of the calf in ballet dancers; also by the hypertrophy of a hollow viscus from obstruction to the exit of its contents—as of the heart from obstruction at the valvular orifices or in the course of the circulation, of the muscular coat of the stomach in stricture of the pylorus, of the intestine above a permanent stricture, and of the bladder in stricture of the urethra. The kidney also may become hypertrophied, owing to the loss or incapacity of its fellow. In bone, hypertrophy of the fibula has been observed in cases of disease, or ununited fracture of the tibia. In all these cases the necessity for increased activity leads to increased development.

In exceptional cases hypertrophy is not dependent upon an increase in the functional activity of the tissue. It may sometimes result from an increased supply of blood to a part, and there are some forms of hypertrophy which are congenital.

The term “hypertrophy” is also frequently applied to those enlargements of organs which result from long-continued irritation; as to enlargements of the lymphatic glands, of the tonsils, and of the prostate gland. Such growths, however, are some of them inflammatory in their nature, others come more properly within the category of “tumors.”

CHAPTER XII.

THE NEW FORMATIONS.

INCREASED nutritive activity of a tissue—as has been seen in the preceding chapter—leads not only to the enlargement of its component elements, but also to the production of new ones. It is this production of *new elements*, and the various structures to which they give rise, that constitutes the New Formations. The New Formations comprise both *inflammatory* and *non-inflammatory* growths.

The *inflammatory* new formations differ from the non-inflammatory in being caused by some *injury*, upon the removal of which the process of growth usually ceases. They are characterized by their *instability*: they either rapidly undergo retroversive changes, or, at the most, form a tissue which in its structure and life is similar to that from which they originated. Their tendency is gradually to approximate to a healthy condition, and, not like the non-inflammatory growths, continuously to increase. These will be treated of in the chapters on “Inflammation.”

The *non-inflammatory* new formations are divisible into—the Regenerations, the Hyperplasias, and the Tumors.

The regeneration of lost or destroyed tissue belongs rather to the domain of normal than of pathological histology, and needs here no further consideration.

The Hyperplasias are new formations of tissue which in structure and formation are similar to that from which they originated. Such new formation has been seen in the preceding chapter to occur in many cases of hypertrophy (numerical hypertrophy). Most frequently, however, a hyperplastic process gives rise to growths which come within the category of tumors. This is seen, for example, in a localized hyperplasia of the papillary layer of the skin leading to the formation of a papilloma.

The tumors, much the most important class of new formations, must be treated of more fully.

The Tumors.

The tumors are new formations which in their development and growth are characterized by their *independence* of the rest of the body; they increase in size by virtue of their own inherent activity, which differs from, and is, to a great extent, independent of that of the surrounding tissues. Having attained a certain size, they either remain permanent, or, more frequently, *tend continuously to increase*.

Whatever be the nature of the tumor it is always the direct product of the elements of a pre-existing tissue, these elements being in some cases migrated white blood-corpuscles. In order therefore to understand the pathology of the tumors, it is necessary to be intimately acquainted with the histology and mode of development of the normal tissues.

ETIOLOGY OF TUMORS.—As tumors are the result of the increased nutritive activity of the elements from which they originate, it will be readily understood that their causes must for the most part be obscure, and that in many cases all that can be said is, that the new growth is the result of the *spontaneous* activity of the elements from which it springs. Sometimes, however, the causes are either wholly or partially ascertainable. They may be divided into *constitutional predisposing*, and *direct exciting*, causes.

Constitutional Predisposing Causes.—That many tumors owe their origin to some constitutional taint, has long been a universally accepted pathological doctrine. The constitutional cause has frequently been regarded as a general one, as consisting either in some alteration in the constitution of the blood, or in some abnormal condition of the physiological processes throughout the entire organism. It was formerly supposed that many new formations were the result of an exudation from the bloodvessels, and that the elements of the growth were produced spontaneously in the exuded structureless blastema. Such growths were looked upon as the local expression of a vitiated constitution of the blood—a *dyscrasia*. Although this hypothesis is now universally abandoned, and all new formations are known to originate from pre-existing cellular elements, an alteration in the constitution of the blood or in the performance of the physiological processes throughout the body, is still regarded by many as playing an important part in their causation.

It is the malignant tumors which are thus supposed to owe their origin to the existence of a *general* constitutional taint. These growths

are said to be of a *constitutional* origin, in contradistinction to the non-malignant growths, which are looked upon as purely *local*. This hypothesis is principally based upon the clinical characteristics of the malignant growths—their tendency to recur after removal, their multiplicity, and the difficulty or impossibility of completely eradicating them.

It is said that the development of malignant growths is frequently preceded by an unhealthy state of the constitution—a cachexia. In the majority of cases of malignant formation, however, there is no evidence of any such cachexia preceding the local growth. The individual is usually in good health at the time of the occurrence of the primary tumor. The general impairment of nutrition and emaciation, which constitute the cachexia, are *secondary* to the local growths, and are in direct proportion to their extent and situation: the more extended the local lesions, the greater the amount of discharge, the more the lymphatics and the digestive organs are involved, the more marked is the attendant cachexia.

The multiplicity of malignant growths is also adduced as an argument in favor of the existence of a general constitutional taint. The fact that malignant growths are frequently multiple, constitutes in itself no ground for such a conclusion. The multiplicity is for the most part a *secondary* phenomenon, the secondary tumors resulting from infection by the primary one. Simple *primary* multiplicity is not so characteristic of malignant as of many other tumors—the lipomata, fibromata, sebaceous tumors, and warts, are all more often *primarily* multiple than cancer. Multiplicity in many cases is evidence rather of a *local* than of a general taint. In scrofula, for example, the lymphatic glands generally are prone to become the seats of new growth, and tumors are often multiple in bone without occurring in other tissues. In such cases there would appear to be a *local* rather than a *general* cause.

The recurrence of the malignant growth after removal at the seat of the operation, may again be owing to the removal having been incomplete, some of the proliferating elements of the growth having been left behind. The cellular elements of malignant tumors often extend into the adjacent tissues for some distance beyond the apparent confines of the tumor, and as the physical characters of the infiltrated tissues differ in no way from those of the healthy, there are no means of certainly determining how far wide of the tumor the incision must be carried in order to include the whole of the affected structures.

(See "Malignancy.") A tendency to local recurrence is a property possessed by many growths, and it can be explained on local grounds without the necessity of admitting the existence of a general taint.

Lastly, the fact that malignant growths are usually followed by the development of similar growths in the lymphatic glands and in internal organs, may in most cases be more readily explained—as will be seen when speaking of "malignancy"—by regarding these as the result of infection by the primary tumor, than by ascribing them to the existence of a common constitutional cause.

The reasons already adduced in a preceding chapter (see "Introduction"), for considering all changes in the constitution of the blood as secondary to local causes, would appear of themselves to be sufficient to render untenable the hypothesis of a *primary* blood dyscrasia. Any abnormal condition of the blood which may be associated with the development of malignant tumors must probably be regarded as resulting either from the absorption of deleterious substances, from the entrance into it of the elements of the growth, from the drain of an attendant discharge, or from interference with the processes of indigestion, assimilation, or secretion, or with the formation of the blood itself. At the same time it must be borne in mind, when discussing the possibility of the development of malignant growths being the result of a "blood disease," that the emigration of white blood-corpuscles may play some part in the process. How far this is the case, as will be seen hereafter, we are at present unable to determine. We know that this emigration constitutes a prominent feature in the process of inflammation, and if future researches should show that it also occurs in the development of malignant new formations whilst in those which are non-malignant it is entirely wanting, it would go far towards rendering the existence of a malignant dyscrasia beyond dispute. In the present position of our knowledge, however, in which it is *only probable* that an emigration of leucocytes may play some part in the development of all new growths, both innocent and malignant in favor of the existence of a blood-dyscrasia as the cause of the development of the latter based upon such considerations would be unjustifiable. If an emigration of leucocytes does take place, it is more in accordance with facts and with the general history of malignant growths to suppose the existence of some peculiarity in the tissues into which the elements escape which determines their special development.

Although there would thus appear to be no necessity to admit the existence of a general constitutional cause in order to explain the

clinical characters which constitute malignancy, there can be no doubt that the development of many growths, both innocent and malignant, is materially influenced by constitutional conditions. Certain conditions of the constitution may *favor* the development of tumors, just in the same way that they favor, for example, the development of syphilitic lesions. In both cases the *local* processes may be modified by the *general* constitutional state. In the case of tumors it must be admitted as possible that the constitutional influence may be sufficient in itself to determine the development of the new growth; and this possibility must be borne in mind in accounting for the development of secondary malignant growths, as although these may result from infection, they may perhaps in some cases owe their origin to the same causes as those which induced the primary one.

The influence of *hereditary predisposition* must also be taken into account in studying the etiology of tumors. This influence is marked in the case of many growths, both malignant and benign. The inherited peculiarity is probably, in most cases, a *local* rather than a *general* one, consisting in some constitutional peculiarity of the tissues from which the new growths originate, some peculiarity which renders them more prone than other tissues to undergo abnormal development. In scrofula, for example, which is a markedly hereditary disease, the tendency of the lymphatic glands to undergo excessive development from very slight degrees of irritation, is probably to be regarded as owing to a predisposition of the glands themselves, and not to any *general* constitutional state. The same is probably true of many other constitutional tendencies. Nævi, tumors of the skin, uterus, mammae, stomach, and of other parts, are again all unquestionably sometimes hereditary; and here also the tendency would appear to consist in a predisposition of the tissues themselves to become the seats of new formations.

The tendency sometimes observed in *particular* tissues to generate new formations, points again to a local cause. The osseous system, for example, may be the seat of new growths—tumors occurring in nearly all the bones, and not being met with in other parts. Tumors may in the same way be multiple in other tissues.

Direct Exciting Causes.—The existence of a direct exciting cause is more capable of demonstration than a constitutional one. It consists either in some direct irritation of a tissue, which is by this means stimulated to increased development; or in the migration or transmission of elements from some primary growth, which by proliferating

themselves, or causing proliferation, in the tissues in which they lodge, constitute the centres of secondary formations.

The direct irritation of a tissue may be owing to mechanical or chemical irritants. Simple mechanical or chemical irritation can, however, under no circumstances be the *only* cause of the development of the growth. The effect of such irritations alone, is to cause an inflammatory formation; in order for them to produce a non-inflammatory one—a tumor, there must be some special predisposition of the tissue itself:—the irritation can merely *determine* its development. The influence of simple irritation in the production of new formations is exemplified by the frequent occurrence of epithelioma on the lips of smokers, from the irritation of the pipe; and on the penis and scrotum of chimney-sweepers, from the irritation of the soot (“chimney-sweepers’ cancer”). The numerous recorded instances of the development of a tumor following some external violence or injury, leave little doubt that these also sometimes stand to one another in the relation of cause and effect. Lastly, as pointed out by Virchow, it is those organs which are the most exposed to irritation from external causes, as the stomach, the uterus, the mammary gland, and the rectum, which are especially liable to become the seats of new growths.

The influence of elements, which have either migrated or been conveyed from their original habitat, in causing the development of new formations is exemplified by the history of malignant tumors. It is by this means that the secondary tumors most frequently originate. The elements, which either migrate spontaneously from the primary growth, or are carried by the blood or lymph streams, may proliferate in the tissue in which they lodge, and so develop into secondary formations.

In other cases it is also exceedingly probable, as pointed out by Dr. Creighton,¹ that the elements (either cells or minute particles) which are transmitted from the primary tumor cause the production of secondary tumors by virtue of an influence on the cells of the tissue where they lodge, which may be termed a *spermatic influence*, and which is strictly comparable with that of the sperm cell on the ovum. In such cases the relation of the primary to the secondary tumor is that of parent to offspring. This subject will be again alluded to when speaking of “*Malignancy*.”

DEVELOPMENT.—In studying the development of the tumors, it is


¹ Discussion on Cancer, “Trans. Path. Soc. Lon.,” 1874, p. 368.

important, in the first place, to bear in mind what has been already stated—viz., that they are in all cases the direct product of pre-existing cellular elements. In their development they also resemble the normal tissues:—every pathological growth has its physiological prototype.

The elements from which tumors most frequently originate are those belonging to the *common connective tissue*, and to the bloodvessels and lymphatic system with which it is so intimately associated. By common connective tissue is meant that tissue which in all parts surrounds the bloodvessels, and is so universally distributed throughout the entire organism. This must be carefully distinguished from the formed connective substances—tendon, cartilage, bone, etc. In this common connective tissue we distinguish two kinds of cells—the stable cells (connective tissue corpuscles), and the mobile cells, which are probably wandering white blood-corpuscles. These cells are in intimate relation with the endothelium of the lymphatics, the latter vessels commencing as serous canals which are universally distributed in the tissue. Further—both the endothelium of the lymphatics and that of the bloodvessels closely resemble in their physiological functions the fixed cells of the connective tissue.

In the process of development of tumors from this common connective tissue, the part which is played by the different cellular elements cannot in our present state of knowledge be certainly stated. It is, however, probable that it is the mobile cells which are principally concerned. These cells are the most active, and consequently the most capable of multiplying by division. It is equally probable that the number of these mobile cells is materially increased by an abundant emigration of white blood-corpuscles from the vessels. Respecting the fixed connective tissue cells and the endothelium of the bloodvessels and lymphatics—it must be admitted that these, although perhaps inferior in their formative powers, also undergo active changes, and thus increase the number of the newly formed elements. In the case of these fixed cells, the process of cell-formation is probably very frequently effected by means of vacuolation and endogenous development (see Fig. 3); and, as was stated when describing the “genesis of cells,” elements which thus originate endogeneously have, for the most part, a destiny different from that of the parent cell.

FIG. 29.



Embryonic ("Indifferent")
Tissue from the Edge of a
Sarcomatous Tumor. $\times 350$.

Whatever be the part played by the respective elements in the process of development of the new growth, the first result of their combined activity is to produce a new tissue, composed of innumerable small roundish-shaped cells, from 18^1_{00} to 28^1_{00} of an inch in diameter, often inclosing a large, round, ill-defined nucleus, which usually only becomes visible after the addition of water or of acetic acid. The cells, which are usually known as *indifferent* cells, possess no limiting membrane, but consist of little masses of protoplasm which are almost in close contact with one another (Fig. 29). They are indistinguishable from the cells of a granulation, and from many of those met with in the rapidly growing connective tissue of the embryo. In some cases the protoplasm continues to increase and the nuclei to divide without any subsequent division of the cell taking place, and thus are produced large irregular-shaped masses of protoplasm containing numerous nuclei. These are the giant or myeloid cells which are met with in the medulla of young bone and in many sarcomatous tumors (see "Myeloid Sarcoma," Fig. 50). Thus the first stage in the process of development consists in the formation of an embryonic tissue, and this embryonic tissue subsequently develops into the tissue of which the new growth is composed. It is often impossible to determine in this early stage of the growth what it will ultimately become—whether a fibroma, a sarcoma, or an enchondroma, etc.

The second stage of the process consists in the development of this embryonic ("indifferent") tissue into the tissue of the permanent growth, and this subsequent development closely resembles that of the immature connective tissue of the embryo. As from the immature connective tissue of the embryo are developed various connective tissue substances—fibrous tissue, mucous tissue, cartilage, bone, etc.—so may this embryonic connective tissue, which constitutes the earliest stage of so many of the pathological new formations, become developed into various tissues, all of which usually more or less resemble the several varieties of the physiological connective tissue. The whole of the primary cells may form the same kind of tissue, in which case the growth will possess the same characters throughout; or it may be complex, some cells forming one kind of tissue and some another. A combination of two or more kinds of structure may thus be met with in the same tumor—as a combination of sarcoma and lipoma, of enchondroma and myxoma, and so on. What determines the ultimate development of the young cells, why they produce such

various forms of growths, is as far from our knowledge as what determines the ultimate destination of the cells in the embryo.

Next to common connective tissue, the *epithelia*, surface and glandular, are the elements from which tumors most frequently originate; and as from connective tissue are usually produced growths of the connective tissue type, so do the growths originating from the epithelia usually resemble epithelium. As to whether all new growths originating in connection with epithelium are solely the offspring of pre-existing epithelial elements, we are unable to speak with absolute certainty. The answer to this question must await a more certain knowledge of the way in which the epithelial structures normally grow and repair themselves. It is regarded as probable by some that the normal production of epithelium is not entirely the result of the multiplication of epithelial cells, but that the mobile cells of the connective tissue by *contact* with epithelial elements may become developed into epithelium. If this be so, it must be admitted as at all events equally probable, that pathological new formations which originate in connection with epithelium may be in part the product of mobile elements belonging to the connective tissue. The process of development from epithelium may take place either by simple division, or by vacuolation and endogenous growth.

Here allusion must be made to the recent investigations of Dr. Creighton on the development of secondary tumors in the liver.¹ Dr. Creighton finds that in various kinds of secondary tumor occurring in the liver, the growth originates from the liver-cells by a process of vacuolation and endogenous formation; and he concludes that in this way liver-cells may originate epithelial cells, connective tissue-cells, spindle-cells, etc., the variations depending upon the nature of the primary tumor.

From the remaining tissues, *muscle* and *nerve*, the development of tumors is comparatively rare, and in nerve-tissue it is even doubtful if formative processes ever occur.

According to the similarity or difference which subsists between the new growth and the tissue from which it grows, tumors are divisible into two classes—*homologous* and *heterologous*. When the tumor resembles in its structure and development the tissue from which it originates, it is said to be *homologous*; when it differs, it is said to be

¹ Discussion on Cancer, "Trans. Path. Soc. Lond.," 1874, p. 368.

heterologous. A cartilaginous tumor, for example, growing from cartilage, is homologous, but growing from any other tissue, as from the parotid gland, it is heterologous. The same variety of tumor may thus be in one case homologous, in another, heterologous. A purely homologous growth is therefore simply a *hyperplasia*—an excessive growth of a tissue in some particular part; any deviation from the type of the parent tissue constitutes heterology.

Heterology, however, is not limited to the production of a tissue which is dissimilar from that from which it originates; a tumor is also said to be heterologous when it differs from the tissue in which it is *situated*, and this may occur without its being the direct product of the latter. It is heterology in this sense that is so characteristic of certain cancers. Those cancers which obviously originate from epithelium, become heterologous, owing to the growth and extension of the epithelium beyond its normal limits (see “Epithelioma,” Fig. 60). The same form of heterology obtains in the case of growths originating from elements which have migrated or been carried from their original habitat, and have developed into a tissue which differs from that in which they are situated.

Heterology is often an evidence of malignancy. Many heterologous tumors, however, are not malignant; slight deviations from the normal type are no evidence of malignancy. A growth primarily homologous may subsequently become heterologous; this is often seen when a long-standing innocent tumor suddenly exhibits malignant characters. A knowledge of the homology or heterology of a growth is therefore an important element in the determination of its innocent or malignant nature.

RELATIONS OF THE TUMOR TO THE SURROUNDING TISSUES.—The relation of the tumor to the surrounding structures will depend upon its mode of growth. Growth may take place simply by the continuous proliferation of the cells of which the tumor is composed. In this case it will merely displace the surrounding parts, and having attained a certain degree of development, a fibrous capsule is often formed around it, by which it becomes completely isolated. The lipomata, fibromata, and enchondromata are usually thus encapsuled. In other cases, growth takes place also at the circumference, by the continuous proliferation of the parent tissue, and the invasion of adjacent structures. There is then no line of demarcation between the tumor and the surrounding part, so that although to the naked eye it may

appear separate, the microscope will discover in the adjacent tissues elements of the new growth. (Fig. 30.) This is very common in the sarcomata and in the cancers, and is a common cause of local recurrence after removal. (See "Malignancy.")

RETROGRESSIVE CHANGES.—The development of the tumor being complete, it becomes sooner or later the seat of retrogressive changes. The time at which these commence varies:—As a rule the permanence and durability of a tumor bear an inverse relation to the rapidity of its growth, and to


the inferiority of its organization. The more rapid the growth, and the more lowly organized the tissue formed, the less its durability, and the sooner do retrogressive changes occur. The cancers and sarcomata, for example, which develop rapidly, and consist for the most part of cells, quickly degenerate; their elements are unstable and soon perish. Osseous tumors, on the other hand, which develop more slowly, and consist of a more highly organized tissue, have a much greater stability, and are but little liable to retrogressive metamorphosis.

The retrogressive changes are similar to those met with in the physiological tissues. Deficient supply of blood is followed by fatty degeneration and in various terminations—softening, caseation, and calcification. Pigmentary, colloid, and mucoid degeneration may also occur. Tumors may also become the seats of an inflammatory process.

MALIGNANCY.—By "malignancy" is understood the property possessed by many tumors of reproducing themselves either locally after removal, or in distant tissues. It is important not to confound the terms "malignancy" and "cancerous." "*Malignancy*" is a purely clinical term, and although in a high degree the property of the cancers, is by no means confined to them; the sarcomata, for example, being in many cases equally, or even more malignant. The term "*cancer*," on the other hand, is used to imply a definite structure, and as such is applied to a certain class of new formations.

The malignant properties of a tumor may manifest themselves either in the tissues immediately adjacent to it or in its neighborhood, in the

FIG. 30.



Scirrhus of the Mamma.—A thin section from the most external portion of the tumor. Showing the invasion by small-celled infiltration of the muscular fibres and adipose tissue in the neighborhood of the gland. $\times 230$.

nearest chain of lymphatic glands, or in more distant parts. Some growths possess these properties in a higher degree than others, so that there are different degrees of malignancy. In many cases the malignancy of a tumor is so far limited that it gives rise merely to a local reproduction, after removal. Such tumors have been separately classified by many surgeons as "recurrent" tumors. (See "Small Spindle celled Sarcoma.") In other growths this tendency to local reproduction is associated with the development of similar growths in the nearest lymphatic glands; whilst in a third class of cases, to one or both of these conditions is added the reproduction of the growth in more distant tissues, especially in the lungs and liver. It will be advisable to treat of these three degrees of malignancy separately.

1. *Reproduction of the Tumor in the Adjacent Structures.*—This is usually the earliest evidence of malignancy in a growth, and gives rise to its persistent recurrence *in loco* after removal by the surgeon. This tendency to the local reproduction of the tumor is principally owing to its mode of growth. The cellular elements of the tumor extend for some distance into the surrounding tissues, so that unless in the operation for its removal the incision be carried for some distance beyond the *apparent* confines of the growth, some of the elements may be left behind and thus constitute the centres of secondary formations. (See Fig. 30.) Such tumors therefore are not usually encapsuled, but extend by peripheral growth. In some cases, however, tumors which possess a distinct capsule may in the same way infiltrate the surrounding structures. The infiltration of the adjacent tissues by the elements of the tumors sometimes gives rise, not only to the continuous enlargement of the primary growth, but also to the development of separate secondary growths in the immediate vicinity of the primary one. Whilst it must be admitted that the reproduction of a malignant tumor in the adjacent tissues is thus in most cases to be ascribed to the infiltration of the latter with the cellular elements, it is possible that the nutrient fluids of the tumor may by their influence upon the surrounding tissues be in part the cause of the abnormal development.

2. *Reproduction of the Tumor in the nearest Lymphatic Glands.*—This is owing to the transmission by the lymph-stream of substances (probably cellular elements) derived from the malignant growth, which become arrested in the nearest lymphatic glands, and there cause the development of secondary formations. These are in all cases of the same nature as the primary tumor. When the lymphatic

glands have developed themselves into secondary growths, they in their turn constitute new centres of infection, and may thus infect more distant glands or the immediately adjacent tissues. The tendency to reproduction in the lymphatic glands varies very much in the different varieties of malignant growths, being, for example, very marked in the cancers, whereas in the sarcomata it is comparatively rare. The reasons for these differences will be seen in the subsequent chapters.

3. *Reproduction of the Tumor in the Distant Tissues.*—This is usually the terminal process in the history of malignant growths. The reproduction of the malignant growth in distant tissues is in the great majority of cases owing to the entry of some of its elements into the blood-stream. The secondary tumors, as in the lymphatic glands, are in all cases of the same nature as the primary one, although they are often softer, more vascular, and show greater activity of growth. They occur as a rule in those organs through which the blood from the primary tumor first passes—that is, in those organs which present the first set of capillaries for the arrest of the transmitted materials. In malignant diseases of those organs, for example, which return their blood through the portal vein, as the stomach and mesenteric glands, it is the liver in which the secondary growths usually first occur, and when this has become involved, it may constitute a secondary centre of infection, and in the same way cause tertiary growths in the lungs. Although this sequence is the rule, there are numerous exceptions. In some cases, the organs which are nearest in the course of the circulation to the primary growth escape, whilst those more distant become affected. This may be owing to one organ being more predisposed to the influence of the infecting materials than another; or to the capillaries of the proximal organ allowing particles to pass through them, whereas those of the more distant one are small enough to arrest them. Lastly, it must be borne in mind that the secondary growths may be entirely independent of the primary one, their origin being due to the same cause.

Although the general dissemination of a malignant growth is thus in most cases owing to the transmission of its elements by the blood-stream, this is not the only way in which it may be brought about. Exceptional cases have been described in which the elements of a tumor have been distributed and caused secondary growths in other ways—as by passing down the trachea, travelling between the layers of the peritoneum, and from the kidneys down the ureters to the bladder, etc.

The secondary tumors, as already stated, are probably either the direct products of the transmitted elements which proliferate in the tissues in which they lodge, or they owe their origin to the spermatie influence of those elements upon cells with which they come into contact.

It is thus obvious that the evidences of malignancy in a tumor will consist—in its invasion of the surrounding structures, the implication of the neighboring lymphatics, and the occurrence of similar tumors in internal organs. As a general rule it may be stated that the more juice a growth contains, and the richer it is in bloodvessels and lymphatics, the more quickly will it infect the lymphatic glands and internal organs; on the other hand, the poorer it is in bloodvessels and lymphatics, the more are its infecting properties confined to the neighboring tissues.

The determination of the innocent or malignant nature of any growth will principally depend upon its microscopical characters, and, as will be seen when considering the various kinds of tumors, the differences which exist in their clinical characters are in great measure to be explained by the differences in their position, minute structure, and mode of growth. Many varieties of tumors are invariably malignant, as the cancers and sarcomata; hence any growth which form its minute structure must be included under these heads, must be regarded as being of a malignant nature.

CLASSIFICATION.—Tumors may be classified upon a histological, or upon a physiological and clinical basis. Although a physiological and clinical classification are much to be desired, in the present state of our knowledge they must be very incomplete. The usual classification of tumors according to their histological characters is consequently here adopted. Such a classification may be most advantageously made in accordance with the classification of the physiological tissues:—

Classification of Tumors.

I. *Type of the fully developed Connective Tissues.*

Type of fibrous tissue . . .	Fibroma.
“ adipose tissue . . .	Lipoma.
“ cartilage . . .	Enchondroma.
“ bone . . .	Osteoma.
“ mucous tissue . . .	Myxoma.
“ lymphatic tissue . . .	The Lymphomata { Hodgkin's Disease. Leukæmia.

II. *Type of Higher Tissues.*

Type of muscle	Myoma.
“ nerve	Neuroma.
“ bloodvessels	Angioma.
“ papillæ of skin or mucous membrane	} Papilloma.
“ secreting glands	
	Adenoma.

III. *Type of Embryonic Tissue.**The Sarcomata.*

Spindle-celled sarcoma	.	.	{ <div> Small spindle-celled sarcoma. <div> Large " " " </div> Melanotic sarcoma. Osteoid sarcoma. </div>
Round-celled sarcoma	.	.	
Myeloid sarcoma.			
			{ <div> Glioma. Alveolar sarcoma. </div>

IV. *The Carcinomata.*

Scirrhus.
 Encephaloid.
 Colloid.
 Epithelioma—Adenoid Cancer.

V. *Cystic Tumors.*

Cysts.

CHAPTER XIII.

THE FIBROMATA.

THE fibromata, fibrous, fibro-cellular, or connective tissue tumors, are tumors consisting of fibrous tissue.

STRUCTURE.—In structure the fibromata present the same variations as those met with in fibrous tissue. Some of them are composed of firm, dense, fibrous tissue, such as constitutes tendons; others are laxer and less fibrous in consistence, more resembling the connective tissue of the cutis. The fibres, which constitute the chief part of the growth, are more or less closely interlaced, and are distributed without any definite arrangement, or grouped in bundles of various sizes. They

are sometimes arranged concentrically around the bloodvessels. (Fig. 31.) Yellow elastic fibres are but very rarely met with. The cells, like those of normal fibrous tissue, are very few in number, and are usually most abundant around the vessels. They are minute, spindle-shaped, fusiform, or stellate bodies, the latter having processes of varying length, which communicate with similar processes from neigh-

FIG. 31.



Section of a Fibrous Tumor of the Skin.—In the neighborhood of the cut bloodvessel *v* are seen some cells; also fibres cut transversely. $\times 200$ and reduced $\frac{1}{2}$.

boring cells. They are often so small and indistinct as in the fresh specimen only to become visible after the addition of dilute acetic acid. The size and number of these cells vary with the rapidity of growth—the slower the growth the more fibrous the tissue, and the smaller and less numerous are the cells.

The fibromata usually contain but few bloodvessels. In the softer growths, however, these are often more numerous. They sometimes form a cavernous network, the walls of which are firmly united to the tissue of the tumor, so that when divided or ruptured they are unable to retract. In such cases, injury to the tumor is often followed by profuse hemorrhage.

DEVELOPMENT.—The fibromata originate from connective tissue, either from the cutis or subcutaneous connective tissue, from the submucous or subserous tissue, from fasciæ, the periosteum, the neurilemma, or from the connective tissue of organs. In the earliest stages of their growth the cells are more numerous than when development is complete. (See “Development of New Formations.”)

SECONDARY CHANGES.—Of these, partial mucoid softening and calcification are the most common. Ulceration also sometimes occurs in these growths which are situated in the skin and submucous tissue.

VARIETIES.—Fibrous tumors present some variations in their cha-

racters, which depend for the most part upon the tissues from which they grow. Two classes may be distinguished.

1. *Soft Fibromata*.—These consist of the looser and less dense form of fibrous tissue. They are met with as diffused growths in the subcutaneous and submucous tissues. In the former situation they often form large pedunculated and non-encapsuled tumors, which are commonly known as *wens*. These are sometimes multiple. A similar increased growth of the subcutaneous tissue is also met with in *Molluscum Fibrosum*. In this disease the large masses which hang down from the thighs, buttocks, and other situations consist simply of loose fibrous tissue. The new growth here often contains numerous large bloodvessels, so that its removal may lead to dangerous hemorrhage.

In addition to these diffused growths, more circumscribed and *encapsuled* fibrous tumors of the soft variety are occasionally met with growing from the scalp, scrotum, labium, intermuscular septa, and other situations.

2. *Firm Fibromata*.—These are composed of dense fibrous tissue resembling that met with in tendons. They are firm, hard, encapsuled tumors, presenting on section a grayish-white, glistening, fibrous appearance. These tumors often occur in connection with bone, especially with the upper and lower jaw, originating either in the centre of the bone or from the periosteum. Growing from the periosteum of the alveolus they constitute simple fibrous *epulis*. They are also met with in the nose, where they form one variety of *nasal polypus*. It is in these firm fibrous growths that the communication of the bloodvessels with cavernous spaces already alluded to, is sometimes found.

Another variety of firm fibrous tumor grows in connection with nerves, and is often described as *neuroma*. True neuromata, however, *i. e.*, new formations of nerve-tissue—are amongst the rarest forms of new formations. These fibrous growths most frequently occur in connection with the superficial nerves. They grow from the neurilemma, and as they increase in size the nerve-fibres become expanded over them. They are very firm, rounded tumors, and are frequently multiple.

The fibroid tumors of the uterus, which are often described as fibrous tumors, appear in most cases to be overgrowths of the involuntary muscular tissue of the organ. They will therefore be considered with the muscular tumors. (See “Myoma.”)

CLINICAL CHARACTERS.—Clinically the fibromata are perfectly innocent. They grow slowly, and have no tendency to recur locally after removal.

CHAPTER XIV.

THE LIPOMATA.

A GENERAL new formation of adipose tissue constituting *obesity*, has already been described under "fatty infiltration." A localized and circumscribed formation, constitutes a *lipoma* or fatty tumor.

STRUCTURE.—The lipomata resemble in their structure adipose tissue. (Fig. 32.) They consist of cells containing fat, and a variable quantity of common connective tissue. The

FIG. 32.



Lipoma.—Some of the cells contain crystallized fatty acids. $\times 200$.

cells, like those of adipose tissue, though usually somewhat larger, are more or less round or polygonal in shape, and are distended with fluid fat. The nucleus and protoplasm are so compressed against the cell-wall by the fluid contents, that although their existence may often be demonstrated by treatment with reagents, they are usually only readily visible when the cell is atrophied and contains less fat. (See Fig. 4 a.) The connective tissue, which varies in amount, usually unites the cells in masses

or lobules of various sizes, and also in most cases forms a thin capsule around the tumor. Bloodvessels are distributed in the fibrous septa.

DEVELOPMENT.—The lipomata grow from connective tissue. Adipose tissue, it must be remembered, is merely connective tissue containing numerous cells which are infiltrated with fat; and its growth consists, either in the infiltration of more of these cells, or in a proliferation of the cells, and an accumulation of fat in those newly developed.

A lipoma in the same way originates by a localized proliferation of cells, which as they are produced become infiltrated with fat. The growth of these tumors is always very slow, and they are usually encapsulated by a layer of fibrous tissue.

SECONDARY CHANGES.—Secondary changes in the lipomata are not common; their fibrous septa may, however, become calcified, or even ossified, and the fatty tissue undergo a process of liquefaction. Soft-

ening may also occur from a mucoid change. Inflammation of these tumors is rare, but when situated in the subcutaneous tissue the skin over them may become adherent and ulcerate.

PHYSICAL CHARACTERS, ETC.—The situation of the lipomata is almost co-extensive with that of adipose and connective tissue. They occur most frequently, however, in those parts in which fat is normally met with, as in the subcutaneous tissue and the inter-muscular septa. They are also occasionally developed in the subsynovial and subserous tissues, in the submucous tissue of the stomach and intestines, and even in internal organs. They sometimes attain an enormous size. They are more or less lobulated, and are usually surrounded by a fibrous capsule which separates them from the adjacent structures. On section they present the ordinary appearance of adipose tissue. Their consistence varies with the amount of fibrous tissue which they contain. They are usually single, although not unfrequently multiple. In their growth they occasionally become pedunculated.

CLINICAL CHARACTERS.—Clinically, the lipomata are perfectly innocent.

CHAPTER XV.


THE ENCHONDROMATA.

THE Enchondromata are tumors histologically resembling cartilage.

STRUCTURE.—Like cartilage they consist of cells and an intercellular substance, which present all the variations observed in the normal tissue. The intercellular substance may be hyaline, faintly or distinctly fibrous, or mucoid. When fibrous, the fibres may be arranged like those of fibro-cartilage, or more or less concentrically around the cells, as in the reticular cartilages of the ear and larynx. (Fig. 33.) When hyaline or mucoid, it is sometimes quite soft in consistence. The cells may be very numerous, or few in proportion to the matrix.

They are round or oval, and occasionally branched and stellate. In the hyaline forms they are usually large and round or oval (Fig. 34);


FIG. 33.



Fibrous Enchondroma.
× 200.

in the fibrous forms they are often smaller and even somewhat spindle-shaped, more resembling those of connective tissue; and in the rarer

FIG. 34.



Hyaline Enchondroma. X200.

mucoid forms, they are more commonly stellate and branched, like those of the umbilical cord. They are either single or arranged in groups, and are usually surrounded by a capsule, as in normal cartilage, although this is often very indistinct. They inclose one or more nuclei and slightly granular contents; sometimes a cell-wall cannot be distinguished. In addition to the intercellular substance, the

growth is usually divided into several lobes by bands of fibrous tissue, in which are contained the bloodvessels. These lobes are often very distinct, so that the growth appears to be made up of several separate tumors. The fibrous tissue in most cases also encapsules the growth and separates it from the surrounding structures, although sometimes this encapsulation is absent, and the tumor is surrounded by a zone of embryonic cells, which infiltrate the adjacent tissues.

DEVELOPMENT.—The enchondromata most frequently originate from bone and common connective tissue, very rarely from cartilage. Cartilage itself, and especially fibrous cartilage, is very closely allied to common connective tissue. It grows from the deeper layers of the perichondrium, which proliferate and form an embryonic tissue; the young cells become cartilage-cells, and these probably form the matrix, which is either homogeneous or fibrillated, constituting in the one case hyaline, and in the other fibrous cartilage. The development of enchondroma from connective tissue is precisely similar to the physiological process.

In the development of enchondroma from osseous tissue, the medulla is the source of the new growth. This proliferates, the osseous trabeculae are absorbed, the neighboring medullary spaces open one into the other, and in this manner a large medullary cavity is produced. In the centre of this, the young cells first formed enlarge and become separated by a homogeneous, or less frequently, slightly fibrillated intercellular substance, and thus is produced a mass of cartilage in the centre of the medullary tissue. This gradually increases till ultimately a layer of fibrous tissue is formed around it, and its further growth takes place from the tissue of its capsule.

Lastly, cartilaginous growths may originate from cartilage itself. These are sometimes seen on the surface of the articular cartilages, in the larynx and trachea, and on the costal and intervertebral cartilages. They are simply local outgrowths from pre-existing cartilage. They rarely attain a large size, and in structure and physical characters more closely resemble normal cartilage than the other forms of enchondroma. They are usually described as *enchondroses*, and must be distinguished from the other forms of cartilaginous tumor.

SECONDARY CHANGES.—Of these, calcification is much the most common. It affects different parts of the growth, commencing in the capsules, and then involving the intercellular substance. Ossification also occasionally occurs; it commences at separate centres, and spiculæ of bone are formed which traverse the tumor in various directions. Fatty degeneration and mucoid softening are common changes, and may lead to the formation of large softened masses which present the appearance of cysts. In rare cases the skin covering the tumor ulcerates, and a fungating mass protrudes.

VARIETIES.—The varieties of enchondroma depend mainly upon the nature of the intercellular substance. There are thus hyaline, fibrous, and mucoid enchondromata; these, however, are usually combined in various degrees in the same tumor. As a rule, those originating from the medulla of bone are of the hyaline and mucoid class, whilst those originating from connective tissue in other situations are more frequently fibrous. The rapidly-growing fibrous forms approach very closely the confines of the sarcomata, the mucoid forms the confines of the myxomata; and these two kinds of growth are often associated in the same tumor.

A variety of enchondroma has been described under the name of *osteo-chondroma*, which in structure more closely resembles bone than cartilage. It consists of a tissue similar to that met with between the periosteum and bone in rickets, which from its resemblance to osseous has been called *osteoid* tissue. This tissue only requires calcifying to become true bone. Like bone it is made up of trabeculæ and medullary spaces, but the trabeculæ, instead of bone-corpuscles and lamellæ, consist of small angular cells without a capsule, situated in an obscurely fibrillated matrix, which in part is calcified. The medullary spaces contain a fibrous stroma and many bloodvessels. The *osteo-chondromata*, although consisting mainly of this osteoid tissue, contain also a small proportion of cartilage. They originate beneath the periosteum, their common seat being the ends of the long bones. Their

growth is very rapid, and they often attain an enormous size. They are much more freely supplied with bloodvessels than the ordinary enchondromata, and hence they are much less frequently the seats of retrogressive changes. They are especially prone to become ossified and converted into true bone.

PHYSICAL CHARACTERS, ETC.—The enchondromata occur most frequently in early life. About three-fourths of them are met with in the osseous system, where they grow either from the medulla or from the periosteum. Their favorite seat is the fingers and toes. The remaining fourth occur most frequently in the parotid gland and in the testicle. They occasionally grow in the intermuscular septa, in the subcutaneous cellular tissue of the mamma, and in the lungs. They are usually single, except when occurring on the fingers and toes, in which situations they are more frequently multiple. They consist of a single tumor, or of several smaller tumors held together by fibrous tissue. The more slowly growing enchondromata are hard, smooth, elastic tumors, often lobulated, and seldom exceeding the size of an orange. Less frequently these tumors grow very rapidly, are quite soft in consistence, and attain a large size.

CLINICAL CHARACTERS.—The enchondromata must for the most part be regarded as innocent growths. They are usually encapsuled, and in most cases produce merely local effects, although these, from the parts involved and the rapidity of growth, are often very injurious. The softer forms, however, and especially those which occur in the medulla of bone and in glands, occasionally exhibit malignant characters. These grow the most rapidly, and are often not limited by a fibrous capsule, but surrounded by a zone of embryonic tissue. Such tumors tend to recur locally after removal, and in rare cases also infect the lymphatic glands, and are reproduced in the lungs. In speaking of the malignancy of these enchondromata, it must be borne in mind that they are sometimes associated with sarcoma, and the malignant properties are probably in all cases to be ascribed to the combination of sarcomatous with the cartilaginous elements.

CHAPTER XVI.

THE OSTEOMATA.

THE Osteomata, or osseous tumors, are tumors consisting of osseous tissue. A new formation of bone occurs under various circumstances. Irritative conditions of the bone and periosteum are often attended by a large formation of new bone. This is seen after fractures, in which there is not only a formation of bone from the bone itself, but also from the periosteum and adjacent fibrous structures ("permanent" and "provisional callus"). Chronic inflammation of the periosteum is also frequently followed by thickening of the bone beneath it. These, however, are inflammatory formations, and have not an independent growth like the osseous tumors.

STRUCTURE.—Osseous tumors resemble in structure normal bone. There are three histological varieties:—

1. *The Eburnated Osteomata*.—These consist of dense, compact, osseous tissue. The lamellæ are arranged concentrically and parallel to the surface of the tumor. There is a complete absence both of bloodvessels and of cancellous tissue.

2. *The Compact Osteomata*.—These are formed of a tissue similar to that of the compact tissue of the long bones; differing only in the arrangement of the Haversian canals and canaliculi, which is less regular than in normal bone.

3. *The Cancellous Osteomata*.—These consist of cancellous osseous tissue, which is usually surrounded by a thin layer of denser bone. The medullary spaces may contain embryonic tissue, a fibrillated tissue, or fat.


DEVELOPMENT.—Osseous tumors originate from bone or its periosteum, from cartilage, and from connective tissue apart from bone.

VARIETIES.—The osteomata are divisible into two classes, according to their seat—the homologous osteomata or *exostoses*, and the heterologous osteomata or *osteophytes*.

The *homologous osteomata* or *exostoses*, are outgrowths from pre-existing bone, growing either from the periosteum, from the articular cartilage, or from the medulla. Those growing from the periosteum

in the fibrous forms they are often smaller and even somewhat spindle-shaped, more resembling those of connective tissue; and in the rarer

FIG. 34.



Hyaline Enchondroma. X200.

mucoïd forms, they are more commonly stellate and branched, like those of the umbilical cord. They are either single or arranged in groups, and are usually surrounded by a capsule, as in normal cartilage, although this is often very indistinct. They inclose one or more nuclei and slightly granular contents; sometimes a cell-wall cannot be distinguished. In addition to the intercellular substance, the

growth is usually divided into several lobes by bands of fibrous tissue, in which are contained the bloodvessels. These lobes are often very distinct, so that the growth appears to be made up of several separate tumors. The fibrous tissue in most cases also encapsules the growth and separates it from the surrounding structures, although sometimes this encapsulation is absent, and the tumor is surrounded by a zone of embryonic cells, which infiltrate the adjacent tissues.

DEVELOPMENT.—The enchondromata most frequently originate from bone and common connective tissue, very rarely from cartilage. Cartilage itself, and especially fibrous cartilage, is very closely allied to common connective tissue. It grows from the deeper layers of the perichondrium, which proliferate and form an embryonic tissue; the young cells become cartilage-cells, and these probably form the matrix, which is either homogeneous or fibrillated, constituting in the one case hyaline, and in the other fibrous cartilage. The development of enchondroma from connective tissue is precisely similar to the physiological process.

In the development of enchondroma from osseous tissue, the medulla is the source of the new growth. This proliferates, the osseous trabeculæ are absorbed, the neighboring medullary spaces open one into the other, and in this manner a large medullary cavity is produced. In the centre of this, the young cells first formed enlarge and become separated by a homogeneous, or less frequently, slightly fibrillated intercellular substance, and thus is produced a mass of cartilage in the centre of the medullary tissue. This gradually increases till ultimately a layer of fibrous tissue is formed around it, and its further growth takes place from the tissue of its capsule.

Lastly, cartilaginous growths may originate from cartilage itself. These are sometimes seen on the surface of the articular cartilages, in the larynx and trachea, and on the costal and intervertebral cartilages. They are simply local outgrowths from pre-existing cartilage. They rarely attain a large size, and in structure and physical characters more closely resemble normal cartilage than the other forms of enchondroma. They are usually described as *enchondroses*, and must be distinguished from the other forms of cartilaginous tumor.

SECONDARY CHANGES.—Of these, calcification is much the most common. It affects different parts of the growth, commencing in the capsules, and then involving the intercellular substance. Ossification also occasionally occurs; it commences at separate centres, and spiculæ of bone are formed which traverse the tumor in various directions. Fatty degeneration and mucoid softening are common changes, and may lead to the formation of large softened masses which present the appearance of cysts. In rare cases the skin covering the tumor ulcerates, and a fungating mass protrudes.

VARIETIES.—The varieties of enchondroma depend mainly upon the nature of the intercellular substance. There are thus hyaline, fibrous, and mucoid enchondromata; these, however, are usually combined in various degrees in the same tumor. As a rule, those originating from the medulla of bone are of the hyaline and mucoid class, whilst those originating from connective tissue in other situations are more frequently fibrous. The rapidly-growing fibrous forms approach very closely the confines of the sarcomata, the mucoid forms the confines of the myxomata; and these two kinds of growth are often associated in the same tumor.

A variety of enchondroma has been described under the name of *osteo-chondroma*, which in structure more closely resembles bone than cartilage. It consists of a tissue similar to that met with between the periosteum and bone in rickets, which from its resemblance to osseous has been called *osteoid* tissue. This tissue only requires calcifying to become true bone. Like bone it is made up of trabeculæ and medullary spaces, but the trabeculæ, instead of bone-corpuscles and lamellæ, consist of small angular cells without a capsule, situated in an obscurely fibrillated matrix, which in part is calcified. The medullary spaces contain a fibrous stroma and many bloodvessels. The osteo-chondromata, although consisting mainly of this osteoid tissue, contain also a small proportion of cartilage. They originate beneath the periosteum, their common seat being the ends of the long bones. Their

amongst it are a varying number of amœboid cells. The bloodvessels, which are not numerous, are readily visible and easily isolated. A few elastic fibres are sometimes seen between the cells.

DEVELOPMENT.—The myxomata always originate from one of the connective tissues. Adipose tissue is their most favorite seat—either the subcutaneous, the submucous, or the inter-muscular adipose tissue. They also grow from the medullary tissue of bone, the connective tissue of organs, from the connective tissue of the brain and spinal cord, and from the sheaths of nerves. They are usually separated from the surrounding structures by a very thin fibrous capsule, fine prolongations from which divide the growth into lobules of various sizes. In exceptional cases they may increase by the continuous invasion of their matrix. Their growth is usually slow, but they may attain an enormous size.

SECONDARY CHANGES.—Of these the most common is rupture of the capillaries, hemorrhage, and the formation of sanguineous cysts; this, however, is less frequent than in the sarcomata. The cells themselves may undergo mucoid or fatty degeneration, and thus be destroyed: this is usually accompanied by liquefaction of the inter-cellular substance.

VARIETIES.—The varieties of myxoma depend principally upon its combination with other growths. The most common is a combination with lipoma. Combinations with sarcoma, enchondroma, and adenoma, are also frequently met with.

PHYSICAL CHARACTERS, ETC.—The myxomata are of a peculiar soft gelatiniform consistence, and of a pale grayish or reddish-white color. Their cut surface yields a tenacious mucilaginous liquid, in which may be seen the cellular elements of the growth. They are most frequently met with in the later periods of life. Growing from the sheaths of nerves, they constitute one variety of so-called neuroma. They may also grow from the placenta, constituting the “uterine hydatids.” When situated in superficial parts they may become pedunculated. In the submucous tissue of the nose they constitute one form of nasal polypus.

CLINICAL CHARACTERS.—Clinically the myxomata are for the most part benign growths. If completely removed they rarely recur. Sometimes, however, they exhibit malignant characters, and recur locally after removal. They probably never reproduce themselves in internal organs. In speaking of their malignancy their occasional association with sarcoma must be borne in mind.

CHAPTER XVIII.


THE LYMPHOMATA.

LYMPHOMA, or Lymphadenoma, is a new formation consisting of lymphatic, or as it is sometimes called, adenoid tissue (the "adenoid tissue" of His). Lymphatic tissue is the tissue composing the follicles of the lymphatic glands and the Malpighian corpuscles of the spleen, and existing in many other parts belonging to the lymphatic system. This tissue is now known to have a much more general distribution than was formerly supposed; it not only constitutes the follicles of the lymphatic glands and the Malpighian corpuscles of the spleen, but also Peyers's glands and the solitary glands of the intestine, the follicles of the pharynx and tonsils, the thymus gland, and the trachoma glands of the conjunctiva. Recently it has also been found to exist in many other situations, as around the bloodvessels of the pia mater and of other parts, in the neighborhood of the smallest bronchi, in the pleura immediately beneath its endothelium, in the peritoneum, in the mucous membrane of the alimentary canal, and in the medulla of bone.

STRUCTURE.—Lymphatic tissue, wherever it exists, possesses the same general structure, and the follicle of a lymphatic gland may be taken as the type, not only of the physiological tissue, but also of the pathological growths.

This tissue consists essentially of a delicate reticulum, within the meshes of which are contained numerous lymphatic elements—the so-called lymph-corpuscles. The reticulum is made up of very fine fibrils which form a close network, the meshes of which are only sufficiently large to inclose a few, or even a single corpuscle in each. The fibrils usually present a more or less homogeneous appearance, and nuclei are often to be distinguished at

FIG. 36.



Lymphoma.—A thin section of a lymphomatous tumor of the mediastinum, from which most of the cells have been removed by pencilling. Showing the reticulated network, and the nuclei in its angles. This network is much coarser than that often met with. $\times 20$.

the angles of the network (Fig. 36). The lymphatic cells, or lymph-corpuscles, which constitute the greater part of the tissue, can in most cases be readily removed from the meshes of the reticulum by the agitation of thin sections in water. They are identical in their characters with the white cells of the blood. As usually seen after death, they are spheroidal, pale, semi-transparent bodies, varying considerably in size, and also presenting slight differences in their structure. Some are granular, and appear to possess no nucleus; in others, a distinct, simple, or compound nucleus is visible, which is usually also granular; others again are much larger, and contain two or even three nuclei (Fig. 37).


The histological characters of the lymphomata, however, vary considerably, according to the age of the growth and the rapidity of its development. In the earlier stages of development, and when growth is taking place very rapidly, the proportion of cells is great, and many of these are much larger than those normally met with in lymphatic glands, containing two, or even more, nuclei. In a more advanced stage, and also in all tumors in which the process of development has been slow, the proportion of cells is smaller, and the reticulum consti-

FIG. 37.



Cells from a Lymphatic Growth in the Liver.—Those to the left are the ordinary lymph-corpuscles which constituted the greater part of the growth. To the right are some of the larger elements. $\times 350$.

FIG. 38.



Lymphoma.—Section of a firm lymphoma of the mediastinum. Showing a very thickened reticulum, within the meshes of which the lymphoid cells are grouped. $\times 200$.

tutes a more prominent part of the growth. (Fig. 38.) The larger cell-forms also are almost entirely wanting; and the reticulum, instead of being exceedingly delicate, is much coarser, and forms a network of broad homogeneous or slightly fibrillated bands. As the reticulum increases the lymph corpuscles gradually diminish in number and become arranged in smaller groups within its meshes. (See Fig. 28.) These variations in the proportion of corpuscles and stroma are precisely analogous to those met with in lymphatic glands as the result

of acute and chronic inflammation. (See "Inflammation of Lymphatic Structures.")

DEVELOPMENT.—The lymphomata originate for the most part from lymphatic tissue, being simply overgrowths of pre-existing lymphatic structures. They are, therefore, usually homologous. They may, however, be heterologous, either owing to the new tissue extending considerably beyond the confines of the old, or to its growth in situations where it is normally almost entirely wanting. This latter condition obtains in Hodgkin's disease, and in certain forms of lymphoma which are malignant.

SECONDARY CHANGES.—The lymphomata do not undergo marked retrograde changes. There is no fatty degeneration, caseation, or softening, such as occurs in scrofulous glands.

PHYSICAL CHARACTERS, ETC.—The physical characters of the lymphomata vary according to the rapidity of their growth. The rapidly-growing forms, in which the cellular elements are so numerous, are of a grayish-white color and soft brain-like consistence, much resembling encephaloid cancer. These often attain an enormous size, and infiltrate the neighboring structures. They have been called by Virchow *lympho-sarcoma*. Those which are more slowly developed, and in which the reticulum constitutes the greater portion of the growth, are much harder in consistence, sometimes being almost cartilaginous. These harder growths rarely attain a large size.

CLINICAL CHARACTERS.—Clinically, the lymphomata are, for the most part, perfectly innocent tumors. They originate most frequently in the lymphatic glands, the gland undergoing a continuous increase in size. Sometimes the enlargement of the glands appears in the first place to be of an inflammatory nature, and to result from some irritation, but upon this being removed the glands, instead of subsiding, continue to increase. In most cases however, no such source of irritation is discoverable. The glands which are especially prone to this disease, are the cervical, the submaxillary, the axillary, the inguinal, the bronchial and mediastinal, and the abdominal glands. Usually only a single gland or a single group of glands is affected; sometimes, however, the growth is more general. As the glands enlarge, they gradually unite, so that ultimately they may form very large lobulated tumors. When occurring in the mediastinum they may invade one or both lungs, and they constitute here the most common form of mediastinal tumor (so-called "Thoracic Cancer"). The lymphatic structures in the intestine may in the same way become enlarged and

project so as to form polypi. The enlargement of the spleen in ague, is also probably of the same nature.

The lymphomata occasionally, however, exhibit malignant properties. This is especially the case in those richly cellular, soft, rapidly-growing forms which are sometimes met with. Such growths may rapidly infiltrate the surrounding structures, involve the neighboring lymphatic glands, and even infect distant parts. It is to these malignant forms that the term *lymphadenoma* is commonly applied. They probably correspond with Virchow's lympho-sarcoma.

In the condition known as "Hodgkin's Disease," and in Leukæmia, lymphomatous growths are met with in various parts of the body.

Hodgkin's Disease.

This disease is characterized by the enlargement of the lymphatic glands in various parts of the body, together with the development of lymphatic growths in internal organs, especially in the spleen and liver. The new growths are precisely similar, histologically, to lymphoma.

The disease was formerly described by Hodgkin, Bright, Wilks, and Trousseau, and was called, after the first-named of these observers, "Hodgkin's Disease." Trousseau designated it "Adénie." It is also known as "Anæmia Lymphatica." Recently its characteristics have been more fully determined by MM. Cornil and Ranvier, and by Drs. Murchison¹ and Sanderson, to whose descriptions the reader is referred for more complete information respecting it.

Histologically, the new growths are precisely similar to those met with in leukæmia. The disease differs essentially, however, from leukæmia in this respect, that the new formation of lymphatic tissue is not associated with any notable increase in the number of the white corpuscles in the blood.

The lymphatic glands are usually the earliest seats of the new growth, and here it differs in no respect histologically from the lymphomata. At first it may be limited to a single group of glands; subsequently, however, the process becomes more general, and the glands throughout the whole body may be more or less involved. The glands as they increase in size gradually become confluent, so as to form large lobulated tumors, and the new growth, which takes place

¹ "Trans. Path. Soc. Lond.," vols. xx. and xxi.


in the earlier stages within their capsules, often ultimately extends so as to infiltrate the surrounding structures.

This new growth of lymphatic tissue, which commences in and often extends beyond the confines of the lymphatic glands, is ultimately followed by the formation of lymphatic growths in various internal organs, but more especially in the spleen and liver. In the spleen, the Malpighian corpuscles become enlarged and form grayish-white nodules and masses. The kidneys, lungs, stomach, muscle, bones, and subcutaneous tissue may all become involved, the new growths occurring either as nodules of various sizes scattered through the organs, or in a more infiltrated form, like many of those met with in leukæmia.

In their physical characters the new growths vary. They are sometimes of a grayish-white color and of a soft brain-like consistence, yielding a milky juice which is rich in cellular elements precisely resembling leucocytes. In other cases they are much firmer in consistence, and they may be as hard as an indurated lymphatic gland. These differences depend upon the relative proportions of the cells and stroma. They have but little tendency to undergo retrogressive changes.

Leukæmia.

This disease, like Hodgkin's disease, is characterized by the development of lymphomatous tissue in various organs, but especially in the spleen. The growth of this new tissue, however, is associated with a large increase in the number of the white corpuscles in the blood. It is this alteration in the blood which gives leukæmia its distinctive characters—hence its name. The disease will be considered subsequently, when treating of "Diseases of the Blood."



CHAPTER XIX.

THE MYOMATA, NEUROMATA, AND ANGIOMATA.

The Myomata.

THE myomata are tumors consisting of muscular tissue. A new formation of muscle has been already described as being frequently associated with the ordinary process of hypertrophy, both of striated and of non-striated muscle—a simple hyperplasia of the elements of the muscle accompanying the increase in their size. (See “Hypertrophy.”)

STRUCTURE.—The myomata consist either of striated or of non-striated muscle. The former are exceedingly rare, only two or three examples having been recorded, and these were congenital.

The myomata of *non-striated* muscle consist, like the physiological tissue, of elongated spindle-cells with rod-shaped nuclei, more or less isolated or grouped into fasciculi of various sizes, with a varying quantity of connective tissue. The connective tissue is often exceedingly abundant, especially in old tumors, so much so, that these growths in the uterus—where they most frequently occur—have commonly been known as “*fibroid*” tumors. The muscular elements either present a more or less regular arrangement, or pass in all directions through the tumor. The bloodvessels, which are usually not numerous, are distributed in the connective tissue.

DEVELOPMENT.—The myomata probably always originate from muscle, they are, therefore, homologous growths. They may form distinctly circumscribed tumors surrounded by a fibrous capsule, or constitute ill-defined irregular masses in the midst of the muscular tissue in which they grow. They not infrequently become pedunculated and form polypi, especially in the uterus.

SECONDARY CHANGES.—Of these, the most frequent is calcification. Hemorrhage, mucoid softening, and the formation of cysts, are also occasionally met with.

Myomata are most frequently met with in the uterus. They may also occur in the prostate, in the œsophagus, and in the stomach and intestines.

Myoma of Uterus.—The uterus is by far the most frequent seat of myomata, and here they constitute the so-called “uterine fibroid.” In most of these muscular tumors of the uterus there is a large proportion of connective tissue—hence the term “fibroid.” This is especially the case in older growths. Those newly developed, however, consist almost entirely of true muscular tissue. They either form firm hard masses, imbedded in the uterine walls, or project into the uterine or abdominal cavities. When projecting into the uterus they constitute a common form of uterine polypus. These tumors are often multiple. Their growth is slow. The older ones are liable to become calcified. They also sometimes undergo mucoid softening which gives rise to the formation of cysts in the tumor.

CLINICAL CHARACTERS.—Clinically, the myomata are perfectly innocent.

The Neuromata.

The neuromata are tumors consisting almost entirely of nerve-tissue. The term “neuroma” has been applied to many growths found in connection with nerves: fibrous, myxomatous, and gummy tumors growing within the nerve-sheath have been included under this head. True neuroma, however, is rarely met with, and is amongst the least frequent of all the new formations.

STRUCTURE.—The neuromata most commonly consist of ordinary medullated nerve-fibres; they therefore resemble in structure the cerebro-spinal nerves, from which they most frequently grow. The nerve-fibres are associated with more or less connective tissue. Virchow has also described, as exceedingly rare formations, tumors composed of non-medullated fibres, and of ganglionic nerve-tissue.

DEVELOPMENT.—The neuromata always originate from pre-existing nerve-tissue,—either from the cranial or from the spinal nerves. Their growth is slow, they rarely attain a large size, but usually exist as small single nodules.

The most frequent seat of these growths is the extremities of divided nerves, where they are sometimes found after amputations. They exist, in this situation, as spherical or elongated enlargements of the divided extremity of the nerve; and they are usually intimately connected with the cicatricial tissue of the stump, from which they can only with difficulty be isolated. They may also occur in the course of the nerves in any situation, either as single or multiple nodules.

CLINICAL CHARACTERS.—Clinically, the neuromata are perfectly innocent tumors. They often cause considerable pain.

The Angiomata.

The angiomata, or vascular tumors, are tumors consisting of blood-vessels held together by a small amount of connective tissue. They include the various forms of nævi, the erectile tumors, and aneurism by anastomosis. They may be divided into two classes—the *simple* angiomata, in which the new vessels resemble normal arteries, veins, or capillaries; and the *cavernous* angiomata, in which the blood circulates in a cavernous structure similar to that of the corpus cavernosum penis.

SIMPLE ANGIOMATA.—These include the various forms of nævi, and telangiectasis. They consist of tortuous and dilated bloodvessels, held together by a small quantity of connective and adipose tissue. The vessels are most of them of new formation; some, however, may be the original vessels of the part which have become considerably enlarged. They most commonly partake of the nature of capillaries, but in other cases the arterial or venous characters predominate. These growths are usually small, superficial, slightly elevated masses; although they sometimes form larger tumors. Their color is red, violet, or purple, according to the character of the blood which they contain. The first-named is much the most frequent.

CAVERNOUS ANGIOMATA.—These include the venous vascular tumors, erectile tumors, and aneurism by anastomosis. They consist of an erectile cavernous tissue, closely resembling that of the corpus cavernosum penis. The growth is made up of irregular fibrous alveoli, which communicate freely with one another, and are lined with an endothelium similar to that of the veins. These spaces are distended with blood, usually venous, which is supplied to them by numerous tortuous vessels, and circulates in them with varying degrees of rapidity. These growths are commonly of a bluish color. They may be diffuse or form distinctly circumscribed tumors. They often exhibit distinct pulsation. Their favorite seat is the skin and subcutaneous tissue. They may also occur in the orbit, in muscle, and in the liver, spleen and kidneys.


CHAPTER XX.

THE PAPILOMATA.

THE Papillomata are new formations resembling in structure ordinary papillæ, and like these they grow from cutaneous, mucous, or serous surfaces.

STRUCTURE.—They consist of a basis of, often richly cellular, connective tissue supporting bloodvessels, which terminate in a capillary network or in a single capillary loop, the whole being enveloped in a covering of epithelium. (Fig. 39.) The epithelial covering varies in

FIG. 39.



Papilloma.—Showing a single enlarged papilla. (Rindfleisch.)

character in different growths. In those of the skin, it is often very abundant, and the superficial layers are hard and stratified, forming a dense firm covering. In those originating from mucous surfaces, the epithelium forms a thinner investment, and is of a much softer consistence; whilst in those growing from serous membranes it often constitutes only a single layer.

The growth may be simple—consisting merely of enlarged papillæ,

as in a common wart; or it may be more complex, the papillæ being very numerous, and giving off secondary and tertiary offsets. If the investing epithelium be very abundant, it may so inclose the whole mass as to give to it a more or less regular outline. More commonly, however, this is not the case, and the epithelium not being sufficient to fill up the spaces between the papillæ, the growth presents a branched, villous, or cauliflower appearance. The bloodvessels are often very numerous, and are sometimes dilated and tortuous.

DEVELOPMENT.—The papillomata always originate from the skin, from mucous, or from serous membranes. They most frequently grow from pre-existing papillæ; sometimes, however, they occur where no papillæ exist, springing directly from the sub-epithelial connective tissue: this is the case in the stomach and larynx. Their growth is usually slow. The individual tumors rarely attain a very large size, the larger forms being for the most part constituted of several smaller growths.

SECONDARY CHANGES.—Of these, ulceration and hemorrhage are the most frequent. They occur especially in those growths which originate from mucous surfaces. The hemorrhage is often very abundant, and may even endanger life. This is not unfrequently the case in the papillary growths of the bladder and intestine.

VARIETIES.—The varieties of papillary tumors depend principally upon their seat. Those growing from the skin include *warts* and *horny growths*. Warts are firm, have a dense epithelial covering, and are less prone to ulceration and hemorrhage than those growing upon other parts. Horny growths appear usually to originate in the sebaceous follicles, by a continuous proliferation of their epithelium. The epithelium, together with the sebaceous secretion, forms a projecting horn, which increases by growth at its base. Such formations hardly come within the definition of papilloma. Larger and more vascular papillary tumors may, however, occur on cutaneous surfaces—such are the *condylomata* and *venereal warts* met with around the anus and upon the external male and female genital organs as the result of irritating secretions.

The papillomata of mucous membranes are softer and more vascular than the preceding, they have a less dense epithelial covering, and are more prone to ulceration and hemorrhage. Many of them come within the category of mucous polypi. They are met with on the tongue, in the larynx and nose, on the gastro-intestinal mucous membrane, on the cervix uteri, and in the bladder. In the bladder and intestine

they are often exceedingly vascular, and give rise to profuse hemorrhage. Here they are not unfrequently confounded with villous epithelioma.

Papillomata of serous membranes never form distinct tumors. They are met with most frequently as small outgrowths from the synovial membrane in chronic diseases of joints.

CLINICAL CHARACTERS.—Clinically, the papillomata are innocent growths. They may, however, prove fatal from continuous ulceration and hemorrhage: this is especially the case, as already mentioned, in papilloma of the bladder and intestine. In these situations they are easily mistaken for epithelioma: the symptoms of both are very similar, and it is often only after death that they can be distinguished. In the papillomata the epithelium is *homologous*, being situated only upon the surface of the papillæ, and in no case growing *within* their connective tissue basis. In the epitheliomata, on the other hand, it is *heterologous*, and it is met with at the base of the tumor in the subjacent connective tissue. (See Fig. 60.) It is important to remember that a growth which is primarily a simple papilloma may subsequently become an epithelioma. (See “Epithelioma.”)

CHAPTER XXI.

THE ADENOMATA.


THE Adenomata—or as they are more commonly called, *glandular tumors*—are new formations of gland-tissue.

STRUCTURE.—In structure the adenomata resemble the racemose or tubular glands.

The racemose adenomata consist of numerous small saccules or acini, lined with small epithelial cells, which are often two or three layers deep. The acini communicate with each other, and are grouped together, being merely separated by connective tissue, in which are contained the bloodvessels. (Fig. 40.) The connective tissue varies in amount. It may resemble the normal tissue, or if growing rapidly it will be much more richly cellular. It sometimes contains spindle elements.

Many new growths, however, contain a large amount of this gland-tissue which are not strictly adenomata. All growths originating in glandular organs may be associated with more or less glandular structure. In the mamma, for example, sarcoma, myxoma, and other forms

FIG. 40.



Adenoma of the Mamma. $\times 300$. (Rindfleisch.)

of tumor, are often so intermingled with the gland-tissue of the organ that it becomes difficult to say which is the predominant structure. In many cases it is evident that the development of such tumors is accompanied by an increase of the gland-tissue amongst which they grow. Thus are produced mixed forms—*adeno-sarcoma*, *adenomyxoma*, etc.

The tubular adenomata grow from mucous membranes, and consist of groups of tubules lined with epithelium. They will be again alluded to hereafter.

DEVELOPMENT.—The adenomata always originate from pre-existing gland-structures, of which they are simply local hyperplasias. Their growth, which is usually slow, takes place by the development of diverticula from the saccules or tubules of the gland, and by a proliferation of the inclosed epithelium. The new growth may remain in intimate relation with the adjacent gland, or it may ultimately become separated from it by a fibrous capsule.


SECONDARY CHANGES.—The most frequent of these is fatty degeneration of the epithelium, which may give rise to the formation of small caseous masses in the growth. Dilatation of the saccules and tubules so as to form cysts, and mucoid softening, are also common.

VARIETIES.—*Adenoma of Mamma.*—The mamma is much the most common seat of adenoma. Here two varieties must be distinguished—one, in which there is a general hyperplasia of the whole gland; the

other, in which the process is limited to smaller or larger groups of lobules. The former constitutes hypertrophy of the mamma, the latter is the *chronic mammary* or *adenoid* tumor. Pure adenoma of the mamma, however, is not common. (See Fig. 40.) Many tumors in this situation described as adenomata are fibrous, sarcomatous, or myxomatous growths in which is contained a varying amount of gland-tissue. The distinction between such growths and pure adenoma is often difficult. The adenomata are either superficially or deeply seated in the gland, from which they are usually separated by a loose fibrous capsule. They are commonly round or oval in shape, lobular, and of a hard elastic consistence. On section they often present a lobulated appearance, their racemose structure being sometimes visible to the naked eye.

Adeno-Fibroma and Adeno-Sarcoma of Mamma.—These are more common than pure adenoma. In them the connective tissue between

FIG. 41.



Adeno-Fibroma of Mamma.—Showing new growth of gland-structure and of connective tissue. $\times 100$; reduced $\frac{1}{2}$.

the acini of the gland undergoes active changes, and forms a fibro-cellular, spindle-celled, or round-celled structure, amongst which the acini are embedded. (Fig. 41.) In many cases it is difficult to determine whether the glandular structure is really increased in amount,

whilst in others it is so abundant that such is evidently the case. These tumors form lobulated masses of a firm consistence. They grow slowly, and rarely ulcerate. In most cases they contain cysts. Many of these are lined with cylindrical epithelium, and are formed by the dilatation of the ducts of the gland; others appear to originate from localized softenings of the tissue. These cysts are usually very numerous, and they often attain a large size. They are frequently seen much flattened, so as to present the appearance of fissures running amongst the growth. In other cases they are more or less completely filled by masses of the tumor which have grown into their interior. This formation of cysts in these tumors has given rise to the terms *cystic-sarcoma*, *cystic-adenoma*, etc.

Adenoma of Mucous Membranes.—The glandular structures of mucous membrane are also common seats of adenoid growths, and here they are usually of the tubular variety. In the nose, the pharynx, the stomach, the intestines, the vagina and uterus, these growths are sometimes met with. In course of time they usually gradually project above the surface of the membrane so as to form a polypus, and thus constitute one of the forms of *mucous polypi*. In consistence they are soft and somewhat gelatinous, and often present a semitranslucent appearance. Their surface resembles in color the surrounding mucous membrane. The formation in them of cysts, by the dilatation of their tubules, is exceedingly frequent: the cysts usually contain a soft mucoid substance. Adenoma of mucous membranes often become cancerous. (See “Cylindrical Epithelioma,” or “Adenoid Cancer.”)

CLINICAL CHARACTERS.—Clinically, the adenomata are perfectly innocent; they are, however, very liable to be confounded with growths possessing malignant properties. A tumor, also, which is primarily a simple adenoma, may subsequently become cancerous. The anatomical distinction between cancer of a gland in its earlier stages and a simple glandular tumor is often exceedingly difficult, especially in the mamma and mucous membranes. In cancer the growth appears to commence by a proliferation of the epithelium within the ducts of the gland, and as the epithelium only subsequently becomes heteroplastic, the determination of the nature of the tumor in this stage is necessarily attended with considerable difficulty. This will be again alluded to when speaking of cancer. (See “Structure of Carcinoma.”) In sarcomatous tumors, again, originating in the connective tissue of a gland, the ducts of the gland, filled with epithelium, are often seen imbedded in the new growth, and thus the appearance of adenoma may be closely simulated.

CHAPTER XXII.

THE SARCOMATA.


THE sarcomata are tumors consisting of embryonic connective tissue. Of these there are several varieties, depending upon the size and configuration of the cells, and the nature of the intercellular substance. They include what have generally been known in this country as *fibro-plastic*, *fibro-nucleated*, *recurrent-fibroid*, and *myeloid* tumors. Many growths formerly described as "cancers" also belong to this class of new formations.

Connective tissue in its embryonic condition is an immature tissue in a state of rapid development. In its most immature state it differs from the fully developed tissue in consisting almost entirely of small round cells, whilst its intercellular substance, instead of being fibrous, is soft and amorphous. This is the common condition of connective tissue in the primary stages of all rapid formative processes, as already described when speaking of it as the tissue from which many tumors of the connective-tissue class originate. (See "Development of New Formations.")

In the process of development of this embryonic into mature connective tissue, the cells diminish in number, many of them assume a spindle shape, and the intercellular substance fibrillates. Similar changes are seen in inflammatory conditions of connective tissue. Here also many of the small round cells which constitute the "granulation tissue" become spindle cells, and the granulation tissue ultimately develops into the fibrous tissue of the cicatrix. In the sarcomata, however, the connective tissue retains the embryonic state throughout its growth, there is a progressive formation of embryonic tissue; and although the process of development may occasionally proceed in certain parts of the tumor to the formation of a more highly developed structure, as fibrous tissue, cartilage, or bone, so that a mixed form of tumor is produced, it usually ceases at the embryonic stage.

STRUCTURE.—The sarcomata may thus be defined as tumors consisting of connective tissue which throughout its growth retains the

embryonic type. The *cells*, which constitute nearly the whole of the growth, consist for the most part of masses of nucleated protoplasm, and rarely possess a limiting membrane. They present many variations in size and form; as a rule, however, they preserve the same general characters in the same tumor. There are three principal varieties—the *round*, the *fusiform*, and the *myeloid* cells.



Cells from a Spindle-celled Sarcoma. $\times 350$.

The *round* cells are many of them indistinguishable from lymph-cells or white blood-corpuscles. Others are somewhat larger, and contain an indistinct nucleus with one or more bright nucleoli: these more closely resemble the cells of a granulation.

The *fusiform*, or *spindle-shaped* cells, are the so-called “fibro-plastic cells.” (Fig. 42.)

They are long narrow cells, terminating at each end in a fine prolongation. Some of them may be broader, approaching the epithelial type; others more or less stellate. They are sometimes slightly granular, and they inclose a long oval nucleus, with or without nucleoli. In size they vary considerably. These cells represent a higher state of development than the round cells, resembling those met with in embryonic tissue which is in the process of forming mature connective tissue. (See Fig. 84 B.)

The *myeloid*, or mother cells, are much larger than either of the preceding, and are analogous to the cells met with in the medulla of foetal bone. (See Fig. 50.) They are large irregular-shaped masses of nucleated protoplasm, for the most part more or less spherical, and often possessing numerous offshoots. They are finely granular, and contain several round or roundly-oval nuclei, each with one or more bright nucleoli. The nuclei may be exceedingly numerous, one cell containing as many as thirty. Both the cells and nuclei vary considerably in size.

An *intercellular substance* exists in all the sarcomata, although it is usually small in quantity, the cells lying in nearly close apposition. It may be perfectly fluid and homogeneous, or firmer and granular, or more or less fibrillated. Chemically it yields albumen, gelatin, or mucin.

The *bloodvessels* are usually very numerous, and are either in direct contact with the cells, or separated from them by a little fibrillated tissue. Their distribution is very irregular, and their walls often con-

sist of embryonic tissue similar to that of the growth which they supply ; hence the frequency with which rupture and extravasation of blood take place.

DEVELOPMENT.—The sarcomata always originate from connective tissue—either from the subcutaneous, the submucous, or the subserous tissues, the fasciæ, the connective tissue of organs, the periosteum, or the medullary tissue of bones. Their growth may take place in two ways, by the multiplication of their own elements—*central* growth, and by the continuous invasion of their matrix—*peripheral* growth. A peripheral growth is the great characteristic of the sarcomata ; they usually increase by the continuous invasion of their connective-tissue matrix, so that no line of demarcation exists between the two. They frequently also invade other tissues, the elements of the growth extending for some distance into the surrounding structures. This infiltrating tendency of the sarcomata varies considerably in the different varieties, being much more marked in the round-celled than in the spindle-celled and myeloid growths. A purely central growth is less common. A sarcomatous tumor, however, often becomes encapsuled, and growth takes place within the capsule ; but even in this case the capsule is sometimes merely that of the part within which the growth originates, as the periosteum, or the capsule of a lymphatic gland.

SECONDARY CHANGES.—The most important of these is fatty degeneration. This always occurs to a greater or less extent in the older portions of the growth, causing softening, or the production of cyst-like cavities. It is frequently associated with destruction of the blood-vessels and hemorrhage : the latter may give rise to the formation of sanguineous cysts. (See “Blood-Cysts.”) Calcification, ossification, and mucoid degeneration are less common. The occurrence of calcification, ossification, and pigmentation is influenced by the predisposition of the matrix from which the growth is produced : thus calcification and ossification are more prone to occur in tumors originating in connection with bone, pigmentation in those originating from the cutis or eyeball.

VARIETIES.—Although all the sarcomata possess the same general characters, they present many histological and clinical differences, which may serve as bases for their classification. The occurrence of various secondary changes—pigmentation, mucoid degeneration, and the formation of cysts, impart their respective characters to the growth ; hence *melanotic-sarcoma* and *cystic-sarcoma* have been described as

distinct varieties. This is to a certain extent justifiable, inasmuch as sarcomata which have undergone these transformations, in many cases possess the property of reproducing the same characters when they occur secondarily in other parts. Then again, as already stated, sarcomatous tumors are sometimes complex in their structure, and are associated with other tissues belonging to the connective-tissue group. A combination of sarcoma with fatty, cartilaginous, osseous, and mucous tissue, is thus not uncommonly met with. This is owing to the embryonic tissue exhibiting a tendency to develop into the different varieties of connective tissue exhibiting a tendency to develop into the different varieties of connective tissue. (See "The New Formations.") The mixed forms—*chondro-sarcoma*, *osteo-sarcoma*, *myxo-sarcoma*, etc., are thus produced. The following histological classification, based upon the three different forms of cells already described, is the one generally adopted. It must, however, be borne in mind, that all the varieties of cells may be found in the same tumor, although the majority are usually of the same type; hence, the *majority* will determine the class to which the growth belongs.

Spindle-celled Sarcoma.

These tumors, which include the growths described by Paget in this country as "fibro-plastic," and "recurrent fibroid," are the most common of all the sarcomata. They consist mainly of spindle-shaped and fusiform cells, lying nearly in close contact, with a little homogeneous or slightly fibrillated intercellular substance. The cells, which contain well-marked oval nuclei, with one or more nucleoli, are parallel to one another, and are arranged in bundles which pass in all directions through the growth, often giving to it a somewhat fibrous appearance. In those portions of the section in which the bundles of

spindle-elements have been cut transversely, they present the appearance of round cells. The cells vary considerably in size in different tumors, hence the division into *small* and *large* spindle-celled growths.

Small Spindle-celled Sarcoma.—In these the cellular elements are small, often not more than $\frac{1}{800}$ th inch in length, and the intercellular substance is occasionally imperfectly fibrillated. (Fig. 43.) These growths approach therefore

FIG. 43.




Small Spindle-celled Sarcoma.—From a tumor of the leg. $\times 200$.

the confines of the fibromata, and histologically they must be regarded as occupying an intermediate place between embryonic and fully-developed connective tissue. They grow from the periosteum, the fasciæ, and from connective tissue in other parts. They are usually tolerably firm in consistence, of a whitish or pinkish-white color, and for the most part present, on section, a translucent somewhat fibrillated appearance. They are often encapsuled, much more frequently so than the other varieties of sarcoma, but they are very liable to infiltrate the surrounding structures, and to recur locally after removal.

Large Spindle-celled Sarcoma.—The cellular elements in these tumors are much larger than in the preceding. The cells are plumper, and the nuclei and nucleoli are especially prominent, and frequently multiple. (Fig. 44.) The intercellular substance is more scanty, and there is a complete absence of any fibrillation. These growths are much softer in consistence than the small-celled variety. They are of a pinkish-white color, and are often stained by extravasations of blood, and sometimes in parts are almost diffuent from extensive fatty degeneration. They grow rapidly, and are usually exceedingly malignant.

FIG. 44.




Large Spindle-celled Sarcoma.—To the left—the cells have been separated by teasing, so that their individual forms are apparent; to the right—they are in their natural state of apposition, such as would be seen in a thin section of the tumor. (Virchow.)

MELANOTIC SARCOMA.—This is a variety of sarcoma in which many of the cells contain granules of dark-colored pigment. By far the greater number of melanotic tumors are sarcomata, and most of the growths which were formerly described as “melanotic cancers,” belong in reality to this class of new formations. Not only are these melanotic tumors most frequently sarcomata, but the majority of them consist mainly of spindle-shaped cells—hence they are described in the present section.

The melanotic sarcomata originate principally in two situations—in the choroid coat of the eye, and in the superficial integuments. In both of these situations pigment is a normal constituent of the tissues, and this tendency of structures normally containing pigment to originate melanotic growths, is exceedingly characteristic. (See “Pigmentary Degeneration.”) These tumors usually consist of spindle-

shaped cells, although in some cases the prevailing type of their elements is round or oval. (Fig. 45.) The pigment, which gives to them their distinctive characters, consists of granules of a brownish or dark sepia color, which are distributed within the cells. (Fig. 45 c.) Frequently, only a very small proportion of the cells are pig-

FIG. 45.



A Melanotic Sarcoma of the Penis.—A. A thin section, showing the general arrangement of the elements. $\times 200$. B. A section from the peripheral part of the growth, showing the “indifferent cells,” amongst which are small isolated pigmented elements. At *a*, a bloodvessel is seen. $\times 200$. C. Some of the elements separated by teasing. In these the pigment-granules are well seen. $\times 400$.

mented, whilst in other tumors the pigmentation is much more universal; in all cases, however, a large number of the elements will be found to be quite free from pigment.


These melanotic tumors are amongst the most malignant of the sarcomatous growths. Although they have comparatively but little tendency to extend locally, they are disseminated by means of the bloodvessels, and occasionally also by the lymphatics, and thus reproduce themselves often very rapidly in distant tissues. In doing so, although they almost invariably maintain their melanotic characters, the degree of the pigmentation of the secondary tumors varies considerably. Whilst many of them may be perfectly black in color, others may be much paler, and perhaps only partially streaked with pigment. The secondary growths are soft, usually distinctly circumscribed, and often encapsuled. They may occur in almost every organ of the body—the liver, the spleen, the kidneys, the lungs, the heart, the brain and spinal cord, and also the lymphatic glands and sub-

cutaneous tissue, may all be simultaneously involved. I have observed, that when occurring in internal organs, the pigmentation is not always limited to the secondary nodules, but that many of the cells proper to the organ itself are filled with granules of similar pigment, which is most abundant in those cells which are immediately adjacent to the new growth. This pigmentation of the cells of the organ often extends for some distance beyond the confines of the tumor.

OSTEOID SARCOMA.—This which is often known as “osteoid cancer” is a variety of sarcoma (usually of spindle-celled sarcoma) in which the growth is either more or less calcified, or has partially become converted into true bone. As a primary growth it is met with almost exclusively in connection with bone, growing either from the periosteum or the medulla. The osteoid characters are usually reproduced in the secondary tumors occurring in the lungs and in other parts.

Simple calcification is much more common than true ossification. Here the growth merely becomes infiltrated with calcareous salts, which may be dissolved out with a little dilute hydrochloric acid, when the characteristic sarcomatous structure becomes revealed. In other cases this calcification is associated with the conversion of portions of the tumor into true, although usually imperfect bone. When this has occurred, there will be seen amongst the spindle-shaped, round or myeloid elements, tracts of tissue consisting of broad bands of granular intercellular material, infiltrated with calcareous salts, inclosing

FIG. 46.



Osteoid Sarcoma.—From a secondary tumor of the lung. Showing the calcification of a spindle-celled growth, and the formation of broad bands of calcified intercellular material inclosing spaces which contain round and oval cells. $\times 200$.

spaces which contain spindle, oval, or round cells. (Fig. 46.) This structure often much resembles cartilage. In other parts of the growth, where the change is more advanced, may be seen lacunæ and even canaliculi such as are found in true bone.


In these osteoid growths it is most important to recognize the exist-

ence of the sarcomatous element, inasmuch as it is the presence or absence of this which determines the innocent or malignant nature of the growth. Osteoid sarcoma must be carefully distinguished from the simple osseous tumor.

Round-celled Sarcoma.

This is of softer consistence than the spindle-celled growths, and from

FIG. 47.



Round-celled Sarcoma.—A thin section of a small round-celled sarcoma of the liver. $\times 200$.


its frequent resemblance in physical characters to encephaloid, it is sometimes known as “medullary,” “encephaloid,” or “soft” sarcoma. Histologically, it is elementary embryonic tissue, consisting mainly of the round cells already described, imbedded in a scanty, and usually soft, homogeneous, or finely granular intercellular substance. (Fig. 47.) The cells usually resemble those met with in the most elementary embryonic tissue; less frequently they are larger, and contain large round or oval nuclei, with bright nucleoli. There is an almost complete absence of fusiform cells, and of the partial fibrillation which is so frequent in the more highly developed spindle-celled variety.

The round-celled sarcomata are of a uniform soft, brain-like consistence, somewhat translucent or opaque, and of a grayish or reddish-white color. On scraping the cut surface, they yield a juice which is rich in cells. They are exceedingly vascular, the vessels often being dilated and varicose, and from their liability to rupture, they frequently give rise to ecchymoses and to the formation of sanguineous cysts. (See “Blood Cysts.”) They grow from the cutis, the subcutaneous cellular tissue, the periosteum, the fasciæ, and from the connective tissue of organs. They extend rapidly by peripheral growth, infiltrate the surrounding structures, reproduce themselves in internal organs, and often involve the lymphatic glands. From their clinical and physical characters, these tumors are very liable to be confounded with encephaloid cancer: they are distinguished by the absence of an alveolar stroma, and by the uniformity in the character of their cells.

GLIOMA.—This is a variety of round-celled sarcoma growing from the neuroglia or connective tissue of nerve. It consists of very small round cells, imbedded in an exceedingly scanty, homogeneous, granu-

lar, or slightly fibrillated intercellular substance. (Fig. 48 *a*.) Some of the cells may possess fine prolongations which, by communicating with one another, form a somewhat reticulated structure. These tumors occur in the gray and white substance of the brain, in the

FIG. 48.



Sarcomatous Tumors from the Brain.—*a*. A glioma of cerebellum. This represents the appearance ordinarily presented by these growths. *b*. A comparatively rare form of sarcoma, which consists of large nucleated cells inclosed within the meshes of a vascular network. The development of this tumor took place in the brain subsequently to that of spindle-celled growths—primarily in the thigh, and secondarily in the lung. $\times 200$.

cranial nerves, and in the retina. In the retina they usually commence as a minute nodule, which may gradually increase until it projects as a large fungating tumor from the orbit. They are not encapsuled, and although they may occasionally infiltrate the tissues in which they lie and cause secondary growths in their immediate vicinity, they very rarely reproduce themselves in the lymphatic glands or in internal organs. They are liable to small hemorrhages into their structure, and sometimes become more or less caseous.

ALVEOLAR SARCOMA.—This is a rare form of round-celled sarcoma, which was first described by Billroth. The cells, which are large, sharply defined, round or oval in shape, and inclose prominent round nuclei, are separated from each other by a more or less marked fibrous stroma. In some parts this stroma forms small alveoli within which the cells are grouped; but careful examination will always show that in most parts of the section the stroma really penetrates between each individual cell. It is this last-named character which serves to distinguish these tumors from the cancers, with which, in many cases, they may easily be confounded. The accompanying drawing, made from a preparation kindly lent to me by Mr. R. J. Godlee, shows well their microscopical characters. (Fig. 49.)

FIG. 49.



Alveolar Sarcoma.—From tumor of the skin. $\times 200$.

Alveolar sarcomata are met with principally in the skin, bones, and muscles. In the skin, where they are often multiple, they lead to ulceration. They tend to recur locally, and also to reproduce themselves in internal organs.

Myeloid Sarcoma.

This, which is the well-known *myeloid* tumor, is somewhat allied to the spindle-celled growths. It possesses, however, certain histological peculiarities which probably depend upon the characters of the tissue

FIG. 50.



Myeloid Sarcoma. (Virchow.)

from which it grows. Myeloid tumors nearly always occur in connection with bone, and most frequently originate in the medullary cavity. They consist of the large, many-nucleated cells already described as “myeloid cells,” which resemble the cells of the medulla in a state of excessive nutritive activity, together with numerous fusiform cells like those met with in the spindle-celled varieties. There are also some smaller round and oval elements. The large myeloid cells, which give

to these tumors their distinctive character, are usually much more numerous in those growths which originate in the medullary cavity than in those which spring from the periosteum. These various forms of cells are nearly in close contact, there being very little intercellular substance. (Fig. 50.) The growths are sometimes very vascular, so much so as to give rise to distinct pulsation. They often contain cysts.

Myeloid tumors almost always grow in connection with bone, the heads of the long bones being their favorite seat. They are also frequently met with springing from the periosteum of the upper and lower jaws, where they constitute one form of *epulis*. When originating within the medullary cavity, the compact tissue of the bone becomes expanded over them, and they thus often communicate on palpation the peculiar sensation known to surgeons as "egg-shell crackling." These tumors are for the most part of firmer consistence than the other varieties of sarcoma; many of them are firm and fleshy, although others are softer, more resembling size-gelatin. They are not pulpy and grumous like the soft sarcomata, neither do they present the fasciculated appearance of the spindle-celled varieties. Their cut surface has a uniform succulent appearance, often mottled with patches of red. They are often encapsuled by the periosteal covering of the bone from which they grow. They are rare after middle life, and are the least malignant of all the sarcomata.

Psammoma.

Allusion must here be made to a rare form of growth occasionally met with in the brain and its membranes which is known as psammoma. This growth, although having but little resemblance to the sarcomata, is usually classified with this group of new formations. Its characteristic feature is that it consists largely of calcareous particles. The calcareous particles are contained in the concentric bodies already described as the corpora amylacea, where they give rise to the so-called "brain sand": hence the name of the growth. The calcified corpora amylacea, associated with a varying quantity of a cellular and fibrillated tissue, and bloodvessels, make up the growth.

Psammoma is usually met with growing from the membranes of the brain, or from the choroid plexus. In the latter situation it often contains numerous cysts. It is of no pathological importance except when of sufficiently large size to produce symptoms from pressure.

Blood-cysts.

Tumors are occasionally met with into which so much hemorrhage has taken place as to mask their real nature, and to give to them the appearance of blood-cysts. The nature of these blood-cysts has only recently been understood. They are now known to be in the majority of cases soft, round, or spindle-celled sarcomata. They consist of broken-down blood coagula, surrounded by an ill-defined layer of soft sarcoma tissue. The microscope will also usually reveal sarcomatous elements amongst the altered blood. These growths are exceedingly malignant, and hence the recognition of their sarcomatous origin is all important.

Clinical Characters of the Sarcomata.

The sarcomata occur most frequently in early and middle life, and, next to the cancers, are the most malignant of the new formations. They are especially characterized by their great tendency to extend locally and to infiltrate the surrounding structures, so that they are exceedingly prone to recur *in loco* after removal. They comparatively rarely infect the lymphatic glands, and in this respect present a marked contrast to the cancers. They are also very liable to become generally disseminated, although this is not usual in the earlier stages of the disease. The secondary growths occur most frequently in the lungs. The dissemination is effected by means of the blood, and this is owing to the thinness of the walls of their bloodvessels and to the immediate contact of these with the cells of the growth—conditions most favorable to the entrance of the cellular elements into the circulation. The dissemination of the sarcomata is, on this account, sometimes more rapid than that of the cancers. In the latter, extension in the early stage takes place principally by the lymphatics, and dissemination by the blood only occurs later in the disease. The secondary sarcomata usually resemble the primary one, but in exceptional cases the several varieties may replace one another.

These malignant properties, as has been seen, are possessed by the different varieties of sarcoma in very different degrees. As a rule, the softer and more vascular the tumor, and the less its tendency to form a fully developed tissue, the greater is its malignancy. The soft, round-celled, and large spindle-celled varieties are thus usually much

more malignant than the firmer, small spindle-celled growths. Their infiltrating powers are much greater, they sometimes infect the lymphatic glands, and they tend to reproduce themselves very rapidly in internal organs. Many of the small spindle-celled tumors, after removal, never recur, whilst others recur locally several times, and ultimately reproduce themselves in distant parts. As a rule, largeness of the spindle elements and the existence in many of them of more than one nucleus, is an evidence of special malignancy. The presence of a capsule limiting the growth must also be taken into account in judging of the degree of its malignancy. It must, however, be borne in mind that in a growth distinctly encapsuled the sarcomatous elements may invade the adjacent structures. The myeloid growths are the least malignant; they may however also, in exceptional cases, give rise to secondary growths in internal organs.

CHAPTER XXIII.

THE CARCINOMATA.

THE Carcinomata, or Cancers, are new formations consisting of cells of an epithelial type, without any intercellular substance, grouped together irregularly within the alveoli of a more or less dense fibroid stroma.


The term "*cancer*" has been so commonly applied indefinitely to any growth possessing malignant properties, that "*cancerous*" and "*malignant*" have come to be regarded by many as synonymous terms. It is important, however, clearly to distinguish between them. A *cancer* is a growth possessing the above-named definite structure; a *malignant* growth, on the other hand, is one which, independently of its structure, tends to reproduce itself in adjacent or distant tissues. (See "Malignancy.") "*Cancerous*" is an *anatomical* term; "*malignant*" is a *clinical* one.

The Cancers include the four following varieties: *Schirrus*, *Encephaloid*, *Colloid*, and *Epithelioma*. Of these, the first three possess the same general characters, although they present certain structural

and clinical differences which serve to distinguish them. Epithelioma constitutes a more distinct variety. It will be well, in the first place, to describe the characters common to the larger and more important group, and then those which are peculiar to its individual members. Epithelioma will be considered subsequently.

STRUCTURE.—In structure, scirrhus, encephaloid, and colloid cancer so far resemble one another, that they all consist of cells of an epithelial type, without any intercellular substance, grouped together irregularly within the alveoli of a fibrous stroma. (See Fig. 54.) Although there is no intercellular substance, a certain amount of liquid exists between the cells. It is this liquid which exudes from the freshly-cut surface of the cancer, and the number of cells which it contains give to it a milky appearance.

The *cells* are characterized by their large size, by the diversity of their forms, and by the magnitude and prominence of their nuclei and nucleoli. (Fig. 51.) In size they vary from $\frac{1}{800}$ to $\frac{1}{100}$ of an inch in diameter; the majority being about five times as large as a red blood-corpuscle. They are round, oval, fusi-form, caudate, polygonal—exhibiting, in short, every diversity of outline. These variations in form are principally owing to the mutual pressure to which in their growth they are subjected. The nuclei, which are large and prominent, are round or oval in shape, and contain one or more bright nucleoli. The nuclei are,



Cells from a Scirrhus of the Mamma. $\times 350$.

perhaps, most frequently single; two, however, are frequently met with, and in the softer and more rapidly growing cancers they may be more numerous. The cells rapidly undergo retrogressive changes, hence they usually contain molecular fat. They are many of them exceedingly destructible, so that sometimes more free nuclei than cells are visible. Cells precisely similar to these are met with in other morbid growths, and also in the normal tissues. There is thus no *specific* "cancer-cell." It is the general character of the cells, together with their mode of distribution in the meshes of a fibroid stroma, that determines the nature of the growth to which they belong. The appearance presented by these cells grouped within the alveoli of the cancer sometimes closely simulates, in the earlier stages of growth, that of simple adenoma. (See Fig. 40.) In adenoma, however, the cells resemble the epithelium of the gland; they are smaller, and less

irregular in size and shape, and less closely packed than those of carcinoma. In many cases, indeed, they form merely a single layer lining the walls of the acini.

The *stroma* varies considerably in amount, being much more abundant in some varieties of cancer than in others. It consists of a more or less distinctly fibrillated tissue, arranged so as to form alveoli of various forms and sizes, within which the cells are grouped. (Fig. 52.) These alveoli communicate with one another, so as to form a continuous cavernous system. The characters of the stroma vary with the rapidity of its growth:—if this is rapid it will contain some round and spindle-shaped cells (see Fig. 56); if, on the other hand, it is slow or has altogether ceased, the tissue will contain but few cells, and will be denser and more fibrous in character. The latter is the condition in which it is most commonly met with.

FIG. 52.



The Alveolar Stroma from a Scirrhus of the Mamma. The cells have been removed by pencilling. $\times 200$.

Within the stroma are contained the *bloodvessels*. These are often very numerous, and form a close network. They are limited to the stroma, and only in very exceptional cases do they encroach upon the alveoli.¹ This distribution of the bloodvessels is important, as distinguishing the cancers from the sarcomata. In the latter, the vessels are not supported by a stroma, but ramify amongst the cells of the growth; hence the facility with which these tumors become generally disseminated.

In addition to the bloodvessels the cancers also possess *lymphatics*. These accompany the bloodvessels, and, as has been shown by MM. Cornil and Ranvier, communicate freely with the alveoli. This explains the great tendency of cancer to infect the lymphatic glands.

DEVELOPMENT.—The question of the genesis of cancer involves that of the genesis of epithelium generally. It is maintained by many Histologists that epithelium can only originate from epithelium, and that the strata of cells set aside in the embryo for the production of

¹ In soft, rapidly growing cancers the bloodvessels have occasionally been observed to project as tufts into the alveoli. See case reported by Mr. Marcus Beck and Mr. Arnott, "Trans. Path. Soc. Lond.," 1874, p. 224.

the epithelial tissues is the source from which all epithelium is subsequently derived. Others admit that epithelium may also originate from connective tissue. (See "Development of the New Formations.") A like difference of opinion exists as to the source of the epithelioid cells of cancer. By many—as Waldeyer, Thiersch, and Billroth—they are regarded as originating only from pre-existing epithelium. Others—amongst whom are Virchow, Lucke, Rindfleisch, and Klebs—maintain that they may also be derived from cells belonging to the connective tissue. It is also believed by some—as Köster—that many cancers originate from the endothelium of the lymphatics.

The difficulty of determining the genesis of cancer is partly owing to the fact that it usually originates in structures where epithelium is normally abundant, as in the mamma, skin, and alimentary canal; and that this normal epithelium is always, from the earliest stage of the growth, the seat of active proliferation. In cancer of the mamma, for example, the first change often observable is an accumulation of epithelium within the ducts of the gland, a condition very similar to that already described as occurring in the development of an ordinary glandular tumor. (See "Adenoma.") Soon, however, the epithelium alters its characters. The cells become larger and more irregular in shape, and their nuclei are more prominent. The epithelium is then found outside the ducts, amongst the inter- and peri-glandular connective tissue, which is also in a state of active proliferation, and is infiltrated with small round cells. The outline of the ducts ultimately becomes completely annihilated, and the epithelioid cells are seen in alveoli formed of a fibroid tissue. The question arises as to whether the epithelioid cells, which constitute the ultimate cancerous growth, originated from the epithelium normally existing within the ducts of the gland, which in the process of development has extended beyond the ducts into the surrounding tissue; or, whether they originated in the connective tissue—the proliferation of the glandular epithelium being merely a secondary process, and resulting from the irritation of the inter-glandular growth. Although this question cannot be regarded as finally settled, the former view is certainly the more probable one, and the doctrine of the epithelial origin of cancer is now steadily gaining ground. The tendency which is exhibited by the cells of cancerous growths to maintain the type of the epithelial structures in the vicinity of which they grow, is greatly in favor of this view that they are derived from the epithelium. In cancers situated

near the cutaneous surfaces, for example, the cells are usually of the squamous type, whereas in those growing in connection with the glands of mucous membranes, they more commonly resemble the epithelium of the gland. In adopting this doctrine of the origin of cancer it is obviously impossible to admit the occurrence of primary growths in situations where no epithelium normally exists—as in bone and muscle; and we now know that tumors in these tissues which were formerly described as cancers, are in reality sarcomata. With regard to the new formation of the epithelial elements, the researches of Dr. Creighton, which show that this takes place principally by a process of vacuolation, and endogenous growth, has been already alluded to. (See “Development of New Formations.”) -

The *stroma* of cancer, although partly the pre-existing connective tissue of the part in which the cancer originates, is in the main a new formation, and it may probably be regarded as the result of the irritation of the connective tissue by the infiltrating epithelial cells.

The cancers in their growth never become encapsuled, but gradually infiltrate the surrounding structures. This process of infiltration is very characteristic, and is more marked in cancer than in any of the malignant growths. A zone of small-celled infiltration is seen for some distance around the confines of the tumor, so that there is no line of demarcation between it and the normal structures. (See Fig. 53.)

SECONDARY CHANGES.—The most important of these is fatty degeneration. This always occurs to a greater or less extent in all the varieties of cancer. The more rapid the growth, the earlier does this retrogressive change take place, and the greater is its extent; hence it is usually most marked in encephaloid. It produces softening of the growth, which is often reduced to a pulpy cream-like consistence. Pigmentation, mucoid and colloid degeneration may also occur. Calcification is very rarely met with.

VARIETIES.—The term “cancer” was so vaguely applied by the older pathologists—nearly all malignant formations being included under this head—that considerable confusion has resulted in the classification of cancerous growths. Peculiarities in situation, structure, and appearance have given rise to special names, hence the terms—“*osteoid*,” “*chondroid*,” “*cystic*,” “*villous*,” and “*hæmatoid*” cancer. A cancer containing large quantities of pigment was described as a distinct variety, under the name of *melanotic* cancer. Melanotic cancer, however, is comparatively rare. The majority of tumors which

are thus designated are in reality sarcomata. (See "Melanotic Sarcoma.")


The most convenient classification, and that which is now generally adopted, divides the cancers into four groups: *scirrhous*, *fibrous*, or *chronic* cancer; *encephaloid*, *medullary*, or *acute* cancer; *colloid* or *gelatiniform* cancer; and *epithelial* cancer, or *epithelioma*, including *adenoid* cancer. This division is based principally upon the relative proportion of the stroma, and upon the type of the epithelial elements.

Scirrhous Cancer.

Scirrhous, *fibrous*, or *chronic* cancer is characterized by the large amount of its stroma and by the chronicity of its growth. The slowness in the development of scirrhous probably accounts in great measure for the peculiarities in its structure and physical characters.

The epithelial growth, although at first it may be luxuriant, quickly subsides. The elements soon atrophy and undergo retrogressive

Fig. 53.



Scirrhous of the Mamma.—A thin section from the most external portion of the tumor, showing the small-celled infiltration ("indifferent tissue") of the muscular fibres and adipose tissue in the neighborhood of the gland. $\times 200$.


changes. They are most abundant in the external portions of the tumor, where growth is taking place; in the central portions they may be almost entirely wanting. The accompanying figures (Figs. 53 and 54) show the appearances presented by scirrhous of the mamma in the earlier stages of its development.

The degeneration of the epithelial elements is probably owing

to the excessive growth of the stroma, and to the subsequent induration and contraction which it undergoes. It quickly assumes the characters of cicatricial tissue, and becomes hard and indurated. This causes obstruction and obliteration of the bloodvessels which it contains, and it is probably to this interference with the vascular supply that the arrest in the development of the cancer is owing. The whole of the central portions of the growth may thus ultimately consist simply of dense fibroid tissue, amongst which are contained atrophied epithelial cells and fatty debris (Fig. 55), the periphery being the only part where the epithelial structure is visible. The amount of atrophy and contraction varies considerably in different cases.

The physical characters of scirrhus are in the same way due to the abundance of its stroma. The growth is firm and hard, and it is


FIG. 54.



Scirrhus of the Mamma.—A portion of the tumor somewhat internal to that represented in Fig. 53, showing the characteristic alveolar structure of the cancer $\times 200$.

usually depressed in the centre, owing to the contraction of the fibroid tissue; this is very characteristic of scirrhus of the breast, where it causes puckering of the superjacent structures. On section the tumor presents a grayish-white glistening surface, sometimes intersected with

FIG. 55.



Scirrhus of the Mamma.—A section from the more central portions of the tumor, showing the atrophy of the epithelial cells, the diminution in the size of the alveoli, the fibroid tissue, and the fatty debris. *a*, earlier stage; *b*, more advanced. $\times 200$.

fibrous bands. The more external are less firm than the central portions of the growth, and yield, on scraping, a juice which is rich in nucleated cells, free nuclei, and granules.

Scirrhus is most commonly met with in the female breast, and in the alimentary canal—especially in the œsophagus, pylorus, and rectum. It also occasionally occurs in the skin. The secondary growths to which it gives rise are often encephaloid.

Encephaloid Cancer.

Encephaloid, *medullary*, or *acute* cancer, is very closely allied to the preceding, from which it differs merely in the greater rapidity of its growth, and the consequent small amount of its stroma, and the softness of its consistence. Encephaloid and scirrhus cannot be regarded

as in any way constituting distinct varieties of cancer. There are all intermediate stages between them, and the differences in the rapidity of their growth, and consequently in their structure and physical characters, constitute their only distinctive features.

The epithelial growth in encephaloid is rapid and abundant, and the cells, for the most part larger than in scirrhus, quickly undergo fatty degeneration, so that often more free nuclei than cells are visible. The proportion of stroma is very small, and owing to the rapidity of its growth, it is much less fibrous than that of scirrhus, and does not undergo a similar cicatricial contraction. (Fig. 56.) The bloodvessels are often very abundant, and the tissue supporting them being soft and non-resisiant, hemorrhage occasionally takes place.

FIG. 56.



Encephaloid Cancer.—From a secondary cancer of the liver. Showing the large size of the alveoli and the thinness of their walls. In the latter small cells are visible. The large epithelial cells are commencing to undergo fatty metamorphosis. $\times 200$.

Encephaloid cancer is of a soft brain-like consistence, the central portions, where fatty degeneration is most advanced, often being completely diffuent. The tumor is sometimes more or less lobulated. On section, it presents a white pulpy mass, much resembling brain-substance, which is often irregularly stained with extravasated blood.


Encephaloid is much less common than scirrhus cancer. It is most frequently met with in internal organs as a *secondary* growth. It also sometimes occurs primarily in the testis and mamma. Many growths formerly described as encephaloid cancer, are soft sarcomata. (See "Round-celled Sarcoma.")

Colloid Cancer.

The growths described under the name of *colloid*, *alveolar*, or *gelatiniform* cancer, although sometimes regarded as constituting a distinct variety of cancer, are simply one of the preceding forms which have undergone a mucoid or colloid change. The frequency with which non-cancerous growths which have undergone these forms of degeneration have been confounded with colloid cancer has already been alluded to. (See "Colloid Degeneration.")

The alveolar structure in colloid cancers is very marked. The alveoli have very thin walls; they are large, distinct, and more or less spherical in shape. This large size and distinctness of the alveoli is owing to their distension with the softened substance. Within them is contained the gelatinous colloid material, which is a glistening, translucent, colorless, or yellowish substance, of the consistence of thin mucilage or size-gelatin. In the main it is perfectly structureless; within it, however, are imbedded a varying number of epithelial cells, which also contain the same gelatinous substance. (Fig. 57.) These cells present a peculiar appearance: they are large and spherical in shape, and are distended with drops of the same gelatinous material as that in which they are imbedded. (See Fig. 57.) Many of them

FIG. 57.



Colloid Cancer.—Showing the large alveoli, within which is contained the gelatinous colloid material. $\times 300$. (Rindfleisch.)

display a lamellar surface, their boundary being marked by concentric lines. It would appear that the colloid change commences in the cells, which become gradually destroyed in the process. In other cases the cells, with the exception of slight fatty metamorphosis, are but little affected, and the substance distending the alveoli is more viscid and mucoid in character. This is due to a mucoid degeneration of the intercellular substance, rather than to a colloid change commencing in the cells. (See "Mucoid Degeneration.")

Colloid cancer is most frequently met with in the stomach, in the intestine, in the omentum, and in the peritoneum.

Epithelioma.

Epithelioma, or *epithelial cancer*, must be regarded as constituting a much more distinct variety of cancer than either of the preceding, although transitional forms between it and scirrhus are occasionally met with. It differs from the other varieties of cancer in always growing in connection with a cutaneous or mucous surface—the junction of the two being its usual seat—and in its epithelial elements closely resembling the squamous variety of epithelium.

The cells of epithelioma are in the main indistinguishable from those met with on the cutaneous surfaces, and on the mucous membrane of




the mouth. They vary in size from $\frac{1}{32}$ to $\frac{1}{16}$ of an inch in diameter, the average being $\frac{1}{16}$. They contain usually a single nucleus; frequently, however, the nuclei are multiple. (Fig. 58.) They are often considerably flattened and distorted in shape, owing to the pressure to which in their growth they are subjected, but they never present those numerous varieties in outline which are met with in the other varieties of cancer, neither do they exhibit the same marked

tendency to undergo fatty degeneration. The arrangement of these cells is peculiar:—some of them are situated in irregular tubular-shaped lobules which communicate with each other; others are less regularly grouped in masses of various sizes amongst the meshes of a stroma. As the cells increase in number they tend to become arranged concentrically in groups so as to form globular masses. These masses are the “*concentric globes*,” or “*epithelial nests*,” which are so commonly met with wherever squamous epithelium is undergoing rapid growth, and which, although not distinctive, are exceedingly characteristic of epithelioma. As the epithelium multiplies, the peripheral layers of cells become flattened by pressure against the surrounding structures, whilst those in the centre remain more or less spherical in shape like those of the deeper layers of the epidermis. (Fig. 59.) The cells may be so closely packed as ultimately to become hard and dry like those of the nails and hair, and the globes are then of a brownish-yellow color and of a firm consistence. These globes are often large enough to be readily visible to the naked eye,

and owing to the arrangement of the epidermic scales, they usually present a fibrous appearance.

The stroma presents every variation between rapidly growing embryonic, and an incompletely fibrillated tissue. It may be tolerably

FIG. 59.




Epithelioma of the Ltp.—Showing the concentric globes of epithelial cells. $\times 100$.

abundant, or almost entirely wanting. It rarely forms such a marked alveolar structure as that which characterizes the other varieties of cancer, but usually consists simply of a small-celled infiltration surrounding the epithelial elements, which may ultimately become developed into a more or less completely fibrillated tissue. (Fig. 60.)

With regard to the development of epithelioma, there can be no doubt that its epithelial elements are derived from the epithelium of the skin or mucous membranes, or from that of the glands which are situated in these tissues. The growth commences by a proliferation of this epithelium, which as it increases becomes *heterologous*, extending beyond the normal limits into the subjacent connective tissue, and even into muscle, bone, and other structures (Fig. 60); and it is this heterologous development of epithelium which is the essential characteristic of epithelioma. This extension of the epithelium into the subjacent connective tissue produces in the latter an irritative growth, and thus the groups of epithelial elements are always surrounded by

a small-celled ("indifferent") tissue, the small-celled growth preceding the epithelial invasion. (Fig. 60.)

FIG. 60.



Epithelioma of the Tongue.—A thin vertical section, showing the excessive epithelial growth upon the surface of the papillæ, and the extension of the epithelial elements into the subjacent connective tissue. The sub-epithelial tissue is infiltrated with small ("indifferent") cells, amongst which are seen the epithelial elements both single and forming concentric globes. $\times 100$.

Epithelioma usually presents itself in the first place either as a small foul ulcer with indurated edges, or as a subcutaneous induration or nodule which subsequently ulcerates. The surface of the ulcer is frequently papillated and villous, owing to the irregular growth of the corium. The tumor itself is firm in consistence, often more or less friable, and on section presents a grayish-white granular surface, sometimes intersected with lines of fibrous tissue. The cut surface yields on pressure a small quantity of turbid liquid, and in many cases also a peculiar, thick, crumbling, curdy material can be expressed, which often comes out in a worm-like shape, like the sebaceous matter from

the glands of the skin. This latter is very characteristic. It is composed of epithelial scales, and on being mixed with water it does not diffuse itself like the juice of other cancers, but separates into minute visible particles. If it is very abundant, the cancer is soft and friable, and the material can be seen in the cut surface as small scattered opaque dots.

Epithelioma has its primary seat in the immediate vicinity of the cutaneous or mucous surfaces, the point of junction of the two being its favorite habitat. It appears in most cases to owe its origin to some external source of irritation. It is rare in the young, and is most frequently met with in the lower lip at the junction of the skin and mucous membrane, on the tongue, prepuce, scrotum ("chimney-sweep's cancer"), labia, eyelids, cheeks, and in the uterus and bladder. As it extends it may involve any tissue—muscle, bone, and tendon may be alike implicated. It usually infects the lymphatic glands, but very rarely occurs in internal organs.

Cylindrical Epithelioma or Adenoid Cancer.—These terms are applied to those forms of epithelial cancer which grow from mucous membranes with columnar (cylindrical) epithelium, as from those of the stomach and intestines. In these tumors the epithelial elements are similar to those of the mucous membrane from which they grow. They are cylindrical in shape, and are arranged perpendicularly to the walls of the alveoli in a manner precisely analogous to that of the columnar epithelium on the mucous surface. There is rarely a formation of concentric globes, and the growths are of a soft, and often gelatinous consistence. These tumors cause secondary growths in the lymphatic glands, and sometimes in the liver and lungs, which possess the same characters as the primary cancer. The distinction between them and simple adenomata is often exceedingly difficult. (See "Adenoma of Mucous Membranes.") *Guichon 794.*

CLINICAL CHARACTERS OF THE CANCERS.—In speaking of the clinical characters of the cancers, it is important in the first place to make a distinction between epithelioma and the other varieties. Epithelioma, as far as its malignancy is concerned, occupies a very inferior position to scirrhus, encephaloid, and colloid. These latter varieties of cancer possess in the highest degree malignant properties. They extend locally, invading indiscriminately the tissues amongst which they grow, and reproduce themselves in the lymphatic glands and in

internal organs. In the process of dissemination, however, they present some peculiarities which distinguish them from growths which are sometimes equally malignant—viz., the sarcomata. The cancers are characterized by their great tendency to reproduce themselves in the neighboring lymphatic glands. This implication of the lymphatics is usually much more marked than in the sarcomata, in which it but comparatively rarely occurs, and this is probably owing to the communication of the lymphatic vessels with the alveolar spaces of the cancerous growth. The general dissemination in internal organs, on the other hand, is often effected much less readily in cancer than in sarcoma, and the course of the former is therefore sometimes more protracted than that of the latter. This difference is explained by the difference in the distribution of the bloodvessels: in cancer, these are contained in the stroma, and very rarely come into contact with the cells of the growth; whereas, in the sarcomata, they ramify amongst the cells, and their walls being composed of thin embryonic tissue like that of the growth which they supply, dissemination through the medium of the blood is rapidly and readily effected. In cancer, the lymph being so important a medium of infection, the reproduction of the growths in internal organs may be considerably delayed: the progress of the disease becomes arrested by the lymphatic glands, and its further dissemination is often only effected after these have become very generally and extensively involved.

With regard to the difference in the clinical characters of these three varieties of cancer—the dissemination of encephaloid takes place much more rapidly than that of scirrhus, owing to the greater rapidity of its growth, its greater vascularity, and the greater activity of its epithelial elements. Colloid is somewhat inferior in the degree of its malignancy to both scirrhus and encephaloid.

Epithelioma is of all the cancers much the least malignant. It extends locally, and may infect the neighboring lymphatics, but it comparatively rarely reproduces itself in internal organs. This is probably owing to the size and character of the epithelial elements, which render them much less liable to be transmitted by the blood and lymph-streams than are the cells of the other varieties of cancer.

In all the varieties of cancer there is a tendency for the secondary growths to repeat the characters of the primary one. This is most marked in epithelioma. In scirrhus, the secondary growths in internal organs, although sometimes resembling the primary tumor, are often

more rapidly developed, are softer and more vascular, and in accordance with the distinction which has been made between scirrhus and encephaloid, they must be regarded as belonging to the latter variety of cancer.

CHAPTER XXIV.

CYSTS.

IN addition to the new growths already described, there is a large class of formations, many of which cannot be regarded as "tumors" in the strict application of this term. These are the *cysts* or *cystic tumors*.

A *cyst* is a cavity containing liquid or pultaceous material, which is separated from the surrounding structures by a more or less distinct capsule. It may be a new formation, or a pre existing structure which has become distended by its own secretion, or by extravasation into it. The former, only, comes within the category of new growths, although, for the sake of convenience, it will be advisable to consider them both under one head.

There are thus two principal modes by which cysts originate—one, the most frequent, by the gradual accumulation of substances within the cavities of pre-existing structures, which are, for the most part, products of their own formation, being in some cases a secretion, and in others a cell-growth; the other, by the independent formation of a cyst in the tissues.

The accumulation of secretions and of other products within pre-existing cavities, may be effected in the three following ways:—

1st. By the retention of the normal secretion owing to the closure of the excretory ducts—as so often occurs in sebaceous glands.

2d. By excessive secretion, the cavity being unprovided with an excretory duct—as in the formation of bursæ.

3d. By the extravasation of blood into the cavity—as in hæmatocele.

The independent formation of a cyst may take place—

1st. By the softening and liquefaction of the tissues in some particular part, owing to mucoid or fatty changes. The tissues around

the softened matters become condensed, and ultimately form a kind of cyst-wall.

2d. By the enlargement and fusion of the spaces in connective tissue, and the accumulation of fluids within them. The surrounding tissue becomes condensed, and forms a cyst-wall; and this may, in some cases, become lined with secreting cells.

3d. By the formation of a cyst-wall around foreign bodies, parasites, or extravasated blood.

STRUCTURE.—The wall of the cyst will vary in its nature according as it is a pre-existing or a newly-formed tissue. In the former case, it will possess an epithelial lining which will present the same characters as that of the gland, serous membrane, or other structure, from which the cyst originated. If the cyst is a new growth, it rarely possesses an epithelial lining, but consists simply of a fibrous capsule. The cyst-wall is sometimes firmly connected with the adjacent parts, so that it can only with difficulty be separated; in other cases, the union is much less intimate. Instead of being a distinct structure, it may be simply the surrounding tissue which has become dense and fibrous in character.

The contents of cysts are very various, and may serve as a basis for their classification. In the retention-cysts, they will vary with the nature of the normal secretion—serum, sebaceous matter, saliva, milk, seminal fluid, and other substances are thus found in these cysts, more or less altered in character from being retained in a closed cavity. In the exudation-cysts, serum is the most frequent constituent; and in extravasation-cysts, blood. In those cysts which originate from the softening and breaking down of tissue, the contents are the products of retrogressive tissue-metamorphosis, and usually consist largely of mucin, fatty matters, and serum.

SECONDARY CHANGES.—These may take place in the wall of the cyst, or in its contents. The cyst-wall itself may become the seat of new growths, and produce secondary cysts, villous, glandular, and other structures:—this occurs in many compound ovarian cysts. It may also be the seat of an inflammatory process, which terminates in suppuration and granulation, and by this means the cyst frequently becomes obliterated, its contents being either absorbed or discharged externally, and the cavity closing by granulation. Calcification and ossification of the wall may also occur. The contents of cysts undergo various changes, owing to their retention in a closed cavity. The secretions become altered in character, thickened and viscid.

Epithelial elements undergo fatty changes, and so give rise to cholesterolin crystals. Calcification of the contents is also common.

Cysts may be *simple* or *compound*. A simple cyst consists of a single loculus. A compound or multilocular cyst is one consisting of numerous loculi, which either communicate with one another, or remain isolated. Another variety of compound cyst, consists of a cyst with endogeneous growths, the larger cyst having others growing from its walls. A compound cyst may become a simple one by the destruction of its walls.

Cysts are frequently associated with other growths, hence the terms—"cystic-sarcoma," "cystic-cancer," etc. It is especially in those growths which originate in glandular structures, as in the mamma, testicle, and ovary, that this combination is met with. The cystic development may almost entirely obliterate the structure of the tumor in which it takes place, so that ultimately the latter becomes converted into a combination of cysts. In other cases large portions of the tumor grow into the cystic cavities. Considerable difficulty is thus not unfrequently caused in determining the nature of the original growth.

CLASSIFICATION.—Cysts may be most conveniently classified according to their mode of origin, thus:—

CLASSIFICATION OF CYSTS.

I. *Cysts formed by the accumulation of substances within the cavities of pre-existing structures.*

A. RETENTION CYSTS.—Cysts resulting from the retention of normal secretions. These include—

- a. *Sebaceous Cysts*.—These are formed by the retention of secretions in the sebaceous glands. Such are comedones and atheromatous tumors.
- β. *Mucous Cysts*.—These are formed by the retention of secretions in the glands of mucous membranes.
- γ. *Cysts from the retention of secretions in other parts*, including—Ranula, from occlusion of the salivary ducts; Encysted Hydrocele, from occlusion of the tubuli testis; cysts in the mammary gland, from obstruction of the lacteal ducts; simple and some compound cysts of the ovary, from dilatation of the Graafian follicles; and simple cysts of the liver and kidneys.

- B. EXUDATION CYSTS.—Cysts resulting from excessive secretion in cavities uprovided with an excretory duct. These include Bursæ, Ganglia, Hydrocele, and many cysts in the broad ligament.
 - C. EXTRAVASATION CYSTS.—Cysts resulting from extravasation into closed cavities. These include Hæmatocele, and some other forms of sanguineous cysts.
- II. *Cysts of independent origin.*
- A. CYSTS FROM SOFTENING OF TISSUES.—These are especially common in new formations, as in enchondroma, lipoma, sarcoma, etc.
 - B. CYSTS FROM EXPANSION AND FUSION OF SPACES IN CONNECTIVE TISSUE.—These include—
 - α. *Bursæ*, originating from irritation and exudation into the tissues.
 - β. *Serous cysts in the neck* (often congenital).
 - γ. *Many compound ovarian cysts*.¹
 - C. CYSTS FORMED AROUND FOREIGN BODIES, EXTRAVASATED BLOOD, AND PARASITES.
 - D. CONGENITAL CYSTS.—These include many Dermoid cysts. These appear often to be the remains of blighted ova. They contain fatty matters, hair, teeth, bones, etc.

γ²

CHAPTER XXV.

INFLAMMATION.

THE morbid processes which have thus far been described have been mainly characterized by some alteration in the nutrition of the histological elements of the tissues; either by a diminution in their nutritive activity, as in atrophy and the degenerations, or by an increase, as in hypertrophy and the new formations. In the process of *inflammation* an alteration in nutrition also plays a prominent part, but

¹ See Dr. Wilson Fox, on Cystic Tumors of the Ovary, "Med. Chir. Soc. Trans.," vol. xlvii.

changes in the bloodvessels and in the circulation are its essential and most important constituents.

Inflammation may be defined to be the succession of changes which takes place in a living tissue as the result of some kind of injury, provided that this injury be insufficient immediately to destroy its vitality. With regard to the nature of the injury, it may consist in some direct damage to the tissue, either by mechanical or chemical agents, or by substances conveyed to it by means of the bloodvessels or lymphatics: or the injury may be indirect, as in some cases of inflammation of internal organs arising from exposure to cold. In all cases, however, some injury of the tissue—an injury which impairs, and if of sufficient intensity would destroy, its vitality—precedes the occurrence of the local changes which constitute the inflammatory process.

The exact nature of these changes has, for the most part, been ascertained during the past ten years, mainly owing to the experimental researches of Professors Cohnheim, Stricker, and Burdon Sanderson. The method of investigation has consisted in the artificial production of inflammation in the lower animals, and the observation of the process as thus induced. The process comprises—

1st. *Changes in the bloodvessels and circulation.*

2d. *Exudations of liquor sanguinis and migration of blood-corpuscles ; and*

3d. *Alterations in the nutrition of the inflamed tissue.*

It will be well, in the first place, to consider each of these separately, in the order in which they occur, and subsequently to endeavor to point out how far a causal relation exists between them.

I. CHANGES IN THE BLOODVESSELS AND CIRCULATION.—Changes in the bloodvessels and circulation, resulting in increased vascularity, have ever been regarded as playing a most important part in inflammation, as upon them principally depend those signs of the process which are most obvious during life. The redness, heat, and swelling, which are so constantly met with in inflamed tissues, are in great measure due to the attendant hyperæmiâ. The swelling, however, is in most cases dependent rather upon the effusion than upon the overfulness of the bloodvessels.

These changes in the bloodvessels and circulation are essential constituents of inflammation, both in vascular and in non-vascular tissues. In the latter, which comprise the cornea and cartilage, they take place in the adjacent vessels from which these tissues derive their nutritive supply. The nature of these vascular changes has been studied by

the artificial production of inflammation in transparent tissues, in which the circulation can be readily observed: the web, mesentery, and tongue of the frog, and the wing of the bat, being most convenient for this purpose. The phenomena, as observed in the mesentery of the frog, which has been previously curarized, may be thus briefly described:—

The first effect of injury of the mesentery—mere exposure to the air being sufficient for the purpose—is to cause *dilatation* of the arteries, and after some interval, a similar dilatation of the veins and capillaries. The dilatation of the arteries commences at once, and is not preceded by any contraction. It gradually increases for about twelve hours, and is accompanied also by an increase in the length of the vessels, so that they become more or less tortuous. This enlargement of the bloodvessels is associated at the commencement of the process with an *acceleration* in the flow of blood; this, however, is soon followed by a considerable *retardation* in the circulation, the vessels still remaining dilated. These alterations in the rapidity of the blood-flow cannot be owing to the increase in the calibre of the vessels, which remain throughout dilated.

It has, however, long been known that the acceleration of the blood-flow in an injured part—the so-called *determination* of blood, which was so correctly described more than thirty years ago by Dr. C. J. B. Williams—is not necessarily followed by retardation. It may gradually subside without retardation or any of the essential phenomena of inflammation taking place. Cohnheim has consequently stated in his more recent researches on inflammation, that the dilatation of the vessels and the increased velocity of the blood-current which ensue immediately after the infliction of the injury are only temporary and accidental. They may even, in some cases, be followed by contraction before the permanent dilatation commences. The permanent dilatation and diminished velocity, on the other hand, come on slowly and are permanent, and these must be regarded as the proper vascular phenomena of inflammation. These proper phenomena may be induced without the previous occurrence of the accidental ones.¹

In studying the retardation of the circulation in the dilated vessels of the mesentery, it will be found that this sometimes commences somewhat suddenly, and that it is usually first observable in the veins. It gradually increases, until, ultimately, in some of the capillaries the

¹ "Neue Untersuchungen über die Entzündung." Cohnheim, 1873.

blood-stream completely stagnates. This constitutes the condition known as *inflammatory stasis*.

As the circulation becomes slower, the white blood-corpuscles (leucocytes) accumulate in the veins. Their natural tendency to adhere to the sides of the vessel is increased, so that they may nearly fill the tube. At the same time they exhibit active movement, by means of which they penetrate the walls of the vessels and pass into the surrounding tissues. Having escaped, they continue to exhibit active movements, and they may probably multiply by division. The red corpuscles also accumulate, more especially in the capillaries, and they likewise escape through the walls of these vessels, although in smaller numbers than the white. The mesentery thus becomes so filled with escaped corpuscles that but little else is to be distinguished. These phenomena will be described under the head of "Exudation of Liquor Sanguinis and Migration of Blood-Corpuscles."

Stasis.—The phenomena of stasis can be better studied in a more localized inflammatory process, such as may be produced by the application of some injurious agent—a small piece of nitrate of silver, for example—to the tongue of the frog. The stagnation of the blood which ensues is described by Cohnheim in his more recent researches, already alluded to, as being most marked in those capillaries which are situated in the more central portions of the inflamed area—here there is absolute stasis and no emigration of corpuscles takes place. Outside this there is an area in which the blood is circulating very slowly, and here both red and white corpuscles pass out of the capillaries; whilst more externally still, the white corpuscles only escape from the veins. The central area, which usually dies, is thus surrounded by an enormous number of red and white corpuscles; and the red corpuscles which have accumulated in this area are so closely packed that their outlines can scarcely be distinguished.

II. EXUDATION OF LIQUOR SANGUINIS AND MIGRATION OF BLOOD-CORPUSCLES.—Another constituent of the inflammatory process consists in the exudation of the liquor sanguinis and the migration of the blood-corpuscles.

a. *Migration of Blood-Corpuscles*.—The migration of the white blood-corpuscles (leucocytes) through the walls of the bloodvessels was first described, although very incompletely, by Dr. W. Addison in 1842.¹ This observer stated as the result of his researches, that

¹ "Experimental and Practical Researches on Inflammation." Trans. Prov. Med. Association, 1842.

in inflammation these corpuscles adhered to the walls of the vessels and passed through them into the surrounding tissues. In 1846 Dr. Augustus Waller described more fully the same phenomenon, and from his description there can be little doubt that he actually observed the emigration of the corpuscles.¹ Both these observers concluded that the escaped blood-corpuscles became pus-corpuscles. Their observations, however, were but little thought of and were soon forgotten, and it was not until 1867, when similar investigations were instituted quite independently by Professor Cohnheim, of Berlin—to whose minute researches we must ascribe most of our present knowledge on this subject—that the emigration of blood-corpuscles came to occupy an important place in the pathology of inflammation.²

The emigration may be observed in the mesentery of a frog which has previously been paralyzed by the subcutaneous injection of curare. The changes in the bloodvessels and in the circulation, and the accumulation of blood-corpuscles in the part, have been already described; it remains only to consider the phenomena of emigration.

The white blood-corpuscles (leucocytes) which have accumulated in large numbers, especially in the veins, remain almost stationary against the walls of the vessel, the blood-current passing by them, although with much diminished velocity. Those immediately adjacent to the wall, gradually sink into it, and pass through it into the surrounding tissue. In doing so they may be observed in the various stages of their passage. At first small button-shaped elevations are seen springing from the outer wall of the vessel. These gradually increase until they assume the form of pear-shaped bodies, which still adhere by their small ends to the vascular wall. Ultimately the small pedicle of protoplasm by which they are attached gives way and the passage is complete, the corpuscle remaining free outside the vessel.

The red corpuscles at the same time pass out of the vessels, but in less considerable numbers, and their transit is mainly through the walls of the capillaries, in which they have more especially accumulated. As has already been stated, in those portions of the inflamed tissue where absolute stasis has occurred, no emigration takes place.

β. *Exudation of Liquor Sanguinis*.—Associated with the passage of the blood-corpuscles through the walls of the vessels, is an exudation of the liquor sanguinis. The exuded liquor sanguinis, which

¹ Phil. Magazine, vol. xxix., 1846.

² "Ueber Entzündung und Eiterung," Virchow's "Archiv," vol. xl.

constitutes the well-known *inflammatory effusion*, differs from the liquid which transudes as the result of simple mechanical congestion, inasmuch as it usually contains a larger proportion of albumen and fibrinogenous substance, a proportion which increases with the intensity of the inflammation. It also contains an excess of phosphates and carbonates.

The most characteristic feature of inflammatory effusion is the large number of cell-structures which it contains. These are the direct product of the inflamed tissue, and are in no case generated spontaneously in the effused liquid. Most of them are migrated blood-corpuscles, others are derived from the proliferating elements of the tissue. The quantity and nature of the effusion will thus vary with the tissue inflamed, and with the severity of the inflammatory process. In non-vascular tissues, as cartilage and the cornea, exudation can only occur to a small extent from the neighboring vessels, and hence the effusion is small in quantity. In dense organs, as the liver and kidney, owing to the compactness of the structure, a large amount of effusion is impossible, and what there is, is so intermingled with the structural elements of the organ that it does not appear as an independent material. In the kidney it escapes into the urinary tubes, and so appears in the urine. The effusion is most abundant, and constitutes an important *visible* constituent of the inflammatory process, in inflammation of those organs which possess a lax structure and in which the vessels are but little supported, as the lungs, and in tissues which present a free surface, as mucous and serous membranes.

III. ALTERATIONS IN THE NUTRITION OF THE INFLAMED TISSUE.—The remaining constituent of the inflammatory process consists in alterations in the nutrition of the elements of the inflamed tissue.

The question as to how far the cellular elements of the tissue participate in the process of inflammation is one which even in the present state of our knowledge, owing to the difficulties which beset the histological examination of inflamed structures, admits only of an incomplete answer. The subject has, for the most part, been investigated in the lower animals, in which inflammation has been artificially induced. In man, the study of the primary changes is difficult, owing to the fact that the process can rarely be observed in its earlier stages. These changes will be more fully described when considering inflammations of particular organs and tissues; it will be sufficient in the present place merely to indicate their general characters.

The alterations in nutrition which accompany inflammation are in

certain tissues characterized by an exaltation of the nutritive functions of some of the cellular elements involved in the inflammatory process. This is evidenced by an increase in the activity of those elements which normally exhibit active movements, as the amoeboid cells of connective tissue and of the cornea. Cells, which under normal circumstances undergo no alterations in form, and exhibit no active movements, become active, sending out processes, and undergoing various alterations in shape. (Fig. 61.) This increase

FIG. 61.



Amoeboid Leucocytes. (v.
Recklinghausen.)

in the activity, and variation in the form of the cells, is usually accompanied by the growth of their protoplasm, and frequently by its division, or by vacuolation and endogenous development, and thus by the formation of new cells. In many cases the protoplasm, as it increases in bulk, becomes cloudy and granular, so much so as frequently to completely obscure any nuclei which it may contain. This occurs especially in epithelial elements, and it constitutes the condition known as "*cloudy swelling*." It is well seen in the glandular epithelium of the kidney in acute tubal nephritis. (See Figs. 104 and 105.)

A few years ago the cells of nearly all tissues were believed to exhibit these active changes in inflammation, and many of the young elements which abound in inflamed parts were regarded as the products of their proliferation. As our methods of histological investigation improve, however, and especially since the introduction of the chloride of gold process by Cohnheim, it has become increasingly obvious that the part which is played by the cells of the tissue in inflammation is much less than was formerly supposed; and that in many cases the young elements which infiltrate the inflamed structure are solely escaped leucocytes.

The physiological peculiarities of the cellular elements appear to influence very considerably their liability to undergo these active changes in inflammation. The cells in which active changes undoubtedly occur are those which are *normally* active, and in which growth and proliferation are associated with the maintenance of the tissue of which they are constituents. Such are epithelial elements. The activity of these is increased in the process of inflammation, and it is in inflammation of the skin, of mucous membranes, and of glandular structures, that cellular activity and proliferation are so constantly met with. The same is also true to a less extent of endothelium, as is exemplified in inflammations of serous membranes. In those cells, on the other

hand, in which normally no active changes take place, as the fixed cells of connective tissue and of the cornea, and probably also those of cartilage, it is doubtful if any activity is manifested in inflammation. The age of the cells probably also influences their tendency to become active, the younger being less stable and more prone to proliferate than the older elements.

Although the earlier alterations in the nutrition of the cellular elements of inflamed tissues are thus in many cases those of increased activity, the subsequent ones are characterized by *impairment* of nutrition. The well-known effect of inflammation is to injure the part affected by it. This injurious influence is in great measure due to the blood-stasis, and to the infiltration of the tissue with the inflammatory products which have escaped from the bloodvessels. (See "Suppuration.") If the inflammatory process be of considerable severity and stasis be induced in a wide area of the tissue, the nutrition may become completely arrested, and gangrene be the ultimate result. (See "Causes of Gangrene.") In inflammations of less intensity, the destruction of the tissue is less marked, the young cells which infiltrate it may undergo fatty degeneration and become absorbed, and the part thus recover more or less completely its former condition. (See "Inflammation of Common Connective Tissue—Resolution.") In other cases, especially in the less severe forms of inflammation, many of the young cells, both the emigrants and the offspring of the tissue proliferation, undergo progressive changes and lead to the development of a permanent tissue, which is, however, for the most part, inferior in its organization to the parent structure. (See "Inflammation of Common Connective Tissue—Organization.") This tendency of the new elements in inflammation to undergo progressive development will vary with the tissue involved, and with the intensity of the inflammation. The more intense the inflammation, the more abortive are the young cells, and the less is their tendency to form a permanent tissue. (See "Acute and Chronic Inflammations.")

In connective tissues, these changes in the cells are necessarily accompanied by changes in the intercellular substance. The latter are for the most part characterized by softening. In common connective tissue the fibres in the first place become succulent and less distinct, and ultimately they are completely destroyed; in cartilage the matrix softens and liquefies; in bone, the lime-salts are removed, the lamellæ disappear, and the osseous structure becomes converted into medullary tissue. Hence the destructive effects of the inflam-

Having thus briefly described the succession of changes which occurs in the process of inflammation, it remains to consider in what way these result from the injury of the tissue, and how far a causal relation subsists between them.¹

The first apparent change which follows the injury of the tissue consists in the dilatation of the bloodvessels, and in an *acceleration* of the flow of blood. Respecting the cause of this primary vascular phenomenon—the physiological investigations of Ludwig and Lovén show that a similar dilatation of the vessels and increase in the activity of the circulation is produced by the excitation of a sensory nerve in those parts in which the nerve originates. It was stated in a previous edition of this work that the primary vascular phenomena in inflammation were probably in the same way owing to an injurious impression received by the sensory nerves being reflected by the vaso-motor centre to the vessels. The more recent investigations of Cohnheim, however, which have been already quoted, render it necessary that this statement should be modified. Cohnheim produced precisely the same phenomena in the tongue of the frog after everything had been cut through with the exception of the lingual artery and vein; and also, in another case, after complete destruction of the brain, medulla, and spinal cord, as those which occur in the intact animal; and he considers that the vascular dilatation and accelerated blood-flow are due to the *direct* influence of the injurious agent upon the *walls of the bloodvessels*. These experiments therefore appear to prove conclusively that the initial injury of the bloodvessels is not *necessarily* reflex, although they do not exclude the probability that it may be so under certain circumstances.

With regard to the cause of the *retardation* of the blood-stream which so quickly succeeds its acceleration, and which may ultimately terminate in complete stasis—this also is due to some alteration in the properties of the walls of the bloodvessels through which the blood passes. That the retardation and ultimate stagnation of the blood-stream in acute inflammation is due to alterations in the walls of the bloodvessels and not to changes in the blood itself, was first shown by Mr. Lister.² Mr. Lister proved experimentally that the blood removed

¹ The following conclusions are in the main those arrived at by Professors Stricker and Burdon Sanderson.—Holmes's System of Surgery, vol. v. See also Lectures on the "Pathology of the Process of Inflammation," by Professor Sanderson.—Lancet, vol. i. 1876.

² "On the Early Stages of Inflammation."—Philosoph. Trans., 1858.

from an inflamed part did not differ in the tendency of the red blood-corpuscles to cohere to one another from normal blood; and he concluded, that the accumulation of the red corpuscles in inflamed tissues, and their adhesion to one another and to the walls of the vessels, was owing to their natural tendency to cohere together when in *abnormal* circumstances, such as occurs after their removal from the body; and that this cohesion in inflammation which leads to stasis was due to impaired vitality of the walls of the bloodvessels. This conclusion of Mr. Lister has been verified by more recent investigations. Ryneck has shown that stasis may be produced in the web of a frog, in which milk or defibrinated blood has been injected in place of the normal blood; and also that in vessels, the vitality of which has been destroyed by the injection of poisonous metallic substances, no stasis can be produced.¹ The investigations of Prof. Cohnheim are still more conclusive. Cohnheim ligatured the ear of a rabbit at its base with the exception of the median artery and vein, and emptied the vessels of their blood by the injection of a weak saline solution. He then injected into the emptied vessels various irritating solutions, and on again allowing the blood to flow through the thus injured vessels all the phenomena of inflammation ensued.² He also conducted a series of experiments, the result of which was to show that if the circulation in any organ (as the ear of the rabbit) be completely arrested for a sufficient length of time by the simultaneous occlusion of the arteries and veins, and then the blood be again allowed to circulate, dilatation of the vessels, hyperæmia, stagnation, exudation, and all the phenomena of inflammation take place in the part, these phenomena being obviously due to the damage sustained by the walls of the bloodvessels owing to their privation, for a sufficient length of time, of arterial blood. (See "Hemorrhagic Infarction.") The results of these experiments appear to be conclusive, and show that the retardation and ultimate stagnation of the blood-stream in inflammation, are owing to some impairment of the vital properties of the walls of the bloodvessels with which the circulating blood comes in contact.

In explanation of the phenomena of emigration, which takes place coincidently with the retardation of the blood-stream, the passage of the white corpuscles was formerly supposed to be due to their inherent

¹ Ryneck, "Zur Kenntniss der Stase des Blutes in den Gefassen Entzündeter Theile."—*Rolle's Untersuch. aus dem Institute für Phys. u. Histol. in Graz.*

² "Neue Untersuchungen über die Entzündung." Cohnheim, 1873.


activity, by virtue of which they penetrated the vascular walls, passing through stomata which were believed to exist between the endothelium. The red corpuscles were supposed to be forced through these openings by the increased blood-pressure. That neither of these explanations is sufficient to account for the phenomena is shown by Cohnheim's recent investigations. These appear to prove conclusively that the emigration of the blood-corpuscles and the exudation of the liquor sanguinis are, like the other phenomena of inflammation, due to some *impairment of the vital properties of the walls of the blood-vessels*. Hence, as so well stated by Dr. Burdon Sanderson, inflammation is simply the aggregate of those results which manifest themselves in an injured part, as the immediate consequence of the injury to which it has been exposed, an injury, it must be borne in mind, which, if of sufficient severity, would have led to its death.

The remaining constituent of the inflammatory process, the alteration in the nutrition of the inflamed tissue, succeeds the changes in the circulation and the exudation. Respecting the cause of the increased nutritive activity of the cellular elements which characterizes inflammation in certain tissues, it is probable that this is, for the most part, the result of the stimulation of the cells by the liquor sanguinis exuded from the bloodvessels; although the well-known experiment of Professor Stricker on the transplanted cornea of the frog, quoted in the last edition of this work, cannot now be accepted as conclusive. The impairment of nutrition produced by inflammation is due partly to the stagnation of the blood, and partly to the injurious influence of the products which escape from the bloodvessels. (See "Suppuration.")

SUPPURATION.—Suppuration, and the formation of abscesses, is a very frequent result of the inflammatory process; it occurs much more frequently, however, in some inflammations than in others. As a rule it may be stated that the more intense the inflammation the more abundant is the formation of pus.

The essential constituents of pus are cells and a liquid in which they are suspended. The liquid has an alkaline reaction and closely resembles the liquor sanguinis. It contains various kinds of albumen, fatty matters, and inorganic substances. The cells, or *pus-corpuscles* (leucocytes), are indistinguishable from the white corpuscles of the blood. As seen after death, they are spherical, spheroidal, or irregu-

lar-shaped, semi-transparent bodies, from $\frac{1}{2500}$ to $\frac{1}{1000}$ of an inch in diameter, containing a varying number of granules, and usually one or more distinct nuclei. (Fig. 62.) The addition of dilute acetic acid causes the cells to swell up; they become more spherical and transparent, and the nuclei are rendered more apparent. The size of the corpuscles and nuclei, and the number of the granules, present manifold variations. Pus-corpuscles, like white blood-corpuscles, lymph-corpuscles, and many other young cell-forms, all of which are included under the common term of *leucocytes*, are masses of contractile protoplasm. They possess the power of spontaneous movement, and when living undergo continuous alterations in form, and migrate in the tissues. (See Fig. 61.) They may also multiply.



The mode of origin of pus has been the subject of much controversy. The liquid ingredient proceeds directly or indirectly from the blood, it is the exuded liquor sanguinis: about this there is no dispute. The difference of opinion which exists is respecting the origin of the formed elements. Without discussing the theories which have been advanced by different pathologists, it must now be admitted that there are at least *two* sources from which the cells of pus may be derived, one from the *blood*, and the other from the *inflamed tissues*.

It has been seen that in the process of inflammation innumerable white blood-corpuscles pass out of the vessels into the surrounding tissues, and as these are indistinguishable from pus-corpuscles, it must be conceded that one mode of origin of pus is from the blood. Further, the white blood-corpuscles may multiply, and it is probable that by this means the production of pus may be greatly increased.

The other source from which the cells of pus are derived is from the cellular elements of the inflamed tissue. In certain tissues, as has been seen, especially in epithelial and endothelial tissues, the cells are the seat of active changes in inflammation; they may multiply and form new cells, and the more intense the inflammation, the more lowly organized are the newly formed elements, and the less is their tendency to form a permanent tissue. Some of these newly formed cells constitute pus-corpuscles. These, in this case, must be regarded as young elements resulting from the proliferation of the tissue, which are of low vitality, and soon perish.

Although the formed elements of pus may thus be derived both

from the blood and from the inflamed tissue, there can be no doubt that the former is their principal source, and that they are in the main migrated blood-corpuscles. In the earlier stages of the inflammatory process, they are mostly, if not all, emigrants; but in the later stages it must be admitted that they may also, in certain tissues, be derived from the cells of the inflamed part.

Such being the modes of origin of pus, it is evident that the more abundant the escape of blood-corpuscles, and the more active the proliferation of the elements of the inflamed tissue, the greater will be the formation of pus, and hence the greater its tendency to collect so as to form an abscess. It is consequently in those inflammations which are the most *concentrated* and the most *intense*—provided that the injury be not sufficiently severe to cause complete stasis—that the formation of pus is most abundant. The greater the injury sustained by the walls of the bloodvessels, the more readily will the blood-corpuscles penetrate them, and hence the more abundant will be the formation of pus. In inflammations of less intensity the escape of blood-corpuscles is less abundant, and the proliferation of the tissue less active, so that pus is not produced in sufficient quantities to cause its collection in the form of an abscess; it merely infiltrates the part, and may require for its recognition the use of the microscope.

Pus exercises a most injurious influence upon the surrounding tissues. The pus-corpuscles appear to be endowed with the power of absorbing the tissues with which they come in contact, or, at all events, of causing their liquefaction. Hence the softening and disintegration of the tissues which constitute such a destructive element in intense inflammations.

Pus which has remained for any length of time in the tissues undergoes certain changes. Its elements may undergo fatty metamorphosis, and thus be rendered capable of absorption. If pus is long confined in a closed cavity, its liquid portions may become absorbed, and its cells atrophy, so that it gradually dries up into a caseous mass, which may subsequently become calcified.

VARIETIES OF INFLAMMATION.—Inflammation exhibits certain variations in its characters according to the *severity* of the injury which produces it. The more severe the injury upon which the inflammatory process depends, the greater, *cæteris paribus*, will be the damage sustained by the bloodvessels, and consequently—if instantaneous stasis be not produced—the greater will be the tendency to the exudation of liquor sanguinis and to the emigration of leucocytes. Hence it is

in inflammations of considerable severity that the vascular phenomena are often so pronounced, the formation of pus so abundant, and the softening and disintegration of the tissues so considerable. Such inflammations, inasmuch as the action of the injury which produces them is for the most part of short duration, are sometimes designated *acute inflammations*.

In inflammations of less intensity, in which the damage sustained by the bloodvessels is less severe, the textural changes occupy a more prominent place, whereas the vascular phenomena are usually much less marked, and the formation of pus is less abundant. The precise nature of these textural changes must obviously partly depend upon the characters of the tissue which is the seat of the inflammation, but variations are also produced according to the severity of the injurious agent. It appears to me to be probably true—that the less severe the injury which produces the inflammation, the more do the resulting textural changes tend to be limited to the *connective tissue which is immediately adjacent to the bloodvessels and lymphatics*; whereas in inflammations of somewhat greater intensity more distant elements become involved. This is seen, for example, in inflammatory processes in the kidneys and in mucous membranes. In the former, the least severe forms of inflammation are characterized anatomically by an increase in the connective tissue around the bloodvessels (see “Interstitial Nephritis”); whilst in inflammations of somewhat greater intensity, the prominent textural change consists in swelling or proliferation of the epithelium within the tubules (see “Tubal Nephritis”). In mucous membranes also, the more severe inflammations are attended by epithelial proliferation, the less intense and more chronic by changes in the submucous connective tissue. With regard to the cellular elements in which these active changes take place—they are probably emigrants, and the cells which are the most intimately connected with the lymphatic system—viz., the cells of the connective tissue, and the endothelium of the lymphatics and of the bloodvessels. The tendency of the textural changes resulting from the least severe forms of inflammation to be limited to the connective tissue immediately adjacent to the bloodvessels and lymphatics, gives to these inflammations certain peculiarities. The new tissue originating around the vessels—consisting in the earlier stages mainly of small round cells, but ultimately becoming developed into a more or less completely fibrillated structure (an adenoid or fibrous tissue)—leads to an *induration* of the organ in which it is situated, and very constantly,

to the subsequent atrophy and retrogressive metamorphosis of its other histological elements. These changes will be more fully considered when treating of inflammation of the individual organs and tissues. (See "Inflammation of Common Connective Tissue," "Cirrhosis of the Liver," etc.) These least severe forms of inflammation, inasmuch as the injury which produces them is not only of slight severity, but is usually also prolonged in the duration of its action, are often known as *chronic inflammations*.

Inflammations are also divisible into *infective* and *non-infective*. The infective inflammations are those the products of which possess infective properties, owing to which the substances which are absorbed from the inflamed part tend to cause secondary inflammations of the tissues with which they come into contact. (See "Acute Tuberculosis" and "Pyæmia.") In these inflammations, Dr. Sanderson states, the tissues surrounding the seat of the injury are involved and infiltrated with the inflammatory products. The non-infective inflammations, on the other hand, are characterized by the absence of these infective properties.

Inflammations have received different names, according to the *nature* of the injury upon which they depend. Those inflammations which result from external injuries, mechanical or chemical violence, are called *traumatic*. Inflammations in which the nature of the injury is not obvious, are usually called *idiopathic*. The nature of the injury may give to the inflammatory process certain peculiarities. The contagium of smallpox, for example, gives rise to inflammation of the skin, constituting the "rash;" that of syphilis, to certain inflammations of the skin, mucous membranes, and other tissues; and that of typhoid fever, to inflammation of the intestinal lymphatic structures. In all these and numerous similar cases, the nature of the injury impresses upon the inflammation certain peculiarities, and in so far as the former is specific, the latter may be called *specific* inflammations. Lastly, inflammatory processes may be modified by the existence of certain *constitutional* peculiarities. This is the case especially in Scrofula.

Of these different varieties of inflammation, scrofulous inflammation, the infective inflammations, including pyæmia and acute tuberculosis, and the specific inflammations resulting from the poison of syphilis, will be separately considered in the next following chapters. The pathology of inflammation will then be still further elucidated by the study of the process as it occurs in the various tissues and organs.

CHAPTER XXVI.

23

SCROFULOUS INFLAMMATION.

THE process of inflammation when occurring in the Scrofulous usually presents certain peculiarities.


The constitutional condition known as Scrofula—a condition usually inherited but often acquired—is characterized by certain pathological tendencies. Of these the most important is an abnormal susceptibility of certain tissues to injury, and a peculiarity in the products and in the course of the inflammation which the injury induces. This susceptibility, more or less general, is commonly most marked in the mucous membranes and in the lymphatic glands, although the skin, bones, and joints are frequently affected. The part, however, which is the most prone to suffer varies considerably in different cases.

Not only is there this susceptibility to inflammation, but the inflammatory process tends to be exceedingly protracted; it is readily reinduced, and the alterations produced in the part differ from those caused by inflammation in healthy persons. When inflammation occurs in a healthy individual, if it does not cause the death of the part, the inflammatory products either become absorbed, or the process leads to suppuration, or to the formation of a vascular fibro-nucleated tissue. In scrofulous inflammation the absorption of the inflammatory products is very much less readily effected; they tend to *infiltrate* and *accumulate* in the tissue, where by their pressure they interfere with the circulation, and so lead to retrogressive and *caseous* changes. There is but little or no tendency to the development of new blood-vessels, and hence there is no organization of the new growth.

These peculiarities of inflammation in scrofulous subjects are to be in great measure ascribed to that inherent low vitality of the tissues which obtains in this disease, and also to certain peculiarities in the histology of the inflammatory products. Virchow long ago pointed out the richly cellular character of the products of scrofulous inflammation, the tendency of the cells to infiltrate the tissue, and the extreme tardiness with which the infiltration becomes absorbed. Quite recently Professor Rindfleisch has stated that these cells are, for the

most part, *larger* than those met with in healthy inflammations; and that, this being the case, their removal by passage into the lymphatics is less readily effected.¹ In tubercle, the close relation of which to scrofulous lesions is well known, the existence of large cell-forms is almost constant, and their pathological significance in both these allied products will be more fully considered when speaking of tuberculosis. (See "Tubercle and Acute Tuberculosis.") This largeness of many of the young cells in scrofulous inflammation, and their marked tendency to infiltrate and accumulate is well shown in the accompanying drawing (Fig. 63).

FIG. 63.



Scrofulous Inflammation of a Bronchus.—Section of a small bronchus of a markedly scrofulous child, the subject of bronchitis, which terminated in miliary tuberculosis. The deeper structures of the bronchial wall are seen to be extensively infiltrated with cells, most of which are *larger* than those met with in the less extensive infiltration of healthy inflammation. The infiltration extends to and invades the walls of the adjacent alveoli, which are seen at the upper part of the drawing. The cavity of the bronchus contains a little mucus, *m.* $\times 200$, reduced $\frac{1}{4}$.

These histological peculiarities of the products of scrofulous inflammations not only lead to an extensive and obstinate infiltration of the affected tissues, but, as insisted upon by Rindfleisch, they must in the parenchyma of organs, as in the glands and viscera, cause, by the

¹ "Ziemssen's Cyclopædia of Practical Medicine," vol. v. Article, Chronic and Acute Tuberculosis, by Rindfleisch.

pressure they exercise, more or less obstruction of the bloodvessels, and so interfere with the vascular supply. To this interference with the vascular supply, and to the inherent low vitality of the cellular elements, is to be mainly ascribed the retrograde changes and *caseous metamorphosis* which are so characteristic of scrofulous lesions.

CHAPTER XXVII.

TUBERCLE AND ACUTE TUBERCULOSIS.

By acute tuberculosis is understood a general infective disease, which is characterized anatomically by the occurrence of numerous minute nodular lesions more or less generally disseminated in the various organs and tissues. The generally disseminated nodular lesions, which are characteristic of the disease, appear to be inflammatory growths, resulting from the distribution of infective materials (probably minute particles), by means of the bloodvessels or lymphatics from some primary inflammatory product. They are, therefore, the anatomical results of an infective inflammatory process, and they constitute what have long been known as *miliary tubercles*.

General Pathology of Acute Tuberculosis.—Our knowledge of acute tuberculosis and of its anatomical result—*tubercle*, has until recently been involved in obscurity; but at the present time—owing in great measure to scientific experimental research—it may be regarded as being much more complete. According to the older doctrines, which were based upon the teaching of Laennec, tubercle was looked upon as a specific non-inflammatory growth which originated spontaneously in the tissues. Further, this new growth was characterized by the regular succession of changes which it invariably underwent;—it was first gray and translucent, then became opaque, and ultimately caseous. Hence in its earlier stages it was known as *gray*, in its later as *yellow* tubercle. Caseous metamorphosis was held to be such a distinguishing peculiarity of the growth, that all caseous masses were regarded as tubercular, and the term “tubercle” came to be applied indiscriminately to all pathological products which had

undergone this form of degeneration, and which in their color and consistence somewhat resemble soft cheese. Caseation, however, as already stated, although most frequent in tuberculous and scrofulous lesions, is a common result of the retrograde metamorphosis of many growths which are destitute of or contain but few bloodvessels, and which consist of closely-crowded cellular elements. (See "Caseation.") For this much wider extension of the pathological significance of caseous degeneration we are mainly indebted to Professor Virchow.¹

The infective nature of acute tuberculosis was first promulgated about twenty years ago by Buhl, who stated that in the majority of cases of this disease, indurated masses which had become caseous existed in some part of the body, and that to the absorption of substances from these infective centres the general development of the tubercle was owing. He further stated that in those cases in which the tubercles were confined even to limited portions of a single organ, they were also secondary to caseous lesions. Buhl's theory, therefore, implied that the origin of the infective substances was necessarily associated with caseous metamorphosis of the primary inflammatory induration. A modification of this view of the nature of the infecting lesion has since been rendered necessary, both by the results of post-mortem observations, and also by those which have been obtained from the artificial production of tuberculosis in the lower animals.

The experimental investigation of acute tuberculosis was commenced by Villemin in 1865, and subsequently followed out by Burdon Sanderson, Wilson Fox, Cohnheim, Klebs, and others. The methods of investigation consisted either in the inoculation of various inflammatory products—for the most part products of *chronic* inflammations (caseous or not caseous), or in the production of a local inflammatory induration by the introduction of setons and of other foreign bodies beneath the skin. Of inflammatory products Dr. Sanderson found that none proved so active as that obtained from the indurated lymphatic glands of an animal already suffering from the disease. In both cases, after a certain lapse of time, disseminated inflammatory lesions were produced in various organs and tissues—lesions which presented a special tendency to become caseous at their centres. It was also found that the distribution of these lesions varied according as the infective materials were introduced into the bloodvessels or

¹ For further information on the history of "tubercle," the reader is referred to the commencement of the chapter on "Pulmonary Phthisis."

Lymphatics, clearly proving that they resulted from the dissemination of infective substances by means of the blood and lymph streams. The results of these experiments therefore show, in the first place, that caseation of an inflammatory product is not necessary in order for it to constitute an infective focus; and, secondly, that the development of the general tuberculosis is not due to anything specific in the substances inoculated, but that the products of various inflammatory processes (for the most part of inflammations of slight intensity), may constitute the infective agents.

The lesions produced, however, in artificial tuberculosis differ somewhat from those met with in the natural disease as it occurs in man. They differ both in their anatomical distribution, and, to a less extent, in their pathological tendencies. The differences in anatomical distribution are principally confined to the brain and lungs. The brain, which is so frequently affected in the natural disease in man, is rarely so in artificial tuberculosis. In the lungs, the first structural changes which take place in artificial tuberculosis are stated by Dr. Klein to consist in the development of adenoid tissue around the perivascular lymphatics, the implication of the alveoli being only a secondary part of the process.¹ In the natural disease in man, on the other hand, the changes commence in the alveoli themselves. Respecting differences in their pathological tendencies—it is to be remarked that caseation occurs much more extensively in the lesions artificially induced than in man. In the former, also, diffused tracts of consolidation are more frequently associated with the miliary lesions, and they may even constitute the predominant structural changes. These differences, however, are probably to be regarded as resulting from differences in the intensity of the infective agent, and from peculiarities in the morbid tendencies of the tissues involved, and not as any evidence of a want of analogy in the pathology of the two diseases.

Passing on to consider acute tuberculosis as it is met with in man, it must in the first place be stated that it occurs most frequently in those who are *scrofulous*, and one or more masses of inflammatory induration which have become *caseous* are, in the great majority of cases, to be found in some part of the body after death; sometimes in the lungs, sometimes in the bronchial glands, sometimes in the glands of the mesentery, etc. Caseous metamorphosis, as was seen in the

¹ "On the Lymphatic System of the Lungs," by Dr. E. Klein. "Proceedings of the Royal Society," No. 149, 1874.

preceding chapter, is exceedingly common in scrofulous inflammation, owing to the marked cellular infiltration and consequent anæmia which characterize the process; and *it is these products of scrofulous inflammation which are the most common cause of acute tuberculosis*. Much less frequently cases of acute tuberculosis are met with, in which there exists some inflammatory lesion which is not caseous, as a simple induration, an inflamed bone, or an ulcerated mucous membrane; whilst in exceedingly rare cases it is stated that no such product of a previous inflammatory process has been discoverable.

The results of post-mortem observations of the natural disease thus appear to justify the same conclusions respecting the nature of the infective substances as those derived from the experimental investigation of the disease in the lower animals, and in a man also it must therefore be regarded as in the highest degree probable that any inflammatory product may, under certain circumstances, give rise to a tuberculous process, and that although caseation of the product is most frequent, it is not essential in order for it to constitute an infective focus. Why such inflammatory products should in some cases be infective, whilst in others they remain inert, we are unable certainly to explain. In attempting to answer this question, it must be remembered that such products may *accidentally* become placed in direct communication with the vascular or lymphatic systems; and also that infective substances are much more liable to produce results in some constitutions and in some conditions than in others. It is also possible that the infective properties of an inflammatory product may be determined by atmospheric influence, or by the presence of minute organisms. (See "Septicæmia.")

Whilst acute tuberculosis is thus an infective disease, it is an infective disease of a special kind. In the first place, the infective substances derived from the infective focus—which are probably minute particles—not only exercise their injurious influence over areas of tissue which are for the most part exceedingly small (hence the miliary character of the lesions), but the injury they inflict is of comparatively slight severity. The principal result of their dissemination is consequently to cause a textural change—a new growth of tissue at the seat of the injury. (See "Chronic Inflammations.") In this respect this disease presents a marked contrast to one with which it is closely allied—pyæmia, in which the severity of the injury produced by the infective particles is much greater and gives rise to the formation of abscesses. (See "Pyæmia.") In pyæmia the in-

tensity of the disseminated inflammatory processes is considerable, and the course of the disease is usually acute, whilst in acute tuberculosis the inflammatory processes are much less severe, and the disease tends to run a more chronic course.

HISTOLOGY OF TUBERCLE.—The miliary lesions in acute tuberculosis, although presenting certain differences according to their age, and to the nature of the tissue in which they originate, are tolerably uniform in their histological characters. Their most marked feature is the prominent place which large multinucleated masses of protoplasm, the so-called giant cells, occupy in their constitution. These large cells, which somewhat resemble the myeloid cells met with in sarcomatous tumors, etc., were long ago alluded to by Virchow, Wagner, and others, but it is only during recent years, mainly owing to the researches of Oscar Schüppel and Langhans, that they have come to occupy a prominent place in the histology of tubercle.

The most characteristic features of these multinucleated cells are their large size, the number of their nuclei, and the irregularity of their outline. Some of the larger ones measure as much as $\frac{1}{16}$ inch in diameter. They possess no limiting membrane, but are simple masses of protoplasm, containing numerous round or roundly-oval nuclei, each inclosing a bright nucleolus. (Figs. 64, 65, and 66.)

FIG. 64. *P. L. P.*

A Multinucleated Cell from the Lung in a case of Chronic Phthisis.—Showing the large number of nuclei with bright nucleoli. $\times 400$.

FIG. 65.




A Multinucleated Cell from the Lung in a case of Chronic Phthisis.—Showing the long branched processes, which are continuous with the reticulum of the surrounding indurated growth. Some of the processes are in connection with smaller nucleated elements. $\times 200$.

As many as forty nuclei may occasionally be counted in a single cell. Some of them are much smaller, and contain only three or four nuclei.

Four or five, or even more, of these multinucleated masses are sometimes found in a single tuberculous nodule. Many of these large cells possess long branched processes, in connection with which, and evidently originating from them, are smaller protoplasmic masses, also nucleated and branched. (See Figs. 65 and 66.) The meshes between the branched cells are, according to Schüppel, filled with epithelial-like elements. These elements I have failed to observe, and the meshes I have either found empty, or containing a few lymphoid cells. (Fig. 66.) It would thus appear that the original protoplasmic mass gives origin to a network of large branched cells.

FIG. 66.



Multinucleated and branched Cells from a firm Gray Military Tubercle of the Lung in a case of Acute Tuberculosis.—Wide meshes are seen in the immediate vicinity of the cells inclosing a few lymphoid elements. The branched processes are directly continuous with the adenoid reticulum of the tubercle. $\times 200$.

With regard to the histological elements from which the giant cells originate—this varies. In the lung, Dr. Klein states, that they are derived from the alveolar epithelium. They may also probably originate from the cells of connective tissue, and from the endothelium of the bloodvessels and lymphatics. Their formation takes place either by the fusion of two or more cells, or by the excessive development of one cell. In the latter case, the cell increases in size and its nuclei multiply, but here the process of development ceases—there is no subsequent division of the cell.

Associated with, and surrounding the giant cell and its branches, is a small-celled adenoid-like structure. (Fig. 66.) This small-celled structure, which often contributes largely to the formation of the tu-

bercle, somewhat resembles that of an indurated lymphatic gland, which is commonly known as adenoid tissue. A similar structure, as has been already stated, is also met with in chronic inflammations of the liver, lungs, and other organs. (See "Chronic Inflammations.") It consists, in the main, of lymphoid cells, which are either separated from one another by fine bands of homogeneous transparent looking material, or by a more or less distinctly fibrillated, and sometimes nucleated reticulum, within the meshes of which the cells are grouped. (Fig. 67.) This reticulum is sometimes dense and well marked, whilst in other cases it is much less prominent. In addition to the small lymphoid cells there are often seen some rather larger cells containing one, and in some cases, two nuclei. (See "Scrofulous Inflammation," Fig. 63.)

FIG. 67.



A portion of a Gray Miliary Tubercle of the Lung.—Showing the adenoid-like structure met with in large portions of those nodules. X 230.

An elementary tubercle thus consists of a giant-cell reticulum surrounded by, and in direct histological continuity with, a varying sized zone of small-celled tissue. This is well shown in Fig. 66. (See also Fig. 70.) The larger tuberculous nodules consist of several of these giant-cell systems. (See Fig. 76.)

Although the above characters of tubercle are to be observed in successful preparations and in certain stages of the tuberculous growth, many of them will often be found wanting. The nuclei and processes of the giant cells will not always be seen, and the cells often appear simply as yellowish, somewhat granular masses in which neither nuclei nor processes are visible. In this imperfect state they are to be found in tuberculous nodules from all organs in which tubercle is met with.

It remains to speak of the bloodvessels of tubercle. The vessels of the tissue in which the nodule originates gradually become obliterated in the process of its growth, and there is no new formation of vessels, such as takes place in more highly developed inflammatory tissue. (See "Inflammation of Common Connective Tissue.") The tubercle is therefore, except in the earliest stages of its development, non-vascular. (See "Scrofulous Inflammation.")

Although the structure which has been described is that most commonly met with, it must be borne in mind that all tuberculous lesions are not thus constituted. When treating of the changes in the several organs, it will be seen that the precise histological constitution of the

nodules varies somewhat, according to the characters of the tissue in which they originate. In the lungs, for example, many of them consist largely of accumulations of epithelial cells within the pulmonary alveoli.

Secondary Changes.—Tubercle invariably undergoes more or less retrograde metamorphosis, although the extent of this varies considerably, and in some cases the nodules may become developed into an imperfect fibroid structure. The occurrence of retrograde metamorphosis is mainly owing to the obliteration of the bloodvessels which accompanies the growth of the lesions. The change commences in the centre of the nodule, this being the part first developed, and consequently that which is the furthest removed from vascular supply. The nodule breaks down into a granular fatty débris, so that its central portions soon become opaque and yellowish. (Fig. 68.) In some cases the process of disintegration is rapid, whilst in others it is more

FIG. 68.



One of the Gray Nodules from the Lung in a case of Acute Tuberculosis, which is becoming opaque and soft in the centre. (Diagrammatic.)

gradual. It is usually most marked in the larger and more diffused lesions, and hence it is these lesions which are most commonly of a yellow color and soft consistence ("yellow tubercle"). In other cases the retrograde change is less marked, the reticulum of the nodule becomes denser and more fibroid, and although the imperfect fibroid tissue usually ultimately undergoes in its central parts, more or less fatty metamorphosis, the nodule may remain as a firm fibroid mass.¹ This occurs more especially in the smaller lesions. The extent and rapidity of the retrograde change depends, I believe, partly upon the intensity of the infective process, and partly upon constitutional conditions. The existence of scrofula favors retrograde changes in the tuberculous lesions as it does in all inflammatory products, and it is in those who are markedly scrofulous that tubercle undergoes the most rapid degeneration. (See "Scrofulous Inflammation.") These changes will also be influenced by the intensity of the infective process. The more intense the process, the greater is the tendency to the degeneration and softening of

¹ Dr. Klein states (*loc. cit.*) that the large multinucleated cells undergo a fibroid transformation, and become converted into a dense feltwork of fibrillar tissue, which tissue gradually dies away and becomes caseous.

the nodules; the less intense and more chronic, the more liable are the miliary lesions to become fibroid.

LOCAL TUBERCULOSIS.—Before concluding the consideration of the general pathology of tuberculosis, allusion must be made to those cases in which the tuberculous processes are confined to a single organ. This limited tuberculosis differs from the more general one only in the extent of the distribution of the infected materials. The miliary lesions originate from some retrograde inflammatory product situated usually in the same organ as that in which they occur, and their limitation is probably owing to the infective substances being disseminated by the lymphatics and serous canals and not by the bloodvessels. This will be again referred to in the chapter on “Pulmonary Phthisis,” and also in the following description of the tuberculous process as it occurs in the several organs and tissues.

CONCLUSIONS.—Having thus described the histological characters of tuberculous lesions, it remains to consider the relation which subsists between these lesions and the products of non-tuberculous inflammations. The tuberculous lesions being inflammatory growths—the result of the injurious influence of infective particles upon the small areas of tissue with which they come into contact—why do they differ from the tissue changes which accompany ordinary chronic inflammation? The answer to this question is at present far from satisfactory.

It must be admitted that constitutional conditions influence considerably the occurrence and character of tuberculosis. Tuberculous processes, for the most part, in the scrofulous, and the products of some scrofulous inflammation are the most frequent infective agents. Constitutional conditions may probably thus so modify an inflammatory process as to give to its products infective properties. It must, I think, also be regarded as probable that conditions of the constitution may influence the effect produced by the dissemination of the infective particles, and that these particles, whether derived from a caseous or non-caseous focus, may in certain conditions of the constitution give rise to tuberculous processes, whilst in others their dissemination is unattended by change.

The histological peculiarities of the tuberculous lesions appear also to depend mainly upon constitutional conditions. The influence of scrofula upon the histological changes which accompany inflammation has already been seen, and from the close histological relationship which subsists between the disseminated nodules met with in tubercu-

losis and the more diffused lesions produced by primary inflammation in scrofulous subjects, and also from the fact that it is in the scrofulous that tuberculosis most frequently occurs, we may, I think, assume that the characters of tuberculous lesions also are dependent upon a constitutional state.

In both scrofulous inflammation and tuberculosis there is a tendency to the development of large cell-forms and to the formation of a lowly organized and non-vascularized structure which soon undergoes retrograde changes. These characters of the inflammatory new growth appear to me to be probably due to the inflammatory process occurring in tissues of such low vitality that the cellular inflammatory products are incapable of forming an organized vascular tissue, but merely undergo some increase in size and then tend to slowly degenerate. The protoplasm grows, the nuclei multiply, but the higher manifestation of vitality—the subsequent division of the protoplasmic mass—does not take place, and thus are produced the giant cells.

Tuberculosis of the Pia Mater.


In the pia mater the tuberculous process is associated with inflammation of the meninges, constituting the condition known as *tubercular meningitis*. This is almost invariably a part of a general tuberculosis.

The process is almost exclusively confined to the pia mater as the base of the brain, and the tuberculous nodules—which may easily escape observation—are seen in connection with the small arteries in the Sylvian fissures, and deeply seated between the convolutions. A few scattered granulations are, however, frequently visible on the upper surface of the hemispheres. The inflammatory growth originates in the perivascular lymphatic sheaths which inclose the small arteries of the pia mater. (Fig. 69.) The cells of the sheath multiply, and the process of proliferation commencing at separate centres, numerous small gray nodules are produced around the vessels. These, which are distinctly visible to the naked eye, cause an external bulging of the sheath, and a diminution in the calibre, or even complete obliteration, of the inclosed.

The localized obstructions to the circulation which result from the pressure of the perivascular nodules cause intense hyperæmia of the collateral vessels, and thus the pia mater at the base of the brain becomes exceedingly vascular, there being in some cases rupture of

the vessels and extravasation. This is followed by a more general inflammatory process, a true basic meningitis. A transudation of the liquor sanguinis takes place from the hyperæmic and injured vessels, blood-corpuscles escape, and thus the meshes of the pia mater become infiltrated with a sero-fibrinous liquid, which in many parts has a puriform character.

FIG. 69.



Miliary Tubercle in the Pia Mater.—The dotted line indicates the original size of the tubercular nodule. A. The lymphatic sheath. V. The bloodvessel. F. Proliferation of elements within the sheath. $\times 100$. (Cornil and Ranvier.)

The changes in the pia mater at the base of the brain are attended by softening of the immediately subjacent cerebral substance, which becomes infiltrated with young cells. The lateral ventricles at the same time become distended with serosity (acute hydrocephalus), so that the convolutions on the surface of the hemispheres are seen to be much flattened. The ependyma and choroid plexus also become exceedingly vascular, and the walls of the ventricles, together with the fornix and soft commissure, become much softened. All of these changes are owing, partly to an inflammatory process, and partly to the mechanical obstruction to the circulation caused by the tuberculous growth. In addition, the arachnoid membrane is dry and sticky.

TUBERCULOUS MASSES IN THE BRAIN.—In addition to the miliary lesions occurring in the pia mater in tubercular meningitis, large tuberculous masses are occasionally met with in the brain unassociated with a general tuberculous process. These masses, which vary in size from a hazel nut to a hen's egg, commonly occur in the cerebral sub-

stance, especially at the base of the brain. They are of a pale yellow color and firm consistence, and usually form quite round globular tumors. Their surface is often seen to be covered with minute gray nodules, which extend into the surrounding tissue; and on section, similar nodules are sometimes visible, scattered through the substance of the tumor. In most cases only one or two such masses are found, but occasionally they are more numerous. They occur especially in childhood, and usually in children in whom there is a general tendency to caseation of inflammatory products ("scrofulous" children). When examined microscopically they are found to be made up of an adenoid structure and large branched cells, such as has been already described as so characteristic of tuberculous lesions. This structure, which is best seen in the peripheral portions of the tumor, is often concentrically arranged around bloodvessels, and is found undergoing in different parts fibrous and caseous metamorphosis.


Respecting the nature of these masses and the way in which they originate, it is difficult to speak with certainty. From the fact that miliary nodules are so often to be seen on their surface and extending into the surrounding tissue, it is supposed that they originate by the aggregation of such nodules, that the primary nodule constitutes an infective focus, and so causes a succession of growths in its immediate vicinity. This hypothesis is probably correct. Occasionally the tuberculous mass causes a more general infection, and so gives rise to tubercular meningitis or to a general tuberculosis.

Tuberculosis of Lymphatic Glands.

In the lymphatic glands, tuberculous processes give rise, in the first place, to changes in the peripheral portions of the gland, inasmuch as it is with these that the infective materials which are conveyed by the lymphatic vessels first come into contact.

The active changes in the gland begin with the formation of a large multinucleated branched cell. (Fig. 70.) In the earlier stage of the process small gray nodules are visible scattered through the cortex. These gradually increase in size and become caseous. The gland

FIG. 70.



Tuberculosis of a Lymphatic Gland.—The earliest stage of the process. Showing the giant cell. $\times 200$.

then becomes enlarged, the distinction between its medullary and cortical portions becomes lost, and it becomes changed to a grayish homogeneous mass, in which are varying sized tracts of caseous material. The new growth very frequently undergoes a marked fibroid development, so that the caseous masses are surrounded by a dense fibroid structure. The caseous portions of the gland may subsequently soften, dry up, or calcify.

Tuberculosis of Mucous Membranes.

In mucous membranes the development of tuberculous nodules is usually secondary to some primary inflammatory process which leads to ulceration of the membrane. The intestinal, the urino-genital, and the respiratory mucous membranes may all be the seats of a tuberculous growth.

The Intestine.—In the intestine the occurrence of true tuberculous processes appears to be almost invariably preceded by primary inflammatory changes in the intestinal lymphatic structures. These changes have their seat in the solitary and Peyer's glands, and as in typhoid fever, it is especially these structures in the lower part of the small intestine and in the cæcum which are affected.

The first stage of the process consists in an inflammatory hyperplasia of the lymphatic elements. In Peyer's patches this hyperplasia usually affects isolated follicles in the patch. The solitary glands and certain follicles in the patches thus become swollen, and project with undue prominence above the surface of the membrane. The newly formed elements then undergo retrogressive changes, they soften, the degeneration in the patches commencing at several separate centres, and often extending until the whole patch becomes destroyed. As the result of these primary inflammatory changes an ulcerated surface is produced, the floor and edges of which are more or less thickened, owing to the extension of the inflammatory infiltration into the sub-mucous connective tissue. This a simple ulcer, and its production, so far, is quite independent of a tuberculous process.

What must be regarded as a tuberculous change occurs subsequently to the primary ulceration, and it consists in the development of small nodules of new growth in the floor of the primary ulcer. The development of these nodules appears to take place principally around the bloodvessels, and as these are arranged transversely around the intes-

FIG. 71.



A Tubercular Ulcer of the Intestine.—(Diagrammatic.) a. Epithelial lining. b. Submucous tissue. c. Muscular coat. d. Peritoneum.

tine, the new growth proceeds in the same direction. These secondary nodules of new growth, like the primary inflammatory product soften and become caseous, and thus the process of ulceration gradually extends transversely until the whole circumference of the gut may be destroyed. The ulcer thus produced presents a strong contrast to that of typhoid. Its edges and base are thickened and indurated, and the tuberculous nodules, tending to become caseous, are seen scattered in its floor. (Fig. 71.)

The tubercular ulcer rarely, if ever, heals. Owing to the thickening of the tissues at its base, perforation is quite an exceptional occurrence. In the process of its extension the ulceration is attended by some contraction and narrowing of the gut.

Tuberculosis of the Lungs.

Tuberculous processes occur in the lungs as a part of a general tuberculosis, and also in many cases of pulmonary phthisis. The nature of the resulting inflammatory lesions is similar in both. It will be well, however, in the present place, more particularly to describe these lesions as they occur in the general infective disease. The more limited processes which take place in many cases of phthisis will be again referred to in a subsequent chapter devoted to the consideration of this affection. (See "Pulmonary Phthisis.")


The pulmonary lesions met with in general tuberculosis consist, for the most part, of disseminated nodular growths, which are universally known as miliary tubercles. These growths are of two kinds—the *gray* and the *yellow*. The *gray* are semitransparent nodules of a grayish-white color, varying in size from a small pin's head to a hempseed. They are somewhat spherical in shape, and usually possess a well-defined outline. Sometimes they are firm, and almost cartilaginous in consistence; whilst in other cases they are much softer and almost gelatinous. These softer forms, instead of being semi-transparent, are more opaque and white. The *yellow* are, for the most part, larger than the preceding, many of them much so, some being as large as a pea. They are also softer in consistence, less defined and regular in outline, and they pass more insensibly into the surrounding tissue. Many of them possess a grayish-white translucent margin,

which may be pretty firm in consistence, but never so hard as are many of the gray nodules, whilst their central portions are opaque, yellowish, or caseous.

Both the gray and the yellow nodules are often found associated in the same lung; in other cases the gray nodules only are met with; whilst, less frequently, nearly all the growths are of the yellow variety. The condition of the pulmonary tissue which is situated between the nodules varies considerably. It may be perfectly normal, more or less congested and oedematous, or it may present varying sized tracts of grayish, granular, friable consolidation. A perfectly normal condition of the intervening pulmonary tissue is found in many of those cases, in which all the growths are of the firm, gray variety; but when there are numerous yellow or soft gray nodules, the lungs are nearly always more or less congested or consolidated.

When these nodules are examined microscopically they are seen to exhibit two different kinds of structure—viz., the adenoid structure

FIG. 72.




A small soft Gray Tubercle from the Lung in a case of Acute Tuberculosis.—The whole of the tubercle is shown in the drawing, and it is obviously constituted largely of *intra-alveolar* products. $\times 100$, reduced to $\frac{1}{2}$.

with branched multinucleated cells, which has been already described as that which is the most characteristic of tuberculous lesions, and accumulations of epithelial cells within the pulmonary alveoli (catarrhal pneumonia). There is, however, this marked difference between the various kinds of nodules—that whereas the smaller firm gray ones

are constituted almost entirely of the first-named structure, the larger soft gray, and most of the yellow ones, consist largely of the intra-alveolar accumulations.

Firstly, with regard to the soft gray and yellow nodules. Most of these when examined with a low magnifying power present the appearance represented in Fig. 72, the nodules evidently consisting largely of

FIG. 73.



A portion of a small soft Gray Tubercle from the Lung.—This is from a case of acute tuberculosis, probably in an earlier stage than that from which Fig. 72 was drawn. The figure shows one of the alveoli filled with epithelial elements and a few small cells, with some cellular infiltration of the alveolar wall. $\times 200$.

accumulations within the alveolar cavities. When more highly magnified their constitution becomes more apparent. It is thus seen that the alveolar cavities are filled with epithelial elements and small cells resembling leucocytes, whilst the alveolar walls are more or less extensively infiltrated and thickened with lymphoid cells. (Fig. 73.) In many cases the central portions of the nodules will be seen to have undergone extensive degenerative changes, and to consist merely of a structureless granular debris, so that the accumulations


within the alveoli and the cellular infiltration of the alveolar walls are only visible at their periphery. This is always the case in the distinctly yellow tubercles. (Fig. 74.)

The histological characters of the firmer gray nodules differ somewhat from the preceding. In these the cellular infiltration and thickening of the alveolar wall is much more marked, and as many of the alveolar cavities are occupied by giant cells, these probably originating, as described by Dr. Klein, from the alveolar epithelium.¹ (Fig. 75.) In other cases, the alveolar structure has completely disappeared, and the tubercle, when examined with a low magnifying power, appears as a little somewhat spheroidal mass, the cellular elements of which are seen to be grouped around separate centres. (Fig. 76.) When more highly magnified these centres are seen to correspond with the

¹ These large multinucleated cells are stated by Dr. Klein (*loc. cit.*) to originate either by the fusion of the alveolar epithelium, or by the excessive development of one epithelial cell. Since the publication of Dr. Klein's statement I have frequently observed these cells situated distinctly in the alveolar cavities, and I have little doubt they originate in the way he describes.

large multinucleated branched cells already described, and the small-celled structure grouped around them, as well shown in Fig. 66.

FIG. 74.




A portion of a Yellow Tubercle from the Lung in a case of Acute Tuberculosis.—Showing the degeneration of the central portions of the nodule *c*, and the cellular thickening of the alveolar walls and accumulations within the alveolar cavities at the periphery *p*. $\times 200$.

This is a fully developed tubercle of the lung. The small-celled structure at the peripheral portions of the nodules extends into and produces a thickening of the walls of the alveoli with which the nodule is incorporated. (Fig. 77.) In the tubercles thus constituted, extensive retrogressive changes rarely occur. Degeneration is slow and very incomplete, and the nodule often becomes imperfectly fibroid.

Respecting the cause of these differences in the histological characters of the miliary lesions in the lungs—I believe them to depend upon differences in the age of the nodules, and in the intensity of the tuberculous process.¹ If the intensity of the process

FIG. 75.




A portion of the more external part of a Gray Tubercle from the Lung in a case of Acute Tuberculosis.—Showing the extensive infiltration and thickening of the alveolar walls, and the giant cells within the alveolar cavities. $\times 100$.

¹ Intensity comprises two factors—severity of injury and susceptibility of injured tissue. In scrofula, for example, the susceptibility of the tissue is great, and the intensity of tuberculous processes is often considerable.

be considerable, the nodules will consist in the main of accumulations of epithelium within the pulmonary alveoli, and the nodule will rapidly


FIG. 76.



A firm Gray Tubercle from the Lung in a case of Acute Tuberculosis.—Showing the grouping of the elements around separate centres, the nodule consisting of several giant-cell systems. $\times 33$.

undergo disintegration. (See Fig. 74.) If the process be less intense, the nodules attain a more advanced age, degeneration will be less and

FIG. 77.



A small portion of the most external part of a firm Gray Tubercle from the Lung in a case of Acute Tuberculosis.—Showing the incorporation of the nodule with the alveolar wall *a*. $\times 270$.

rapid and complete, the cellular infiltration and thickening of the alveolar walls will be greater, and the epithelial elements may

Form large multinucleated cells. (See Fig. 75.) Lastly, in the least severe and most chronic processes, the development of multinucleated elements and the formation of the network of branched cells reaches its maximum (see Figs. 66 and 76), degeneration takes place slowly in the central portions of the nodule, and there is often considerable fibroid induration of the new tissue. There is thus a close analogy between the tissue-changes resulting from tuberculosis of the lungs and those which result from other inflammatory processes. (See "Chronic Inflammations.") The degeneration of the lesions is mainly owing to the interference with their vascular supply, caused by the pressure exercised upon the bloodvessels by the intra-alveolar accumulations, and the obliteration of the vessels by the cellular infiltration of the alveolar walls.

CHAPTER XXVIII.

PYÆMIA AND SEPTICÆMIA.

THE diseases known as Pyæmia and Septicæmia result from the absorption and dissemination of certain infective substances, derived usually from the products of some acute inflammation. In these diseases, therefore, as in acute tuberculosis, we have a focus from which the infective substances are derived, the general dissemination of these substances, and the secondary processes to which they give rise.

Pyæmia and Septicæmia are very closely allied, and very frequently associated. They both owe their origin to the development and absorption of the same infective material, but the latter is a somewhat more simple process, and it will therefore be considered first.

Septicæmia.

By Septicæmia is understood the effects which are produced by the presence of putrid matter in the circulating blood.

When a person receives an external injury sufficient to cause an inflammatory process of considerable extent and intensity in the part

injured, this process is usually followed by a general disorder of the vital function, the most prominent symptom of which is pyrexia. This pyrexial state, which so commonly occurs after surgical operations and other extensive local injuries, is what is usually known as *traumatic* or *surgical fever*. Respecting the cause of this fever—it is mainly due to the absorption of some of the products of the local inflammatory process. In other words, it results from the dissemination of certain infective materials from a focus of acute inflammation. This being the case, whatever interferes with the removal of the inflammatory products from an injured part, and thus favors their absorption, tends to induce or to intensify this febrile state. This is exemplified by the well-known effects which are produced by pent-up pus, which effects subside after its evacuation.

Closely allied to this traumatic or surgical fever is the condition known as *septicæmia*. Here, also, there is an absorption of certain infective substances from some local lesion, but the general disturbance of the vital functions to which they give rise is much more considerable. Septicæmia appears therefore to differ from simple traumatic fever mainly in this—that in it the infective process is one of much greater intensity. No sharp line of demarcation, however, can be drawn between the two.

The clinical phenomena of septicæmia, as observed in man, are characterized not only by pyrexia, but also by vomiting, diarrhœa, muscular enfeeblement, affecting particularly the heart and respiratory muscles, and ultimately a condition of collapse which tends to terminate in death. After death the blood is found to be darker and less firmly coagulated than usual. Extreme congestions and ecchymoses are met with in internal organs, especially in the heart, lungs, and gastro-intestinal mucous membrane. The spleen, liver, and other viscera are enlarged, friable, and abnormally vascular; and little patches of lymph are seen on the pleura and pericardium. Metastatic abscesses and all other secondary inflammatory lesions are completely absent. Such pure uncomplicated cases of septicæmia are, however, comparatively infrequent, the septicæmia being usually associated with the development of secondary suppurative inflammations—a pyæmia. A simple septicæmia is perhaps most common in the puerperal state (Puerperal Septicæmia).

In studying the pathology of septicæmia we must consider—1. The nature of the infective material, the septicæmic poison, and the conditions under which it is produced; 2. The circumstances which in-


fluence its absorption and dissemination ; and 3. The changes which it produces in the circulating blood. Our knowledge on all these questions has been derived mainly from the experimental investigation of infective processes induced in the lower animals ; and amongst those who have contributed most largely to this knowledge during the past few years are Professors Billroth, Bergmann, and in our own country, Dr. Burdon Sanderson.¹

1. *The Septic Poison*.—Firstly, with regard to the infective material, and the circumstances under which it is produced. In the natural disease this material originates in connection with some local inflammation, or much less frequently with some wound in which no inflammation is present—as the uterus after delivery. It is in the albuminous liquids of such local lesions that the septic poison is produced, and its production appears to be invariably associated with the *putrid decomposition* of the liquid. In the disease artificially induced in the lower animals the poison is introduced from without, and consists of some putrid liquid, such as a putrid infusion of muscle.

It is well known that the putrid decomposition of albuminous liquids is always associated with the presence in them of *bacteria*, or their germs. (Fig. 78.) In a series of experiments made in the laboratory of the Brown Institution in 1872, with the object of determining the nature of the septic poison, Dr. Sanderson found that these, or similar organisms, abounded in all liquids which were capable, when introduced into the circulation of an animal, of producing septicæmia;² and from the results of these and other more recent experiments, Dr. Sanderson concludes that the *agency of bacteria is essential for the production of the septic poison*.

But although the production of the poison appears to be invariably associated with the presence of bacteria, it is something quite distinct from the process of simple putrefactive decomposition. That this is so has been proved by Bergmann, who has shown that the poison can be produced by the action of bacteria on non-albuminous liquids ; and also that if bacteria be grown in such non-albuminous liquids, the first

FIG. 78.



Fetid Pus from an Empyæmia.—Showing common active bacteria amongst the pus corpuscles. $\times 800$.

¹ "Lectures on the Infective Processes of Disease," by Dr. Burdon Sanderson, delivered at the University of London.—Brit. Med. Journ., Dec. 1877, and Jan. 1878.

² "Trans. Path. Soc. Lond.," vol. xxiii. 1872.

crop, which are rod-shaped, active organisms (bacteria proper) are inert; but as the process of cultivation proceeds, spheroidal and less active organisms are produced (micrococci), and the liquid becomes intensely virulent. From these facts Dr. Sanderson infers that the poison is not a product of the septic disintegration of protein substances, but something much more intimately associated with the existence and growth of the organisms themselves.

The production of the poison being thus dependent upon the agency of bacteria, the question arises whether its effects are due to the direct action of the organisms themselves. In answer to this question, the results of Dr. Ander's experiments appear to be conclusive. This observer has shown that the complete destruction of the organisms in a liquid which has been proved to be septic in no way impairs the virulence of its action.¹

From these investigations we must therefore conclude that the septic poison is a product of the growth of bacteria; but, says Dr. Sanderson, *bacteria are incapable of producing the poison in the healthy organism*. The condition which appears to be necessary for the development of the poison, is some abnormal state of the living tissues, such as is produced by injury. It is in the products of such injured tissues that the development of the bacteria and the production of the poison can take place, and it is consequently in some inflammatory lesion that septic processes most frequently originate.

Inasmuch as the agency of bacteria is essential for the production of the septic poison, it will be readily understood why, in the majority of cases, septicæmia originates from lesions which are in direct communication with the *external air*, and also why the development of septic processes is favored by the crowding together of persons with open wounds; for, as pointed out by Mr. Savory, the decomposition of an animal fluid is hastened by the introduction into it of any other animal matter which is also undergoing active putrefactive change.²

2. *The absorption of the Poison*.—Having discussed the nature of the poison, and the circumstances under which it originates, it remains to consider how it becomes absorbed and is disseminated. Putrid liquids in contact with injured tissues—liquids capable of producing the most intense septic processes if artificially inoculated—do not

¹ "Ander's, "Giftige Wirkung von durch Bakterien betrühte Nahrflüssigkeit." —*Deutsche Zeitschrift für Chirurgie*, vol. vii.

² "Discussion on Pyæmia at Clinical Society," Mr. Savory.—*Trans. Clin. Soc. Lond.*, vol. vii. p. lxxvi.

always become absorbed. The causes which in some cases favor their absorption, and in others prevent it, must be looked for, according to Mr. Savory (*loc. cit.*), in the condition of the tissues with which they are in contact. A fresh wound is a very readily absorbing surface, but Billroth and others have shown, experimentally, that healthy granulations offer a decided obstacle to the absorption of fluids from their surface. When, however, the granulations become partially destroyed, or are in an unhealthy condition, fluids readily permeate them. It must, then, be considered as exceedingly probable that the absorption of the poison is intimately connected with some abnormal condition, due to injury or disease, of the tissues with which it is in contact. The poison, when absorbed, is disseminated by means of the veins and lymphatics.

3. *Changes produced by the poison in the Blood.*—Respecting the changes produced by the septic poison in the circulating blood, and in the organism generally—the results of the experimental production of septicæmia show that these are in direct proportion to the quantity of the poison that is introduced. The poison does not multiply in the body; and in this respect, therefore, as will be seen presently, simple septicæmia appears to differ from the closely allied pyæmia.

The only other fact of importance in connection with this part of the subject relates to the marked tendency to congestive hyperæmia and blood-stasis which characterizes both the natural and the artificially induced disease. The investigations of Professor A. Schmidt and Dr. Köhler appear to prove that these circulatory phenomena are due to the destructive action of the poison on the white blood-corpuscles. Professor A. Schmidt has shown that the phenomenon of blood-coagulation is due to the white cells of the blood, that these cells contain a ferment, and that upon the destruction of some of them, and the consequent liberation of the ferment, the process of coagulation depends. From the experiments of Dr. Köhler¹ it would appear that in septicæmia there is a destruction of the white blood-corpuscles, and that to this destruction, and the consequent tendency to coagulation, the congestive phenomena which characterize the disease are owing. The other phenomena of septicæmia—the fever and collapse—however, are not thus accounted for. These may possibly be due to the action of the poison on the nerve-centres.

¹ "Ueber Thrombose und Transfusion, Eiter und septische Infection und deren Beziehung zum Fibrinferment," by Dr. Armin Köhler. Dorpat, 1877.

14

Pyæmia.

Pyæmia, as already stated, is the result of the development and absorption of the same poison as that which gives rise to septicæmia. The two conditions are thus closely allied, and they are very often associated. The former, however, is much the more frequent.

Pyæmia differs from septicæmia in this respect—that in it the absorption and dissemination of the infective poison gives rise not only to an alteration in the blood and a general disturbance of the vital functions, but also to the production of secondary foci of inflammation—the so-called *metastatic abscesses*. It is the production of these abscesses, and of other more diffused inflammatory lesions which is the distinctive character of pyæmia. It is thus probable that pyæmia is invariably associated with more or less septicæmia, and it may be regarded as a septicæmia in which there are metastatic suppurative inflammations.

Pyæmia being thus the result of the absorption of the same septic poison as that which gives rise to septicæmia, all that has been said respecting the origin and nature of this poison, and the circumstances which influence its absorption and dissemination from the focus of infection, applies equally to both diseases. What remains for consideration are the differences in the effects which are produced. It will be well, in the first place, however, to describe briefly the pyæmic lesions.

The lesions which are the most characteristic of pyæmia are the so-called metastatic abscesses. These present certain peculiarities. In their earlier stages they usually consist of somewhat reddish, friable, granular-looking masses of consolidation, which are surrounded by a thin zone of red hyperæmic tissue. These masses are frequently wedge-shaped, the apex of the cone being towards the centre of the organ. They vary considerably in size, some not being larger than a small pea, whilst others exceed the size of a chestnut. The consolidated mass soon assumes the characters of an abscess. Its more central portions become softened and purulent, and ultimately it tends to become converted into a collection of pus and broken-down tissue, this being still surrounded by a thin red zone of induration. These metastatic abscesses are met with most frequently in the lungs; but they also occur in the liver, spleen, kidneys, and in other internal organs. They are more commonly found near the surface than in

the more internal portions of the organ, often being situated immediately beneath the fibrous capsule.

In addition to these abscesses, more diffused inflammatory lesions are met with in pyæmia. These consist chiefly of suppurative arthritis, and of inflammations of the subcutaneous cellular tissue and serous membranes, especially of the pleuræ and pericardium.

With regard to the pathology of these metastatic abscesses, and other secondary inflammations—they are, for the most part, of *embolic* origin, the emboli originating in the thrombi which form at the seat of the primary lesion. The circumstances under which thrombi formed in the vessels of a part become softened or broken-up, so as to furnish embolic plugs, and the way in which these plugs may set up acute inflammatory processes at the seat of their arrest, will be fully described in the chapters on “Thrombosis” and “Embolism.”¹ It will be sufficient in the present place to state that in the course of a pyæmic process the coagula which form in the vessels at the seat of the pyæmic inflammations become infected with the pyæmic poison, and if they soften or become broken up so as to furnish embolic plugs, these plugs cause secondary suppurative processes in the parts in which they become arrested; and it is to the dissemination of these infective emboli that the formation of the majority of the metastatic abscesses in pyæmia is owing.

The prominent part which is played by thrombosis and embolism in the pathology of these infective processes probably accounts for the fact, that these processes are, in the majority of cases, associated with the development of metastatic abscess and other secondary inflammations—in other words, that pyæmia is so much more common than pure septicæmia. The bloodvessels of the primary lesion which serves as the starting-point of the infective process, usually a local inflammation, can rarely be free from coagula, and hence not only is the absorption of the poison often delayed (or even prevented), but when it does take place, it is usually associated with the dissemination of solid substances derived from the coagula themselves (emboli).

The occurrence of thrombosis and embolism also appears to influence the intensity of the infective processes. In simple septicæmia it has been seen that the changes produced are in direct proportion to the quantity of the poison which is absorbed from the primary lesion. In

¹ The student will do well to read the chapters on “Thrombosis” and “Embolism” as a sequel to the present chapter.

pyæmia, however, Dr. Sanderson says, the poison probably *multiplies in the organism*. In attempting to explain this, the experiments on the cultivation of the septic poison are of much value. These experiments prove not only that the poison can be produced by cultivation, but also that in an artificially produced septic liquid the virulence of the liquid increases with the development of the bacteria. A similar cultivation of the poison probably occurs in pyæmic processes. The original poison produced at the primary lesion increases in the secondary and tertiary foci of inflammation, and not only increases, but perhaps becomes more virulent.

CHAPTER XXIX.

SYPHILIS.

THE lesions occurring in the course of constitutional syphilis are for the most part inflammatory in their nature, but in their seat, distribution, and histological characters, many of them present certain peculiarities which make them quite characteristic of this disease. The primary syphilitic lesion (usually the indurated chancre), the secondary lymphatic gland enlargement, and the subsequent series of changes in the skin, mucous membranes, and later, in the nervous system, bones, and internal organs, are all of them the results of chronic processes, allied to inflammation, induced by the syphilitic poison.

The lesions, however, which must be regarded as especially characteristic of syphilis, are of three kinds—certain fibroid indurations, a form of growth which is known as a *gumma*, or syphilitic tumor, and certain changes in the arteries. These forms of lesion are very frequently associated, and in many cases it is difficult to draw any line of demarcation between them.

1. *Fibroid Changes*.—Fibroid indurations are often associated with the gummata, and many fibroid lesions are simply the remains of what were previously gummy tumors.

Fibroid changes also occur very frequently in the course of syphilis

Quite independently of gummata. These changes consist in the development of a fibroid tissue precisely similar to that met with as the result of ordinary chronic inflammation. (See "Inflammation of Common Connective Tissue.") The new tissue, which originates around the bloodvessels, and in the earlier stages of its growth is more or less richly cellular, produces a fibroid thickening of the affected area. These fibroid thickenings are characterized by their localization and by the irregularity in their distribution. They occupy, for the most part, *small areas*, and the *surrounding tissues are unaffected*. It is these peculiarities in their distribution and localization, and not their histological characters, which render them so characteristic of syphilis.


These fibroid changes are met with in various situations—in the periosteum, in the sheaths of nerve-trunks, in the capsules and interstitial tissue of organs, and in muscle. When occurring in the capsules of organs they present very characteristic appearances. In the liver and spleen, where they are the most frequent, they give rise to an irregularly distributed thickening and puckering of the capsule, which is exceedingly typical of advanced syphilis. The thickened portions are usually connected with dense fibroid septa which pass into the interior of the organ; and there are frequently also numerous strong peritoneal adhesions. The gummy masses, about to be described, are very often to be found in the midst of the fibroid growth.

2. *Gummata*.—These are perhaps more characteristic of syphilis than the fibroid lesions, with which, as already stated, they are frequently associated. The gummata, as usually met with, are moderately firm yellowish-white nodules, having often, on section, somewhat the appearance of a horse-chestnut. They vary in size from a hemp-seed to a walnut, and are surrounded by a zone of translucent fibrous-looking tissue, which sometimes has the appearance of a capsule, and which is so intimately associated with the surrounding structures that the enucleation of the mass is impossible. In the earlier stages of their development, when they less commonly come under observation, they are much softer in consistence, more vascular, and of a reddish-white color; whilst in their most advanced stages, owing to extensive retrogressive changes, they may be distinctly caseous.

When the gummata are examined microscopically they are found to consist in the main of atrophied, degenerated, and broken-down cell-products, imbedded in an incompletely fibrillated tissue. There are, however, some marked structural differences between the central and external portions of the growth. The central portions are com-


posed almost entirely of closely packed granular débris, fat granules, and cholestrin, amongst which may be an exceedingly scanty fibrillated tissue. (Fig. 79, *a*.) Surrounding this and directly continuous with it is a fibro-nucleated structure; whilst the peripheral portion of the growth is a richly cellular and vascular tissue. (Fig. 79, *b*, and Fig. 80.) This peripheral cellular tissue, which is in direct histological con-

FIG. 79.



Gummy growth from Liver.—*a*. Central portions of growth, consisting of granular debris. *b*, Peripheral granulation tissue. *r*. A bloodvessel. $\times 100$. (Cornil and Ranvier.)

FIG. 80.



The Peripheral Portion of a Gummy Growth in the Kidney.—Showing the small-celled granulation growth in the intertubular tissue. $\times 200$.

tinuity with the surrounding structures, consists of small cells, many of which resemble white blood-corpuscles, whilst others are larger and nucleated like the cells of granulation-tissue. These cells are separated by a scanty, homogeneous, intercellular material and numerous new bloodvessels.

FIG. 81.



From a Gummy Growth in the Kidney.—Showing the reticulated structure occasionally met with in the intermediate zone of these formations. $\times 200$.

These cells are separated by a scanty, homogeneous, intercellular material and numerous new bloodvessels.

The three zones above described, which are to be distinguished more or less clearly in most of the fully developed gummy nodules, correspond with three different stages in their growth. The most external zone, consisting of the vascular granulation-tissue, represents the early stage of development, and by the continuous formation of this tissue the growth may continuously increase. The intermediate more

fibrous zone represents the second stage in the process—the development of the granulation-tissue into a more or less completely fibrillated

structure. The characters of this fibrillated tissue vary in different growths. In some the fibrillation is very distinct; in others, the tissue is dense and cicatricial in character; whilst less frequently, it consists of a reticulated structure within the meshes of which are grouped round small cells. (Fig. 81.) The central zone, consisting of the amorphous granular material, represents the oldest portion of the growth—that which has undergone retrogressive changes. The bloodvessels, which only exist in the zone of proliferation, appear to become obliterated in the process of development, and this accounts for the rapid degeneration of the central portions of the growth. It is probable that this obliteration of the bloodvessels may be due to the changes in the arteries about to be described. When the tumor is large, it may sometimes be seen to be made up of several distinct smaller growths, each presenting at its circumference the more perfect cells, whilst its central parts are granular and amorphous.

It would thus appear that the first stage in the process of formation of these gummy growths consists in the production of a highly organized and vascular granulation tissue. This new tissue, although it may develop into an incompletely fibrillated structure, soon undergoes degeneration. The degenerated elements become closely packed in the centre of the growth, whilst proliferation and incomplete fibrillation continue at the circumference. The central portions of the growth occasionally become calcified, but more commonly they gradually become absorbed, and a mere puckered fibrous cicatrix may thus ultimately occupy the seat of the original tumor. This is one of the ways in which the fibroid lesions originate.

The gummata are the new formations most characteristic of syphilis. They are met with in the skin and subcutaneous cellular tissue, in the submucous tissue, in muscle, fasciæ, bone, and in the connective tissue of organs, especially of the liver, brain, testicle, and kidney. They also occur, but much less frequently, in the lungs; as do also simple localized fibroid indurations. When situated in the submucous tissue, the mucous membrane usually becomes destroyed, and a deep ulcer forms. This is seen in the pharynx, soft palate, tongue, larynx, and in other parts. These true gummy ulcerations must be distinguished from the superficial ulcerations resulting from inflammatory processes in the lymphatic structures which also occur in syphilis.


3. *Changes in Arteries.*—Certain changes in the *cerebral arteries* have been recently described by Heubner as characteristic of syphilis. These changes have been brought prominently before English patholo-

gists by Drs. Greenfield, Barlow, and others;¹ and the investigations of the first named of these observers would tend to render it probable that similar changes occur in the arteries in other situations.

In the cerebral arteries the changes produce opacity and marked thickening of the vessel, with considerable diminution in its calibre. It is this diminution of the lumen of the vessel which is especially characteristic.

When transverse sections of the vessels are examined microscopically, the principal change is seen to be situated in the *inner* coat. It is well shown in the accompanying drawings made from specimens kindly lent to me by Dr. Barlow. (Fig. 82.) This coat is con-

FIG. 82.



Syphilitic Disease of Cerebral Arteries.

A. Segment of middle cerebral artery, transverse section—*t*, thickened inner coat; *a*, endothelium; *f*, membrana fenestrata; *m*, muscular coat; *a*, adventitia. $\times 200$, reduced $\frac{1}{2}$.

B. Small artery of Pia Mater, transverse section. Showing thickened inner coat, diminished lumen of vessel, and considerable infiltration of adventitia. The cavity of the vessel is occupied by a clot (? thrombus). $\times 100$, reduced $\frac{1}{2}$.

rably thickened by a cellular growth. The growth, which is limited internally by the endothelium of the vessel (Fig. 82A. *e*), and externally by the membrana fenestrata (Fig. 82A. *f*), closely resembles ordinary granulation-tissue, consisting of numerous small round and

¹ "Trans. Path. Soc. Lond.," vol. xxviii.—*Visceral Syphilis*.

spindle-shaped cells. This tissue appears gradually to undergo partial development into an imperfectly fibrillated structure.

In addition to this change in the intima, the outer coat is abnormally vascular and infiltrated with small cells (Fig. 82A. *a*), and this cellular infiltration usually also invades the muscular layer (Fig. 82A. *m*).

The result of these changes in the inner coat is to diminish very considerably the lumen of the vessel (Fig. 82B); and the consequent interference with the circulation frequently leads to coagulation of the blood (thrombosis) and cerebral softening.

Dr. Greenfield's observations, as already stated, tend to show that similar arterial changes occur in other parts, and that they account for the degeneration of syphilitic gummata.

Syphilitic Disease of the Liver.

The liver is one of the most frequent seats of syphilitic lesions. The most common change is the development of fibroid and gummy growths in the substance of the organ. These growths are strictly localized, the surrounding liver-tissue being healthy. The growths—which are usually connected with fibroid thickenings of the capsule—sometimes consist simply of a dense fibroid structure. More commonly, however, gummy masses are found imbedded in the fibroid growth. In the former case it is possible that the gummy mass may have become absorbed, leaving merely its fibroid cicatrix.

The development of these growths produces very marked alterations in the form of the liver. Scar-like depressions are seen on its surface, and the organ is irregularly, and often very deeply puckered.

A more general fibroid change, not associated with the formation of gummy masses, is occasionally met with in the liver in inherited syphilis. This change closely resembles ordinary cirrhosis, although the intercellular network of the liver is usually more extensively involved.

Lastly, it must be mentioned that the liver in syphilis is frequently amyloid.

It is unnecessary to describe particularly syphilitic lesions in other organs, as they all present the same general characters—viz., strictly localized fibroid, or fibroid and gummy growths.

CHAPTER XXX.


INFLAMMATION OF NON-VASCULAR TISSUES.

Inflammation of Cartilage.

THE phenomena of acute inflammation in cartilage, although comparatively rare in man, have been studied by the artificial production of inflammation in the articular cartilages of the lower animals. They consist in changes in the cartilage itself, and in the vessels of the adjacent synovial membrane and bone, from which the cartilage receives its nutritive supply. Respecting the vascular changes, these are such as have been already described as characteristic of inflammation.

The changes which take place in the cartilage itself have been believed to be accompanied by activity of the cartilage-cells, by their proliferation, and the production of new elements. Such changes are represented in Fig. 83. The results of recent investigations, however, tend to show that the cells of the cartilage really remain inactive in the process of inflammation, and that all the young elements are emigrants from the vessels.

FIG. 83.



Section of Inflamed Cartilage.—*a.* The normal cartilage-cells. *b.* The same enlarged. *d.* Multiplication of cells within their capsules. *e.* Great increase in the number of the young cells, and destruction of the intercellular substance. $\times 250$. (Cornil and Ranvier.)

Whether the cells of the cartilage take any active part in the process or not, their capsules become destroyed, the intercellular substance softens and breaks down, and numerous small cells take the place of the original tissue. (Fig. 83, *e.*) As the process pro-

ceeds the new cells and granular débris escape from the surface of the cartilage, which thus becomes irregular, presenting numerous elevations and depressions: this is acute ulceration of cartilage. Whilst these changes are taking place in the cartilage itself, numerous new bloodvessels are formed from those in the adjacent bone or synovial membrane: the little loops of new vessels dipping into the softened cartilage. This process of softening and ulceration may go on until the cartilage is completely destroyed, or it may become arrested. In the latter case the young cells form a granulation-tissue, this tissue fibrillates, and the lost cartilage thus becomes replaced by a fibroid structure.

When the inflammation is less severe and runs a more chronic course, the cellular infiltration is less abundant, and the intercellular substance is less destroyed. The newly formed cells are more highly organized, and whilst many of them undergo retrogressive changes, others form a fibrillated tissue. Irregular cavities may thus be produced in the midst of a fibrillated cartilage.

Erosion of Cartilage.—Allusion must be made here to a morbid condition of cartilage which is exceedingly common in people past middle life. It consists in the occurrence upon the surface of the articular cartilages of small, yellowish, flocculent spots, which gradually increase in size, and ultimately break down and become destroyed, leaving a superficial erosion. The change occurs most frequently in the cartilages of the patella and of the knee-joint, and in other situations in which the cartilages are especially subjected to pressure.

When the diseased portions are examined microscopically, the most marked alteration is seen to consist in a fibrillation of the intercellular substance of the cartilage. Some of the cartilage-cells are also increased in size and show signs of proliferation; whilst others are in various stages of fatty metamorphosis. The change appears to result from long-continued irritation due to pressure, and it may be regarded as an irritative process, in which, owing to the natural poverty of the vascular supply, retrogressive changes occupy a very prominent place.

Inflammation of the Cornea.

The process of inflammation as it occurs in the cornea has been chiefly studied in the frog. After injury of the frog's cornea, the earliest changes observed consist in the conjunctival epithelium becoming visible, and in the appearance amongst the epithelial cells of

a few leucocytes, which have probably escaped from the hyperæmic vessels of the conjunctiva. The cornea cells and their prolongations then become visible (they are invisible in the healthy cornea, which appears perfectly structureless), and the intercellular substance gradually becomes increasingly opaque, owing to its infiltration with small cellular elements (leucocytes). These are so numerous and increase so rapidly, that they must undoubtedly be regarded as, in the main, emigrant white blood-corpuscles. It was formerly believed that the cells of the cornea exhibited active changes, and that from them many of the new elements which are seen in the intercellular substance were produced; but here, as in cartilage, all recent investigations render it more probable that they really remain inactive. As the number of young elements increases the consistence of the cornea becomes diminished, until ultimately the tissue breaks down and is destroyed.

The inflammatory process may continue until the whole of the substance of the cornea is softened and destroyed; or it may become arrested. In the latter case more or less thickening and opacity of the cornea will result, owing to the increase in the number of cells and the changes in the intercellular substance.

CHAPTER XXXI.

INFLAMMATION OF COMMON CONNECTIVE TISSUE.

COMMON connective tissue is one of the most frequent seats of the inflammatory process, not only the subcutaneous connective tissue, but also the connective tissue of organs and of other parts.

If connective tissue be examined a few hours after the infliction of an injury, it will be found that in place of the fibrillated substance and fixed connective-tissue cells of which it is normally composed, the tissue is infiltrated with small round cells (leucocytes), and that the fibrillated intercellular material has become homogeneous and gelatinous in consistence. The number of these cells gradually increases and the intercellular substance gradually becomes more completely destroyed, being probably consumed by the newly formed elements,

so that ultimately the tissue consists almost entirely of small round cells, held together by a very small quantity of soft gelatinous inter-cellular material.

Respecting the source from which these young elements are derived—*i. e.*, how far they are emigrants, and how far they are the offspring of cells belonging to the connective tissue—there appears to be little doubt that they are almost entirely emigrants. Although it was formerly supposed that the connective-tissue corpuscles multiplied very rapidly in inflammation, and that the newly formed cells were entirely the result of their proliferation, the investigations of Professors Cohnheim, Stricker, and others show that this view is erroneous; and according to Cohnheim, these corpuscles take no part whatever in the inflammatory process. Professor Stricker, however, has observed them undergoing active movements in the inflamed tongue of the frog, and although he has never seen them divide, he concludes that they may probably do so in the latter stages of the inflammatory process. The present state of our knowledge respecting inflammation of connective tissue would therefore appear to justify the conclusion that in the early stages of the process all the young cells are emigrants, but that in the latter stages a few of them may possibly be derived from the proliferation of the fixed connective-tissue cells.

Such being the nature of the changes which more immediately follow injury of the connective tissue, the inflammatory process may terminate in *resolution*, in *organization*, or in *suppuration*.


RESOLUTION.—If the injury sustained by the tissue is not severe, the inflammation may gradually subside, the process terminating in *resolution*. In this case the hyperæmia diminishes, the emigration ceases, some of the young cells undergo fatty metamorphosis and thus become absorbed, others pass into and are removed by the lymphatics, and the tissue gradually returns to its normal condition.

ORGANIZATION.—If the inflammatory process does not terminate in resolution, many of the young cells may become more fully developed, and ultimately form a fibrillated tissue. This *organization* of the inflammatory formation is seen in the healing of wounds by the “first intention,” and also in many of the chronic inflammations of the kidney, liver, and other organs. In order for it to occur it is necessary that there should be a considerable diminution in the intensity of the inflammation. (See “Chronic Inflammations.”)

The process of organization takes place by the production of new capillaries and by the development of the young cells into a granula-

tion-tissue. This granulation-tissue consists almost entirely of young cells, which must be regarded as exhibiting a somewhat higher stage of development than the primary amœboid cells (leucocytes) of the inflammatory formation. They are spherical masses of protoplasm from $\frac{1}{100}$ to $\frac{3}{100}$ of an inch in diameter, usually containing an indistinct round or roundly-oval nucleus, which often only becomes visible after the addition of acetic acid; and although they exhibit slight amœboid movements, they are much less active than the emigrant blood-corpuscles. The further organization of the growth takes place by the development of this granulation-tissue into a fibrillated or adenoid structure. The granulation-cells diminish in number, many of them frequently become spindle-shaped, the newly-formed capillaries gradually disappear, and the richly cellular tissue thus becomes converted into a more or less dense structure, which is characterized by the gradual process of contraction which it undergoes. This is usually known as *cicatricial tissue*.

FIG. 84.



Varieties of new Growth resulting from chronic Inflammation of Connective Tissue.—A an adenoid, B a fibroid structure. $\times 200$.


The characters of this new tissue present certain variations. Sometimes it consists of closely-packed wavy fibres, amongst which are a few elongated spindle-shaped elements, such as are represented in Fig. 84, B. It is equally common for the tissue to assume the appearance shown in Fig. 84, A—a dense homogeneous or obscurely fibrillated material, forming meshes of various sizes, within which are grouped a few lymphoid cells. (See also Figs. 67 and 102.) These two varieties are very frequently associated in the same specimen, the former probably representing a higher degree of development than the latter.

SUPPURATION.—When the injury sustained by the tissue is so severe or so prolonged in its action as to prevent the occurrence of

resolution or of immediate organization, the young cells infiltrate the tissue in such numbers that they may accumulate so as to constitute pus. The pus may either become collected together within the tissue so as to form an abscess, or it may be continuously discharged from the surface, as in a granulating wound. The deleterious influence which the pus exercises upon the tissues with which it comes in contact, causing their destruction and absorption, has already been alluded to. (See "Suppuration.")

Organization after Suppuration.—This constitutes what is generally known as healing by *granulation*, or by the *second intention*. It takes place in wounds in which, from the non-apposition of the wounded surfaces, union by the first intention has not been effected;

FIG. 85.



A Granulating Surface. *a.* Layer of pus. *b.* Granulation-tissue with loops of bloodvessels. *c.* Commencing development of the granulation-tissue into a fibrillated structure. $\times 200$. (Rindfleisch.)

and in other lesions in which the injured tissue presents a free surface communicating with the external air, as in an ulcer. The process of repair after the separation of a dead part (demarkation and separation) is in the same way effected by granulation, as is also the closing

a few leucocytes, which have probably escaped from the hyperæmic vessels of the conjunctiva. The cornea cells and their prolongations then become visible (they are invisible in the healthy cornea, which appears perfectly structureless), and the intercellular substance gradually becomes increasingly opaque, owing to its infiltration with small cellular elements (leucocytes). These are so numerous and increase so rapidly, that they must undoubtedly be regarded as, in the main, emigrant white blood-corpuscles. It was formerly believed that the cells of the cornea exhibited active changes, and that from them many of the new elements which are seen in the intercellular substance were produced; but here, as in cartilage, all recent investigations render it more probable that they really remain inactive. As the number of young elements increases the consistence of the cornea becomes diminished, until ultimately the tissue breaks down and is destroyed.

The inflammatory process may continue until the whole of the substance of the cornea is softened and destroyed; or it may become arrested. In the latter case more or less thickening and opacity of the cornea will result, owing to the increase in the number of cells and the changes in the intercellular substance.

CHAPTER XXXI.

INFLAMMATION OF COMMON CONNECTIVE TISSUE.

COMMON connective tissue is one of the most frequent seats of the inflammatory process, not only the subcutaneous connective tissue, but also the connective tissue of organs and of other parts.

If connective tissue be examined a few hours after the infliction of an injury, it will be found that in place of the fibrillated substance and fixed connective-tissue cells of which it is normally composed, the tissue is infiltrated with small round cells (leucocytes), and that the fibrillated intercellular material has become homogeneous and gelatinous in consistence. The number of these cells gradually increases and the intercellular substance gradually becomes more completely destroyed, being probably consumed by the newly formed elements,

so that ultimately the tissue consists almost entirely of small round cells, held together by a very small quantity of soft gelatinous inter-cellular material.

Respecting the source from which these young elements are derived—*i. e.*, how far they are emigrants, and how far they are the offspring of cells belonging to the connective tissue—there appears to be little doubt that they are almost entirely emigrants. Although it was formerly supposed that the connective-tissue corpuscles multiplied very rapidly in inflammation, and that the newly formed cells were entirely the result of their proliferation, the investigations of Professors Cohnheim, Stricker, and others show that this view is erroneous; and according to Cohnheim, these corpuscles take no part whatever in the inflammatory process. Professor Stricker, however, has observed them undergoing active movements in the inflamed tongue of the frog, and although he has never seen them divide, he concludes that they may probably do so in the latter stages of the inflammatory process. The present state of our knowledge respecting inflammation of connective tissue would therefore appear to justify the conclusion that in the early stages of the process all the young cells are emigrants, but that in the latter stages a few of them may possibly be derived from the proliferation of the fixed connective-tissue cells.

Such being the nature of the changes which more immediately follow injury of the connective tissue, the inflammatory process may terminate in *resolution*, in *organization*, or in *suppuration*.

RESOLUTION.—If the injury sustained by the tissue is not severe, the inflammation may gradually subside, the process terminating in *resolution*. In this case the hyperæmia diminishes, the emigration ceases, some of the young cells undergo fatty metamorphosis and thus become absorbed, others pass into and are removed by the lymphatics, and the tissue gradually returns to its normal condition.

ORGANIZATION.—If the inflammatory process does not terminate in resolution, many of the young cells may become more fully developed, and ultimately form a fibrillated tissue. This *organization* of the inflammatory formation is seen in the healing of wounds by the “first intention,” and also in many of the chronic inflammations of the kidney, liver, and other organs. In order for it to occur it is necessary that there should be a considerable diminution in the intensity of the inflammation. (See “Chronic Inflammations.”)

The process of organization takes place by the production of new capillaries and by the development of the young cells into a granula-

ossification.)¹ The process thus causes enlargement of the bone, or if this be necrosed, it helps to reproduce the lost structure. (See "Necrosis.")

Strictly localized inflammations of the periosteum occur in the formation of nodes, and here also there is frequently a new growth of bone.

NECROSIS.—All conditions interfering with the supply of blood to a bone may be causes of its death. Necrosis may thus result from accident, portions of bone being violently separated from their vascular connections, or, more commonly, the periosteum being stripped off and injured. Much more frequently, however, it is the result of the intense inflammatory processes in the bone or periosteum which have been described.

The bone, when dead, undergoes a gradual process of separation from the adjacent living tissue. This is effected in the same way as the separation of soft structures. (See "Gangrene.") The adjacent living bone becomes the seat of an inflammatory process, a granulation-tissue and pus are produced, and by this means the dead portion is completely separated. When separated it is termed a *sequestrum*.

The removal of the sequestrum from the granulation-tissue with which it is in contact is often only effected with considerable difficulty, especially if it be deeply seated. This difficulty is occasionally due to a more or less thick layer of the old bone surrounding the necrosed portion. Much more frequently, however, it is owing to the participation of the periosteum in the inflammatory process. The inflamed periosteum produces new bone, a capsule of which is thus formed, inclosing the sequestrum. Openings exist in this capsule (*cloacæ*) leading to the dead bone, and through these openings the inflammatory products are discharged. When the sequestrum is quite superficial, its removal is, of course, more readily effected.

CARIES.—By caries is understood that inflammatory disintegration of bone and removal of the dead products which corresponds with ulceration of soft parts. Caries is therefore one of the results of inflammation of osseous tissue.

Inflammatory processes in bone which are of comparatively slight intensity have been seen to be characterized histologically by a gradu-

¹ According to Billroth, this new osseous tissue is partly derived from the bone itself, which, he states, is the seat of a superficial osteitis. (See "Billroth's Surgical Pathology and Therapeutics," translated by Hackley, 4th edit., p. 419.)

ally increasing production of an inflammatory granulation-tissue from the medullary tissue of the bone, and by the absorption of the compact osseous structure. This absorption of the bone appears to be effected entirely by the new inflammatory growth. When these chronic inflammatory processes occur in the superficial layers of the bone beneath the periosteum, the bone is gradually destroyed, and presents a ragged, irregular excavated surface. This is superficial caries, or ulceration of bone. The floor of the ulcer consists of a cancellous structure containing the débris of the disintegrating process. In many cases there is a marked condensation (sclerosis) of the deeper portions of the bone.

Chronic inflammatory processes in the interior of bone cause in the same way absorption and disintegration of the osseous structure. Here, however, the inflammatory products being less readily removed, sometimes undergo, especially in scrofulous children, a kind of caseous transformation, and the carious cavity contains molecular débris, sanious pus, and minute portions of dead bone. The inflammatory granulation-tissue often projects as fungating masses into the cavity.

Caries is frequently associated with more or less necrosis, especially when the process affects the interior of the bone. Small portions of bone are cut off from their vascular supply, become detached, and are found in the carious cavity.

There are two other morbid conditions of bone, which, although probably not coming within the category of inflammation, may be conveniently described in the present chapter—viz., *Mollities Ossium* and *Rickets*.

Mollities Ossium.

Mollities Ossium or Osteomalacia is a rare disease, occurring only in adults. It is characterized by a progressive softening of the bones, owing to an increase in their medullary tissue and the destruction of the compact osseous structure. The bones become so much softened that they can be easily cut with a knife; they are exceedingly light, and bend readily in all directions. On section, the cortical layer is found to be almost destroyed, the bone consisting of a wide cancellous structure containing a reddish, gelatinous, fatty material.

The nature of the disease is obscure. According to Rindfleisch, the change consists in a decalcification of the osseous tissue spreading gradually from the medulla and cancelli of the bone. The decalcified

tissue does not become so completely absorbed as in inflammation of bone, but undergoes more or less of a mucoid change. The medullary tissue is exceedingly vascular, but unlike ordinary granulation-tissue it usually contains much fat. Lactic acid has been found in the bone and in the urine. The urine usually also contains lime salts, which have been removed from the bone and eliminated by the kidneys.

Rickets.

The common disease of children so well known as Rickets, or Rachitis, is mainly characterized by certain alterations in the bones. The bones generally are softened. The ends of the long bones are enlarged, and the flat bones are thickened. As a result of these alterations the bones become deformed, and their growth is frequently arrested.

Respecting the nature of this disease, we are at present unable to speak certainly. It appears, however, to be closely associated with mal-nutrition, and all those conditions which materially interfere with the nutrition of the child may be causes of the disease.

The alteration in the bones may be briefly described—as stated by Sir W. Jenner—as consisting in an increased preparation for ossification, but an incomplete performance of the process. In a growing bone the zone of soft tissue in which the process of conversion of cartilage into bone is taking place is exceedingly narrow, the changes in the cartilage involving only one or two rows of the cartilage cells. In rickets this transition zone is very considerably widened, the changes in the cartilage cells extending for some depth into the cartilage, and the adjacent layer of cancellous tissue being much broader than natural. The junction between the bone and the cartilage is also exceedingly irregular, so that in some parts the cancellous bone extends into the cartilage much further than in others. Usually the alterations in the cartilage in the transition zone are accompanied by the calcification of the new tissue and its conversion into bone. In rickets, however, this calcification takes place very incompletely, and thus a wide layer of imperfectly calcified soft tissue is formed at the growing end of the bone.

A similar abnormality exists in the process of ossification from the periosteum. Here also the zone of vascular granulation-tissue from which the bone is developed is exceedingly broad, the calcification of the tissue is exceedingly incomplete, and the bone thus becomes thick-

ened by a soft, vascular, and very imperfectly calcified structure. As the bone grows, the medullary cavity naturally widens, and the compact tissue becomes absorbed. This imperfect formation of osseous tissue from the periosteum produces, therefore, considerable softening and weakening of the bone.

The changes in the bones in rickets thus closely resemble those which occur in the normal process of ossification. There is the same change in the cartilage cells and in the deeper periosteal layers, but this change involves abnormally large areas. The development of the young cells and the transformation of the new tissue appear also to be more or less incomplete. In many parts the young elements do not form perfect bone-cells, but simply angular cells without canaliculi, whilst in others the development may be much more complete. Various transitions are thus met with between tolerably well developed and very imperfect bone-structure, but owing to the deficiency of lime salts this structure is very imperfectly and very irregularly calcified.

CHAPTER XXXIII.

INFLAMMATION OF BLOODVESSELS.

Inflammation of Arteries.


IN studying the process of inflammation in arteries, it must be borne in mind that the inner coat of these vessels is non-vascular, the blood-vessels being distributed in the external layers.

Acute and chronic inflammations of arteries are a very frequent result of coagulation of the blood within the vessel (thrombosis). Such inflammations will be considered in the chapter on Thrombosis. Independently of thrombi, the most acute inflammatory processes in arteries are for the most part the results of injuries from violence, or of inflammation in adjacent tissues. Here, the histological changes are principally confined to the *external* coat of the vessel. This becomes exceedingly hyperæmic and infiltrated with small cells, the cellular infiltration involving also the middle coat. As a consequence

of these changes in the outer coats the vitality of the intima is impaired or destroyed, and when this has occurred the blood contained within the vessel coagulates.

ATHEROMA.—The more chronic forms of arterial inflammation give rise to changes which are most marked in the deeper layers of the *inner* coat, where they produce those various alterations in the walls of the vessel included under the common term—*atheroma*. In the earliest stage of the inflammatory process the fibrous and elastic lamellæ of the inner coat of the vessel become infiltrated with small cells, which are probably partly emigrants, and partly derived from the cells of these structures. As these cells increase in number they give rise to a swelling beneath the innermost layers of this coat of the artery (Fig. 86). This swelling of the intima is very characteristic.

FIG. 86.



Atheroma of the Aorta.—Showing the cellular infiltration of the deeper layers of the inner coat, and the consequent internal bulging of the vessel. The new tissue has undergone more or less fatty degeneration. There is also some cellular infiltration of the middle coat. *f.* internal, *m.* middle, *e.* external coat of vessel. $\times 50$, reduced $\frac{1}{2}$.

It is, in the earlier stage of the process, of a soft flabby consistence, and the lining membrane which is continuous over it can be stripped off, leaving the diseased tissue beneath. It thus contrasts strongly with the superficial patches of fatty degeneration which result from the fatty metamorphosis of the endothelial and connective-tissue cells of the vessel. (See “Fatty Degeneration of Arteries.”)

The results of the inflammatory process will depend upon its severity. If the process be very acute, the young elements accumulate rapidly and infiltrate the middle and external coats, so that the walls of the vessel become so much softened, that dilatation, aneurism, or rupture

may ensue. Such acute changes are a common cause of aneurism. The inflammation, however, is usually of less intensity and runs a more chronic course, and it is only to these more *chronic* arterial changes that the term "atheroma" is commonly applied.

When the process is less intense and more chronic in its course, it leads to various alterations in the wall of the vessel. The soft cellular infiltration of the deeper layers of the intima usually undergoes retrogressive changes, owing to deficient nutritive supply. These changes commence in the deeper portions of the new tissue. The young cells undergo fatty degeneration, the intercellular substance softens, and thus a soft, yellowish, pultaceous material may be produced beneath the lining membrane of the vessel. This has been termed an *atheromatous abscess*. The lining membrane may ultimately give way and the softened matters be carried away by the circulation, and thus is produced the *atheromatous ulcer*. In other cases the more liquid constituents of the degenerated tissue gradually become absorbed, cholesterin forms, and thus a mass, consisting of broken-down fibres and cells, fatty débris, and cholesterin crystals, with a varying quantity of the original fibrillated tissue, remains in the deeper layers of the inner coat. (Fig. 86.) This may subsequently calcify, and so form a *calcareous plate*.

In the most chronic forms of the atheromatous process there is more or less fibrillation of the new tissue, and thus is produced a *fibroid thickening* of the inner coat of the artery. The organization, however, is rarely complete, more or less fatty débris being usually inclosed in the fibroid stroma.

Respecting the causes of these chronic arterial inflammations included under the common term of atheroma—they are probably mainly due to over-strain of the vessel, the strain exercising a chronic injurious influence. This view has been especially insisted upon by Dr. Moxon.¹ It is supported by the fact that those vessels which are the most exposed to strain, are the most liable to this disease—*e. g.*, the arch of the aorta; also by the exceeding frequency of atheroma in all those conditions in which there is an increase of the blood-pressure. The effect of increased blood-pressure as a cause of atheroma is seen in the liability of athletes to the disease; and also in the frequency with which it occurs in chronic Bright's disease, and in the pulmonary

¹ "Lectures on Pathological Anatomy." Wilks and Moxon, 2d Edit., p. 150.

vessels in cases of mitral obstruction, etc. The occurrence of atheroma is also favored by syphilis.

Inflammation of Veins.

Inflammatory processes in veins are more frequent than in arteries, but here they are in the very great majority of cases secondary to coagulation of the blood within the vein (thrombosis), the coagulum exercising an injurious influence upon the coats of the vessel. These inflammations resulting from thrombosis will be treated of in a subsequent chapter. (See "Results of Thrombosis.")

The inner coat of the veins is, like that of the arteries, non-vascular, and hence acute inflammatory changes are, for the most part, confined to the external and middle coats. Independently of inflammations resulting from thrombosis, the lining membrane of veins appears to undergo no inflammatory changes.

Acute and chronic inflammatory processes in veins are, as already stated, most frequently due to thrombosis. They may also result from violent injury, or from the extension of inflammation from adjacent tissues. The process, when acute, closely resembles that in the arteries. The external and middle coats become infiltrated with cells, the vitality of the intima ultimately becomes impaired or lost, and when this has occurred the blood within the vein coagulates.

CHAPTER XXXIV.

INFLAMMATION OF THE HEART.


INFLAMMATORY processes in the heart may affect the substance of the organ, or the endocardium. They are much more frequent in the last-named situation.

Endocarditis.

Endocarditis is for the most part limited to the valves of the heart, although it is occasionally met with involving more or less of the cardiac cavities. The process is almost exclusively confined to the


left side of the organ, and in the great majority of cases it commences in, and comparatively rarely extends beyond, the confines of the aortic and mitral valves and the corresponding orifices. Further, it is those portions of the valves which come into contact in the act of

FIG. 87.



Inflammation of Aortic Valves.—The earlier stage of the process. Showing the situation of the inflammatory granulations.

FIG. 88.




Inflammation of Mitral Valves.—The earlier stage of process. Valve seen from the auricular surface. Showing the situation of the inflammatory granulations.

closure, and are thus *most exposed to friction*, which are especially involved, and in which the changes usually commence. Thus, in the aortic valves, it is the convex surfaces of the segments which are most liable to be affected, and not the free edge of the segment, but the little band of tissue which passes from its attached border to the corpus Arantii in the centre (Fig. 87); and in the mitral valve, the auricular surface of the segments at a little distance from the attachment of the chordæ tendineæ. (Fig. 88.) When portions of the endocardium apart from the valves are affected, this is frequently due, as pointed out by Dr. Moxon, to the irritation caused by the friction of vegetations or fibrinous clots situated on the valves themselves. (Fig. 89.)

The histological changes accompanying endocarditis closely resemble those already described as occurring in those more chronic forms of arterial inflammation known as atheromatous. The endocardium and inner coat of an artery are very analogous in their structure,

FIG. 89.



Endocarditis due to Friction.—The drawing represents a long vegetation on one of the segments of the aortic valve, which by rubbing on the endocardium below has produced numerous inflammatory granulations (A).

both being non-vascular, and consisting of a layer of connective tissue with an internal endothelial covering. The inflammatory process may be acute or chronic.

ACUTE ENDOCARDITIS.—If the process be acute, the deeper layers of the endocardium become rapidly infiltrated with young cells, and as these increase in number the intercellular substance becomes softened and destroyed, and thus is produced a soft tissue composed almost entirely of cells such as always results from inflammatory processes in connective tissue. The new tissue as it increases tilts up the superjacent endothelium and projects in the form of minute granulations and vegetations upon the surface of the softened valve. (See Figs. 87 and 88.) The endothelial elements probably also participate in the active process.

The above changes take place in an almost non-vascular tissue, and although there is more or less increase of vascularity in the more external endocardial layers, where the capillaries are more numerous, there is rarely any redness or injection of the endocardium seen after death. Neither is there usually any liquid exuded upon the surface

of the granulations. What was formerly regarded as an exuded material, is in the main coagulum which has been deposited from the blood upon the roughened surface of the valve. This deposition of coagulum frequently occurs in endocarditis, the roughened and abnormal endocardium acts as a foreign body and so causes a deposition upon its surface. This must not be confounded with the vegetations themselves. (Fig. 90.)



Acute Endocarditis.—A granulation from the mitral valve, showing a fibrinous coagulum upon the surface of the granulation. $\times 10$. (Rindfleisch.)

The results of this cellular infiltration vary. If the process be very acute the new tissue may break down, and thus a loss of substance result, an endocardial ulcer. This takes place without any accumulation of cells sufficient to form an abscess, the new tissue simply becoming rapidly softened and disintegrating. In rare cases, however, small quantities of pus are found in the deeper endocardial layers. The ulcer is irregularly defined, and its edges are usually swelled and thickened. This *ulcerative endocarditis* is, however, not frequent, the process usually being less acute. The ulceration may lead to perforation of the valve, or to a considerable destruction of its substance. Laceration or aneurism of the valve may also ensue from the pressure exercised by the blood against the damaged tissue. Sometimes the

ulcerative process extends so as to involve the cardiac substance. Ulcerative endocarditis is a grave affection, often giving rise to embolism, and sometimes to a pyæmic process.

When the inflammatory process is less severe, as is much more commonly the case, the new tissue becomes incompletely organized into a fibrillated structure, whilst it undergoes, in part, fatty and calcareous degeneration. These changes may result in the adhesion of the valves, either to one another or to the walls of the heart. They always produce permanent *thickening*, *rigidity*, and *shrinking* of their structure. The new tissue may continue to grow after the severity of the process has subsided, and thus are produced the vegetations and papillary excrescences on the valves which are so commonly met with. (See Fig. 89.) These consist of a lowly organized tissue, which tends to undergo fatty and calcareous changes.

CHRONIC ENDOCARDITIS.—This may be the sequel of acute inflammation, or the process may from its commencement be chronic in its nature. In chronic endocarditis the cellular infiltration is much less rapid and abundant than in the acute form; the intercellular substance consequently becomes much less softened and destroyed, and the new tissue has a much greater tendency to become developed into a fibrillated structure. The result of these chronic processes is therefore to produce a *fibroid thickening* of the endocardium, and so to cause considerable induration and contraction of the valves or valvular orifices. The new tissue sometimes forms papillary growths on the valves, which undergo partial fatty and calcareous changes. (See Fig. 89.)


Myocarditis.

Myocarditis, or inflammation of the cardiac substance, is much less frequent than the preceding. Intense and concentrated inflammations leading to the formation of abscess probably occur only as the result of a pyæmic process. Less intense and more diffuse forms of cardiac inflammation are also not unfrequently met with in association with pericarditis, and, less commonly, with endocarditis. Here the inflammatory process appears, by extension, to involve the immediately adjacent muscular layers of the organ, which are found infiltrated with small cells, the fibres themselves being softened and granular.

In addition to the above, a form of myocarditis must be recognized in which the substance of the heart becomes more generally involved.

In certain cases of acute rheumatism the muscular tissue of the heart is found after death swollen, softened, opaque, and occasionally faintly mottled with slightly yellowish patches. When examined microscopically, the fibres are seen to have lost their striation and to be finely granular, their nuclei are large and prominent, and small cells are found in varying numbers infiltrating the intermuscular tissue. (Fig. 91.) I have met with these appearances in two or three cases of

FIG. 91.



Acute Myocarditis.—From a case of acute rheumatism. *a.* A thin section of the left ventricle made in the direction of the muscular fibres, showing the granular and swollen condition of the fibres, and the prominence of their nuclei. *b.* A transverse section, showing the cellular infiltration of the intermuscular tissue. $\times 200$.

acute rheumatism, and they must, I think, be regarded as evidence of the existence of an acute inflammatory process. The change is most marked in the left ventricle, and it is usually associated with endo- or pericarditis. It is a grave complication of acute rheumatism, and possibly of some other diseases, and is probably more frequent than is generally supposed.

FIBROID INDURATION OF THE HEART.—This, a comparatively rare condition, is probably, in most cases, a result of myocarditis. The change is characterized by the development of a fibrillated tissue between the muscular elements. The process commences in the intermuscular septa around the bloodvessels. This becomes infiltrated with small cells, which tend to become developed into a fibrillated structure. (Fig. 92.) The growth of new tissue gradually extends between the bundles of muscular fibres, so that ultimately each fibre may be surrounded by a tract of dense fibroid tissue. (Fig. 93.) The muscular fibres themselves, owing to the resulting interference with their nutritive supply, atrophy, undergo fatty meta-


morphosis, and are gradually replaced by the fibroid growth. (Fig. 93.) Very frequently the cellular nature of the growth, which I believe to characterize the earlier stages of its development, is not seen, the new tissue being simply fibroid.¹

FIG. 92.



Fibroid Induration of the Heart.—A thin section from the wall of the left ventricle, showing the small-celled growth in the intermuscular septa around the bloodvessels. *a, a*, vessels. $\times 200$.

FIG. 93.



Fibroid Induration of the Heart.—A section from the left ventricle of the same heart as Fig. 92, showing a more advanced stage. The fibroid tissue surrounds the individual muscular fibres, which are undergoing fatty degeneration. $\times 200$.

Fibroid induration of the heart appears in most cases to be induced by inflammatory processes commencing in the peri- or endocardium. When secondary to pericarditis, the change is usually most advanced in the more external portions of the cardiac walls, and it commonly affects both the right and left ventricles. When, on the other hand, an endocarditis is the precursor of the indurative process, the change is more marked in the internal muscular layers, and inasmuch as inflammatory processes in the endocardium occur almost exclusively in the left cardiac cavities, the left ventricle is principally involved. In other cases the fibroid growth appears to be the result of syphilis. (See "Syphilis.")

Although the growth of new tissue is thus usually more advanced in certain portions of the muscular walls than in others, it is by no means uniformly distributed. In some parts it may be very dense, the muscular fibres being entirely obliterated, whilst in others it is entirely wanting, and the muscular elements present a normal appearance.

¹ Dr. Hilton Fagge, in a series of eleven cases of fibroid disease of the heart, found that cellular elements in the new growth were almost invariably absent. (See "Trans. Path. Soc. Lond.," vol. xxv. p. 64.)

Fibroid induration of the heart—excluding that resulting from syphilis—appears to occupy the same pathological position as similar fibroid changes in other organs—*e. g.*, in the liver and kidneys. It must therefore be regarded as the result of a chronic inflammatory process, which might be termed *chronic myocarditis*. Its effect must evidently be to interfere very materially with the motor power of the organ, and it consequently constitutes one of the most grave of all cardiac diseases.

The cardiac walls may become much thickened by the new growth, and the induration of texture is often very considerable. In the specimen from which the accompanying drawings were made, the disease was apparently secondary to an endocarditis, and the walls of the left ventricle were so hard that they cut almost like a piece of tendon.

CHAPTER XXXV.

INFLAMMATION OF LYMPHATIC STRUCTURES.

INFLAMMATORY processes in lymphatic structures usually result from their injury by substances conveyed to them by the lymphatic vessels. They include—*acute* and *chronic* inflammations, and the specific inflammation associated with *Typhoid Fever*. Each of these must be considered separately.

Acute Inflammation of Lymphatic Structures.

Examples of acute inflammation of lymphatic structures are furnished by the inflammation of the glands in the axilla from a wound on the hand, of the glands in the groin from gonorrhœa, and of Peyer's and the solitary glands in the intestine from inflammation of the intestinal mucous membrane.

The process consists in a hyperæmia of the gland together with a rapid increase in the number of the lymph-cells. This increase is probably due, partly to a hyperplasia of the original cells of the gland, and partly to the migration of blood-corpuscles. The cells not only increase in number, but many of them become much larger in size, and

their nuclei multiply. The cells of the tuberculæ also participate in the active process. The increase in the size of the elements and the multiplication of their nuclei usually lead to the production of some larger cells containing three or four nuclei. These have been already alluded to as being frequently met with in lymphatic structures which are undergoing active processes. (See Fig. 37.) Owing to these changes the gland becomes considerably increased in size, soft and pulpy in consistence, and its cortical and medullary parts are no longer distinguishable.

Upon the removal of the injurious influence the process may gradually subside, the new elements undergo disintegration and absorption, and the gland returns to its normal condition (Resolution). In other cases the process goes on to suppuration, the trabeculæ are destroyed, many of the cells become disintegrated, and the loculi of the gland become filled with pus. This is usually associated with inflammation and suppuration of the surrounding connective tissue. In the glands of a mucous membrane the process gives rise to what is known as a follicular abscess.

Chronic Inflammation of Lymphatic Structures.

Chronic inflammations of lymphatic structures result from injuries which are less severe and more prolonged in their action than those which give rise to the acute form. The resulting cellular infiltration of the gland is consequently a more continuous one, and the gland becomes more or less permanently increased in size. The reticulum is also considerably involved. These chronic inflammatory processes differ from the acute, inasmuch as they lead to a gradually increasing development of the reticular structure of the gland. The reticulated network becomes thicker and more fibrous, its meshes become smaller and smaller, the lymph-cells diminish in number, and thus the gland becomes hard and fibrous in consistence. (Fig. 94.)




FIG. 94.

Chronic Inflammation of a Lymphatic Gland.—Showing the increase in the stroma, and the diminution in the number of the lymphoid cells. $\times 200$.

Scrofulous Glands.—In those chronic inflammations of the lymphatic glands which occur in scrofulous subjects, and in which the glands

tend to become caseous, the changes resemble those which have been already described as characteristic of scrofulous inflammation. (See "Scrofulous Inflammation.") The cellular infiltration is considerable, there is but little tendency to absorption, and many of the cells increase in size, and even form multinucleated elements. The gland thus becomes enlarged, soft, and elastic in consistence, and of a uniform grayish-white color. Owing to the obstruction to the circulation caused by the pressure of the cellular infiltration, the gland undergoes retrogressive changes and becomes caseous. The caseous material may subsequently liquefy, or become infiltrated with calcareous particles. Many caseous lymphatic glands are tuberculous. (See "Tuberculosis of Lymphatic Glands.")

Inflammation of Lymphatic Structures in Typhoid Fever.

The inflammatory processes which occur in the lymphatic structures in Typhoid Fever, have their seat in the spleen, in the lymphatic structures of the intestine, and in the mesenteric glands.

The Spleen.—In the spleen the change resembles that which occurs in many of the other acute febrile diseases, although it reaches its maximum in typhoid. The splenic tissue becomes exceedingly vascular and the lymphatic elements increase rapidly in number, so that the organ often attains two or three times its natural size. Many of the new elements enter the blood, thus causing a slight temporary increase in the number of white blood-corpuscles. As the fever subsides, the hyperæmia diminishes, many of the new elements undergo disintegration and absorption, the remainder enter the blood, and thus the organ again attains its normal characters and dimensions.

The Intestinal Lymphatic Structures.—It is in the solitary and Peyer's glands that the most characteristic changes take place in typhoid fever. These structures may be involved throughout the whole of the small and large intestine, but in most cases the process is limited to those in the ileum and cæcum; and those glands are always the most affected which are situated the nearest to the ileo-cæcal valve.

The primary change here consists in a hyperæmia and cellular infiltration of the glands. Many of the cells increase considerably in size, so as to form the multinucleated elements already alluded to. Both Peyer's patches and the solitary glands thus become considerably enlarged and prominent, standing up above the surface of the

intestine. (Fig. 95.) They are of a grayish-white or pale reddish color, and of a soft, brain-like consistence. The surrounding mucous membrane is also exceedingly vascular, and is the seat of an acute catarrhal process. This catarrh is more or less general, and usually precedes the swelling of the glands. The cellular infiltration, in many parts, rapidly extends beyond the confines of the glands into the immediately surrounding and subjacent tissues, and even in some cases into the muscular coat.





Fig. 95.

Typhoid Swelling of Peyer's Patches and Solitary Glands of the Intestine.

The process now passes into the second stage—that of the death and disintegration of the newly formed tissue. This may terminate in various ways. The enlarged glands, many of them, subside, the new elements become disintegrated and are absorbed, and the gland thus undergoes a gradual process of *resolution*. In others, the individual follicles of the gland rupture, discharging their contents externally, and the patches then acquire a peculiar reticulated appearance. The most characteristic termination, however, of the typhoid process, is the separation of the dead tissue as a slough, and the formation of the *typhoid ulcer*.

The process of sloughing and ulceration may, like the cellular infiltration, take place uniformly throughout the whole gland, in which case the whole mass is thrown off, leaving an ulcerated surface corresponding in size with that of the gland. (Fig. 96.) More commonly, however, in the patches, the sloughing commences in different portions of the patch, and small irregular losses of substance result, which may gradually extend until they form one large ulcer.

Fig. 96.



A Typhoid Ulcer of the Intestine.

Although, as already stated, the cellular infiltration may extend beyond the confines of the glands, this is rarely the case with the ulceration. The peripheral infiltration undergoes resolution, and hence the ulcers have the same configuration as the original glands; those originating from the patches being oval, with their long diame-

ter in the direction of the gut; and those originating in the solitary glands being spherical in shape. In rare cases, when there is much infiltration of the surrounding mucous membrane, the ulceration may extend slightly beyond the confines of the glands.

With the sloughing and disintegration of the new tissue the process of infiltration ceases, and hence there is no induration or thickening of the base or edges of the ulcer. The

FIG. 97.



*A Typhoid Ulcer of the Intestine (diagrammatic).—*Showing the undermined edges of the ulcer, and the slough still adherent. *a.* Epithelial lining. *b.* Submucous tissue. *c.* Muscular coat. *d.* Peritoneum.

base is smooth, and is usually formed of the submucous or muscular coat of the intestine. The edges are thin and undermined, and consist of a well-defined fringe of congested mucous membrane. (Fig. 97.) This is best seen when the gut is floated in water. In some cases, however, the sloughing extends deeper

through the muscular layer to the sub-peritoneal tissue, and it may thus cause perforation and peritonitis.

The third stage of the process is that of cicatrization. This takes place by the resolution of the peripheral infiltration, the approximation and union of the undermined edges with the floor of the ulcer, and the gradual formation from the margin of an epithelial covering. The gland-structure is not regenerated. The resulting cicatrix is slightly depressed, and less vascular than the surrounding mucous membrane. There is no puckering or diminution in the calibre of the gut. In some cases, however, cicatrization does not take place so readily, and the floor of the ulcer becomes the seat of a *secondary* ulceration. This usually takes place after the general disease has run its course, or during a relapse. Profuse hemorrhage and perforation more commonly result from the secondary ulceration than from the primary sloughing of the glands.

The Mesenteric Glands.—The change in the mesenteric glands is probably secondary to that in the intestine. These glands become the seat of an acute cellular infiltration, and are enlarged, soft, and vascular. They usually, like many of the glands in the intestine and the spleen, undergo a gradual process of resolution. In rare cases, however, the capsule of the gland is destroyed, and the softened matters may escape into the peritoneal cavity and so cause peritonitis. The enlarged glands may also become caseous, and subsequently calcify.


CHAPTER XXXVI.

INFLAMMATION OF MUCOUS MEMBRANES.

IN mucous membranes inflammatory processes are divided into *catarrhal*, *fibrinous*, and *croupous* inflammations.

CATARRHAL INFLAMMATION.—Catarrhal inflammation, or as it is more commonly called *catarrh*, is much the more frequent. In its milder forms this is characterized mainly by an increased secretion of mucus. The initial hyperæmia of the membrane is followed by an abnormal production of epithelial and mucous elements. The liquid which transudes from the bloodvessels is more abundant, the small spherical cells which constitute the mucus-corpuscles are produced with increased rapidity, and the result is an increased secretion of mucus, rich in cellular elements. These young cells are most of them emigrant blood-corpuscles, others are produced within the epithelial cells by

FIG. 98.



Catarrhal Inflammation of the Conjunctiva.—*a.* Epithelium. *b.* Sub-epithelial connective tissue. Showing the proliferation of the epithelium, and the origin of the young elements within the epithelial cells. (Rindfleisch.)

endogenous multiplication. As the process continues, many of the epithelial elements become loosened and are discharged with the mucus, and within these groups of bodies may occasionally be seen, which are evidently young mucus-corpuscles. (Fig. 98.) The secretion of the mucous glands is also increased. Owing to these changes, the mucous membrane becomes swelled and abnormally vascular. The

increased vascularity is evidenced by redness during life, but after death the blood usually passes out of the vessels, and the membrane may look paler than natural.

If the inflammation be more severe, the vascular phenomena are more marked, the production of young elements is more rapid, they are smaller and not so well developed, the epithelium loosens and falls off more readily, and the secretion becomes puriform from the great number of cellular elements which it contains. Many of these elements are indistinguishable from pus-corpuscles, others are somewhat larger and resemble the corpuscles of normal mucus. Between the corpuscles of mucus and pus there is no line of demarcation, the one passing by insensible gradations into the other. The former are somewhat larger and more regular in shape than the latter, and usually contain only a single nucleus. As the process continues the sub-epithelial tissue is gradually involved, and becomes infiltrated with small cells. Owing to the loss of epithelium, the surface of the membrane may at the same time become more or less irregular, and present numerous irregular abrasions or ulcers.

These changes in the mucous membrane itself are accompanied by inflammation of the lymphatic structures which it contains. The lymph-follicles become enlarged. Their contents may soften and form a minute pseudo-abscess, and this bursting gives rise to a small ulcer. These are the follicular ulcers so often seen in catarrhal conditions of the intestines and pharynx. The ulceration in some cases extends beyond the confines of the follicle. The proper glandular structures may also become involved. Their epithelium multiplies, the glands become choked with the epithelial elements, and they may subsequently atrophy. This is seen in catarrh of the stomach.

The acute process may quickly subside, or it may become chronic. In the latter case the vascularity diminishes, but the multiplication of the epithelial elements continues, and the sub-epithelial tissue becomes more extensively infiltrated with small cells. The mucus thus becomes thicker and more puriform in character.

Chronic catarrhal inflammations of mucous membranes differ from the acute, inasmuch as the sub-epithelial connective tissue is much more extensively involved. This tissue becomes infiltrated with small elements, which may ultimately form a more or less completely fibrillated structure. The membrane thus becomes indurated and thickened, and the pressure exercised by the new tissue may induce atrophic changes in the glandular structures which it contains; by preventing

the exit of their secretion it may also cause them to dilate so as to form cysts. This atrophy of the proper glandular structures is seen in chronic catarrh of the stomach. These changes in the sub-epithelial connective tissue are usually accompanied by enlargement of the lymphatic structures, an enlargement which sometimes gives to the membrane a nodular or granular appearance. This is well seen in the pharynx (follicular pharyngitis). The enlarged lymphatic structures may also constitute the starting-points of an ulcerative process. (See "Tuberculosis of Mucous Membranes.") In some situations, as the stomach and intestine, the membrane often at the same time becomes deeply pigmented.

FIBRINOUS AND CROUPOUS INFLAMMATIONS.—The terms fibrinous and croupous, as applied to inflammations of mucous membranes, have in this country usually been regarded as synonymous, and have been employed to designate those inflammations which lead to the production of a so-called false membrane, such as is seen, for example, in true croup. According to Dr. E. Wagner, and some other continental observers, however, fibrinous and croupous inflammation must be regarded as constituting distinct varieties. Fibrinous inflammations are those intense forms of inflammation in which the exuded inflammatory liquids contain a large proportion of coagulable material, so that they form a kind of membranous layer. They occur as the result of severe injuries of the mucous membrane, such as are occasionally met with in children from attempts to drink out of a vessel containing boiling water. Croupous inflammations, on the other hand, according to these observers, are characterized mainly by a peculiar metamorphosis of the epithelial cells, which become enlarged, and are then converted into a kind of fibrinous material. They occur in croup and diphtheria. This distinction between fibrinous and croupous inflammation, however, has at present met with but partial acceptance in this country. The two appear to be closely allied; and the formation of the membranous layer is probably often due partly to the coagulable inflammatory liquids, and partly to mucoid, or to some similar metamorphosis of the epithelial elements.

Croup.

In croup, the inflammatory process at its commencement is simply a severe catarrhal one, consisting in an increase in the vascularity of the membrane, together with an exudation of liquor sanguinis and

blood-corpuscles, and proliferation of the epithelial elements. This is quickly followed by the production of a membranous layer upon the surface of the tissue, usually known as the *false membrane*. This membrane varies considerably in consistence, being in some cases firm and tough, in others soft and pultaceous. It can readily be removed from the subjacent tissues, and after its removal, the mucous membrane is left partially deprived of its epithelium. If the inflammatory process subsides, the epithelium is reproduced, and the part returns to its normal conditions; but if it continues, a fresh false membrane is formed. As the inflammation extends downwards into the trachea and bronchi, a gradual transition may usually be observed from croupous to catarrhal inflammation. In the upper part of the air passages, where the process is most intense, the false membrane exists, but in passing downwards, where it becomes less severe, the mucous membrane is simply coated with mucus.

Respecting the nature of the false membrane, as already stated, opinions differ. When examined microscopically, it is seen to consist of a finely fibrillated and delicate network of homogeneous material, inclosing in its meshes numerous leucocytes and epithelial cells. This fibrillated network has been commonly supposed to be produced by the coagulation and fibrillation of the inflammatory exudation liquids, but according to Wagner and others it is mainly due to changes in the epithelium. The smaller cells which are inclosed in its meshes are doubtless principally escaped blood-corpuscles; the larger epithelial elements, which are much less numerous, are those belonging to the mucous membrane.

Micrococci appear to be constantly present in the croupous membrane, and also in that of diphtheria. They are usually found as colonies of globular organisms in the upper layers of the membrane. Whether they are essential or merely accidental is at present uncertain.

Diphtheria.

It is impossible in most cases to draw any line of demarcation between the histological changes occurring in diphtheria and those of croup, and it must be borne in mind that by many these diseases are regarded as identical. In diphtheria, however, there is often some cellular infiltration of the submucous tissue, and the false membrane is less readily removed. The circulation also sometimes becomes so

much interfered with, that portions of the tissue lose their vitality, and large ash-colored sloughs are formed, which after removal leave a considerable loss of substance.

Dysentery.

The inflammatory processes occurring in the intestine in dysentery are for the most part limited to the large intestine, although the ileum is also occasionally involved. The inflammation is always most marked in the rectum and descending colon, and it may be stated generally, that it is characterized by the ulceration and sloughing of the membrane to which it gives rise.

The intestinal changes vary considerably, according to the intensity of the inflammatory process. In the milder forms of the disease the changes are most marked on the summits of the folds of the mucous membrane. These are found covered with a grayish-white layer of fibrinous-looking material, which, when scraped off, leaves a superficial loss of substance. The mucous membrane generally is hyperæmic and softened. The submucous tissue is also infiltrated with inflammatory products, and the solitary glands are enlarged and prominent.

When the process is more severe, the submucous tissue becomes more extensively involved, and the superficial layer of fibrinous material extends over wider areas and implicates more deeply the mucous membrane. The thickening of the intestinal wall, however, is much greater in some parts than in others, so that projections are produced upon the inner surface of the intestine, corresponding with those parts which are the most affected. The enlarged solitary glands usually slough, and so give rise to circular ulcers, which rapidly increase. When the process has reached this stage, the muscular and serous coats are implicated, the latter being covered with layers of fibrin which form adhesions with adjacent parts. The intestine is much dilated, and contains blood and disintegrating inflammatory products.

In the most intense forms of the disease, the necrosis is more extensive. According to Rokitansky, large portions of the mucous membrane are converted into black rotten sloughs. The submucous tissue is infiltrated with dark blood and serum, but subsequently it becomes the seat of a reactive suppurative inflammation, by means of which the necrosed portions of tissue are removed.

If death does not occur and the inflammatory process subsides, the

ulcers may gradually heal. When the loss of substance has not been considerable, the edges of the ulcers may, by the contraction of the submucous tissue, become completely approximated. More commonly, however, the loss of substance is so great that portions of the membrane are left, consisting simply of connective tissue.

When the inflammatory process becomes chronic, the changes in the submucous connective tissue become more marked, and the new fibroid growth gives rise to considerable thickening and induration of the intestinal wall, and to more or less contraction and narrowing of the cavity. Sometimes it forms fibrous bands which project into the gut. The formation of abscesses and fistulous passages occasionally occurs in the thickened intestinal wall.

CHAPTER XXXVII.


INFLAMMATION OF SEROUS MEMBRANES.

INFLAMMATORY processes in serous membranes vary in their intensity, and in the amount and character of the effusion.

The process commences, as in mucous membranes, with hyperæmia, exudation of liquor sanguinis and emigration of blood-corpuscles,

together with increased activity of the endothelial elements. The endothelial cells enlarge and become more granular, their nuclei divide, and new elements are formed by endogenous multiplication. (Fig. 99.) The number of new cells which are thus produced, together with some of the older cells, and a large number of emigrant blood-corpuscles, escape with the exuded liquor sanguinis into the serous cavity, where they may probably

FIG. 99.




Inflamed Epiploon of a Rabbit.—Showing the endogenous proliferation of the endothelium. $\times 250$. (Cornil and Ranvier.)

continue to exhibit their formative activity.

Owing to these changes the membrane loses its natural smooth and

glistening appearance, and becomes opaque, roughened, and exceedingly vascular. Its surface at the same time becomes covered with a fibrinous layer, and more or less liquid transudes into its cavity. The coagulable material which exudes from the vessels forms a soft, elastic, membranous or reticulated investment, inclosing in its meshes numerous small cells. This either glues the two surfaces of the membrane together, or, if they are separated by liquid effusion, forms a slightly adherent layer. (Fig. 100.) The exuded liquid varies considerably in amount and is always turbid, thus differing from non-inflammatory effusions. It contains flakes and masses of coagulated fibrin and innumerable cell-structures, the latter being in the earliest stages of the process almost entirely emigrants.

FIG. 100.



Inflammation of the Diaphragmatic Pleura.—Showing the adherent fibrinous layer. *a.* Muscular coat of diaphragm. *b.* Sub-serous tissue. *c.* Serous membrane. *e.* Fibrinous layer. $\times 400$. (Rindfleisch.)

The nature of the subsequent changes will depend upon the intensity of the inflammation, and upon the amount of liquid exuded into the serous cavity. If the inflammatory process subsides, and the liquid exuded is not sufficient to prevent the two surfaces of the membrane from coming into contact, they grow together and form an adhesion. This constitutes the so-called *adhesive inflammation*. The union is effected by the formation of connective tissue. The small round cells imbedded in the fibrinous layer become elongated and spindle-shaped, the fibrinous material fibrillates, and numerous new vessels are formed. Many of the latter atrophy and disappear as the organization becomes complete. The process is thus precisely similar to that which takes place in the union of an incised wound. It is probable also that in some cases union may take place without the intervention of any fibrinous layer, by the formation and growing together of irregular papillary outgrowths from the subendothelial tissue.

If, however, the inflammatory process is severe, or the surfaces of the membrane are separated by a large quantity of liquid effusion, organization and adhesion cannot be thus readily effected. If a large quantity of liquid exists in the serous cavity, the removal of this becomes necessary before union can take place. If the inflammatory process continues, or its severity is great, union is prevented by the formation of pus. These two conditions must be considered separately.

The existence of a large amount of effusion interferes with the adhesion of the serous surfaces, and before this can be effected the absorption of the liquid becomes necessary. In some cases this occurs very rapidly, and it is probably greatly facilitated by the new blood-vessels which are produced in the fibrinous layer. Frequently, however, the process is more prolonged, and the subendothelial connective tissue becomes involved before absorption takes place. When this is the case the connective tissue of the serous membrane becomes infiltrated with small cells, which form a granulation-tissue beneath the layer of proliferating endothelium. The endothelium itself gradually becomes less abundant, and the fibrinous exudation undergoes fatty changes and liquefies. The new granulation-tissue becomes exceedingly vascular from the formation of numerous bloodvessels, and if the inflammation subsides, it gradually develops into connective tissue, and thus a false membrane is formed, rich in vessels, which takes the place of the endothelial layer. As the liquid is absorbed, the two surfaces of the membrane come into contact and grow together, the new vessels becoming gradually obliterated.

If the inflammatory process does not subside, or from its commencement is of considerable severity, it may be attended by the formation of large quantities of pus. In this case the emigration of blood-corpuscles is so considerable that the young elements exist in large enough numbers to give to the exuded liquids a purulent character. The condition is then termed empyema. As the connective tissue becomes involved and a granulation-tissue is formed, this may continue to generate pus like an ordinary granulating wound. If the pus be renewed, the suppuration may gradually cease, the granulation-tissue develop into a fibrous structure, and the union of the serous surfaces thus be effected. The serous membrane becomes greatly thickened, and the new tissue undergoes considerable contraction in the process of its organization.

CHAPTER XXXVIII.

INFLAMMATION OF THE LIVER.

INFLAMMATORY processes in the liver are either acute or chronic. *Acute* inflammations, leading to suppuration, are usually infective in their nature, resulting from the transmission of infective materials from inflammatory lesions in the abdominal organs or in other parts. The processes are consequently most frequently disseminated and confined to small portions of the hepatic substance. The pus-corpuscles—which usually accumulate so as to form an abscess—are almost entirely emigrants, although recent investigations render it highly probable that they may also originate by the endogenous proliferation of the liver-cells.

These suppurative inflammations are in the majority of cases pyæmic, and when pyæmic they are usually multiple. Hepatic abscess is also not infrequently, in tropical climates, associated with dysentery. Here the abscess is commonly single, and it probably results from the absorption of infective substance from the diseased intestine. As other causes of hepatic abscess must be mentioned, inflammation of the bile-ducts, such as sometimes results from gall-stones, etc., and external violence.

Chronic inflammatory processes in the liver, being of much less intensity, lead to a gradual increase in the connective tissue of the organ, and consequently to induration and to the subsequent atrophy of its proper structure. They constitute what is known as *interstitial hepatitis*, or more commonly as *cirrhosis*.

Cirrhosis of the Liver.

Cirrhosis of the liver, or interstitial hepatitis, is characterized by a gradual increase in the connective tissue of the organ and by the subsequent atrophy of the liver cells, so that when examined with a low magnifying power, the acini are seen to be separated by broad tracts of new interstitial growth. (Fig. 101.)

The process usually commences in the connective tissue surrounding the smaller branches of the portal vein, and gradually extends to that

FIG. 101.




Cirrhosis of the Liver.—Showing the growth of connective tissue between the hepatic lobules. a. Lobules. b. New growth of interlobular connective tissue. $\times 16$.

surrounding the larger ones, until ultimately the connective tissue throughout the whole organ may become involved. This tissue becomes infiltrated with small round cells, from which are produced a more or less completely fibrillated structure, containing in the earlier stages of its development numerous new bloodvessels, which are supplied by branches of the hepatic artery. The change is sometimes uniform, but more commonly it is much more advanced in some parts than in others. In many places the new tissue consists almost en-

tirely of cells, whilst in others the fibrillation is much more marked; and if the process lasts long enough, the vascular and richly-cellular growth gradually assumes the characters of cicatricial tissue, the vessels at the same time becoming obliterated. (See Fig. 84.)

FIG. 102.



Cirrhosis of the Liver.—A thin section from the external portion of one of the hepatic lobules. Showing the new growth of connective tissue, and the way in which it involves the intercellular network and causes atrophy of the liver-cells. $\times 200$.

The effect of the new growth is ultimately to cause atrophy of the hepatic cells, and to obstruct the circulation through the portal capillaries and the passage of bile through the bile-ducts. This effect is

materially increased by the process of contraction which the new tissue undergoes. The hepatic cells in the outer zone of the lobules are the first to atrophy; the new tissue insinuating itself between them, so as to gradually involve the intercellular network. (Fig. 102.) The cells then become smaller and undergo fatty metamorphosis; and ultimately they are completely destroyed. Those in the central parts of the lobule are in the earlier stages but little altered, although they are often stained with bile. As the growth extends, however, these also become annihilated, and the whole lobule may be replaced by connective tissue. The cells in the outer part of the lobules are sometimes infiltrated with fat prior to their destruction, the cirrhosis being associated with fatty infiltration. (See Fig. 8.)

The obliteration of the portal capillaries by the new tissue gives rise to ascites, hæmatemesis, diarrhœa, enlargement of the spleen, and to the other results of portal congestion. Obstruction of the bile-ducts, although it may cause staining of the hepatic substance, is rarely such as to interfere with the passage of the bile into the intestine, and hence there is usually but little general jaundice.

Physical Characters.—The physical characters of the cirrhotic liver vary. In the earlier stages of the disease the organ is probably always more or less increased in size; the enlargement being almost uniform, and the edge rounded and thickened. In some cases this enlargement is considerable, and there is but little subsequent atrophy and contraction, the organ being found large and comparatively smooth after death. More commonly, however, atrophy and contraction constitute marked features in the disease, and the liver is small, presenting an irregular nodular surface.

Respecting the causes of these differences in the alterations which the liver undergoes as a consequence of the interlobular growth—we are at present unable to speak certainly. They probably, however, depend in great measure upon differences in the age of the growth, and in its distribution. In the large livers the growth is usually richly cellular and very generally distributed; whilst in the smaller nodular forms of cirrhosis it is commonly more fibrous and less cellular in character, and its distribution is more partial. It is probable that in the larger livers the growth is in an earlier stage of development, and that if the patient had lived, the organ would have diminished in size. When the growth is irregularly distributed it necessarily causes more irregularity in the configuration of the liver than when the distribution is more uniform.

The cirrhotic liver is always increased in consistence. It breaks down less readily under the finger, and is often exceedingly firm and tough. On section, the new tissue is visible to the naked eye surrounding the lobules and in many parts completely replacing them. This gives to the cut-surface a mottled granular appearance, the lobules themselves contrasting with the new interlobular tissue; and this appearance is sometimes increased by fatty infiltration of the cells in the peripheral zone. The capsule also may be thickened, and the organ is frequently stained with bile.

Acute Yellow Atrophy.

This exceedingly rare disease of the liver is characterized by a rapid diminution in its size, and by destruction of the hepatic cells. The organ may, in the course of a few days, be reduced to less than half its natural bulk, being especially diminished in thickness. It is soft and flabby in consistence, bloodless, and of a dull yellow or yellowish-red color. The lobules are indistinguishable. When examined microscopically, the liver-cells are found to be completely destroyed, being replaced by granular debris, fat granules, and pigment. Tyrosin and leucin have been found in the disintegrated liver-tissue.

The pathology of this disease is exceedingly obscure. By some it has been regarded as an acute inflammatory process, by others as a passive degeneration.

CHAPTER XXXIX.

INFLAMMATION OF THE KIDNEY.

INFLAMMATORY processes in the kidney present certain variations according to their intensity. They comprise *suppurative*, *tubal*, and *interstitial* nephritis. Of these, suppurative nephritis, as the name implies, is an intense inflammation leading to the formation of abscess. This usually results from the transmission of infective materials from some primary lesion (pyæmic), or it is associated with some inflammatory condition of the urinary passages. Tubal nephritis is also an

inflammation of considerable intensity, and in it the structural changes have their principal seat in the tubular epithelium. Interstitial nephritis is an inflammatory process which runs a more chronic course, and is of less intensity than either of the preceding. Consequently in it the principal structural changes take place in the connective tissue around the bloodvessels—in the intertubular connective tissue. (See “Chronic Inflammations.”) It must, however, be distinctly borne in mind that these two varieties of histological changes—those in the tubular epithelium and those in the intertubular connective tissue—are very constantly associated. Tubal and interstitial nephritis cannot therefore be separated from one another by any distinct line of demarcation. They might, perhaps, be more correctly designated *acute* and *chronic* nephritis.

Suppurative Nephritis.

Acute inflammatory processes in the kidney attended by the formation of pus, give rise to *renal abscesses*. Such processes, as already stated, are often infective in their nature, resulting from the transmission of infective materials from some primary inflammatory lesion, as in pyæmia; and they also occur in connection with inflammatory conditions of the urinary passages. In the latter they constitute what is commonly known as the “Surgical Kidney.”

The abscesses met with in the kidney as the result of pyæmia are principally confined to the cortex, and they resemble pyæmic abscesses in other organs. They are usually multiple, and are often surrounded by a thin zone of red hyperæmic tissue. Their characters will be more fully described in the chapter on embolism. (See “Results of Embolism.”)


SURGICAL KIDNEY.—This is the name commonly given to those inflammatory conditions of the kidney which result from inflammation of the urinary passages. These renal inflammations frequently occur in association with inflammation of the pelvis of the kidney, such as is produced by a renal calculus; or with those inflammatory changes in the mucous membrane of the urinary tract which result from vesical calculus, urethral stricture, or enlargement of the prostate. In addition to the inflammation of the urinary passage there almost invariably exists some impediment to the escape of the urine.

The changes which occur in the kidney vary. When the abnormal condition of the urinary passages has lasted for a lengthened period, it

is exceedingly liable to give rise to chronic renal changes, which are characterized mainly by more or less cellular infiltration of the inter-tubular connective tissue. ("Interstitial Nephritis.") This cellular infiltration occurs both in the pyramids and cortex. The tubules are in some parts found blocked with epithelium, whilst in others they are wasted or obliterated. The walls of the small arteries are not thickened. Owing to these changes, the kidneys—although at first they may be enlarged—gradually become diminished in size, the atrophy being probably, in most cases, partly due to the pressure exercised by the retained urinary secretion. The capsule becomes adherent and thickened, and when removed, the surface of the organ is found puckered and irregular.

In other cases the process is much more acute, the cellular infiltration of the intertubular tissue is much more abundant, and the young cells (emigrants) accumulate in certain situations in such numbers as to give rise to abscesses. (Fig. 103.) The blocking of the tubes

FIG. 103.



Surgical Kidney.—At the lower part of the figure is seen the cellular infiltration of the inter-tubular tissue, and the blocking of the tubes with epithelium and leucocytes. At the upper part, there is the commencing formation of an abscess. $\times 100$.

with epithelium is then more marked, and many of them are found filled with the emigrant leucocytes. The kidneys in such cases are usually enlarged, pale, and soft; and yellowish-white patches of various sizes are seen on their cut-surface. Many of these patches have purulent centres. They are not commonly surrounded by a zone

of red hyperæmic tissue, as in pyæmia. Yellowish lines and striæ are also often seen extending from the papillæ to the bases of the pyramids. This acute suppurative process may occur independently of the more chronic lesions. Much more commonly, however, it supervenes in kidneys which have already undergone more diffuse interstitial changes, and its supervention appears not unfrequently to be determined by the performance of some surgical operation in connection with the bladder or urethra.

Respecting the pathology of these changes in the kidney, they undoubtedly result from the inflammatory condition of the urinary passages, and the impediment to the escape of the urine with which they are associated. As to the manner in which these conditions produce the kidney changes we are unable to speak certainly. The simple tension due to the urinary impediment is in itself sufficient to cause interstitial changes. Dr. Dickinson believes that in many cases the changes are due to the absorption of septic substances contained in the urine.¹ According to Mr. Marcus Beck, reflex irritation of the kidney caused by injuries of the bladder from calculus, operative interference, etc., is often an important causative element.

Tubal Nephritis.

Tubal, parenchymatous, or acute nephritis is that subacute inflammation of the kidney which constitutes the more acute form of Bright's disease, that form which is characterized by a more or less marked beginning, scanty and highly albuminous urine, and dropsy. In its more advanced stages it is the large kidney of chronic Bright's disease.

The changes which take place in the kidney have their seat mainly in the cortex. They comprise increased vascularity and exudation into the urine tubes, and swelling with, probably, proliferation of the tubular epithelium. The prominence of the vascular phenomena, however, varies very considerably in different cases.


In the most acute cases of Bright's disease—those which are induced suddenly from exposure to cold, the vascular changes are marked. In these cases the contraction of the cutaneous vessels and the check to the function of the skin caused by the chilling of the surface, lead to considerable hyperæmia of the organs. There is

¹ Dickinson on "Disseminated Suppuration of the Kidney." *Med. Chir. Trans.*, vol. lvi.

abundant exudation into the urine-tubes, many of the capillaries at the same time frequently rupture, and thus there is an escape of blood-corpuscles and of liquor sanguinis into the tubes of the cortex; hence the blood and "blood-casts" in the urine which are so characteristic of the early stages of these most acute forms of the disease. In this stage the process may quickly subside, and, with the exception of some swelling and desquamation of the tubular epithelium, no further alterations take place in the kidney.

In the less acute cases the vascular phenomena are much less marked, and the principal changes take place in the tubular epithelium. The epithelial elements become swollen and granular. (Fig. 104.) The granules, which are often so numerous as to occlude the

FIG. 104.



Tubal Nephritis.—The earlier stage of the process. Showing the swelling of the tubular epithelium. In some of the tubes the epithelium has fallen out during the preparation of the section. $\times 230$.

nucleus of the cell, are soluble in acetic acid, and thus differ from molecular fat. This is the condition known as "cloudy swelling." Many small cells are also seen within the tubes, and these have been supposed to be the products of epithelial proliferation. It is probable that some of them are thus produced, although the majority must be regarded as having escaped from the vessels. Owing to these changes the tubes become distended with cellular elements. (Fig. 105.)

In addition to the cell-forms, many of the tubes also contain hyaline cylinders, which are commonly regarded as consisting of coagulated

substances which have escaped from the vessels. By many pathologists, however, this hyaline material is supposed to be the product of a mucoid, or some allied, metamorphosis of the epithelium. The cell-forms contained within the tubes adhere to this hyaline substance, and some of them are washed away and appear in the urine as "epithelial casts." A varying number of emigrant leucocytes are also usually found around the Malpighian tufts.

The alterations which these changes produce in the physical characters of the kidneys vary according to the extent of the hyperæmia. The organs are always considerably increased in size, and more or less abnormally vascular. The capsule separates readily, exposing a perfectly smooth but vascular surface. The consistence is diminished, the tissue breaking with a soft, friable fracture. On section, the in-

crease in the size of the organ is seen to be principally due to the increased thickness of the cortex. This is either of a reddish-brown, or of an opaque-white or pale buff color; these differences depending upon the relative proportion of blood and of accumulated intra-tubular elements. Although in the earliest stage of the most acute forms of the disease the color is redder than natural, it usually soon becomes pale and opaque. This is owing to the swelling of the epithelial elements and to the accumulation in the cortical tubes. The blood becomes expressed from the intertubular vessels, and hence the increased vascularity is most evident in the Malpighian corpuscles, beneath the capsule, and in the pyramidal portion of the organ. The Malpighian corpuscles stand out as prominent red points, and the pyramidal cones are of a deep red color, thus contrasting strongly with the pale opaque cortex.

The termination of the process varies. The increased vascularity and epithelial change may, as already stated, subside, and the inflammatory products passing away in the urine, the organ gradually return to its normal condition. In other cases the disease continues; and although the vascularity diminishes, the vitality of the epithelial elements becomes so much impaired that they undergo retrogressive changes. The cells then continue to come away with the urine, adhe-

FIG. 105.



Tubal Nephritis—a Single Urine Tube.—Showing the accumulation within the tube. In the few epithelial cells which have escaped is seen the granular condition of the protoplasm. $\times 200$.

rent to the casts, but instead of presenting the swelled granular appearance as in the earlier stage of the disease, they contain molecular fat. This fat gradually increases in amount as the degeneration proceeds, until ultimately the cells are destroyed, and it appears as free molecules and granules on the tube-casts.

This fatty degeneration of the epithelium is attended by corresponding changes in the appearance of the organ. The redness diminishes and the Malpighian corpuscles are less prominent. The enlarged cortex presents a uniform yellowish-white tinge, studded with minute yellowish streaks. This is owing to the presence of fat in the tubes of the cortex. This fatty stage, if only slightly advanced, may undoubtedly pass off. The degenerated cells are carried away by the urine, from those which remain in the tubes the fat is probably partially absorbed, the retrograde process gradually ceases, and the organ returns to nearly its normal size and condition. In other cases the degeneration continues, and owing to the loss of epithelium the kidney becomes somewhat diminished in size. This atrophy, however, I believe never occurs without changes in the intertubular connective tissue.

When the inflammatory process is of longer duration, or when the kidneys are the seats of repeated attacks of sub-acute inflammation, the intertubular connective tissue invariably becomes involved. This tissue becomes infiltrated with small cells which ultimately tend to form a fibrillated structure. (See Fig. 106.) The new intertubular growth may gradually increase, and so lead to more or less irregular atrophy of the organ, such as will be described as occurring in interstitial nephritis. (See "Interstitial Nephritis.") In other cases death ensues before any marked atrophy has taken place, and thus the organ may remain smooth and large to the termination of the disease. The intertubular growth is sometimes found thickly studded with fatty granules.

SCARLATINAL NEPHRITIS.—The changes which take place in the kidney in scarlatina have usually been regarded as precisely similar to those which have been just described as tubal nephritis. Recent investigations, however, show that this view requires considerable modification. It has long been known that cases of scarlatina sometimes occur in which the kidney change differs from the type of ordinary acute nephritis; but it is only during the past year, owing to the researches of Dr. Klein, that any exact knowledge of the scarlatinal

kidney has existed.¹ The changes as described by Dr. Klein may be thus briefly summarized :—

The earliest changes, those occurring during the first week of the disease, comprise :—

1. Increase of the nuclei covering the glomeruli of the Malpighian corpuscles.

2. Hyaline degeneration of the elastic intima of minute arteries, especially of the afferent arterioles of the Malpighian corpuscles. This change produces a swelling of the intima, so as in some places to cause a distinct narrowing of the lumen of the vessel. The capillaries of the Malpighian corpuscles are in parts altered in the same way, in consequence of which many of them become impermeable.

These marked and early changes in the Malpighian corpuscles are interesting, as helping to explain those cases which are occasionally met with, in which death occurs from anuria and uræmia, and no catarrhal or other conspicuous alterations are found in the kidneys.

3. Multiplication of the nuclei of the muscular coat of the minute arteries, and a corresponding increase in the thickness of the walls of these vessels.

4. Cloudy swelling of the epithelium in the convoluted tubes, with multiplication of the epithelial nuclei. Granular matter and even blood may also be found in the tubes, and in the cavity of Bowman's capsules. These parenchymatous changes are in the early stages of the disease but little marked.

The later changes, those occurring after the first week, consist in :—

5. A cellular infiltration of the intertubular connective tissue of the cortex (interstitial nephritis), together with an increase in the epithelial changes, and a crowding of the tubes with small round cells (leucocytes). The cellular infiltration commences around the larger vascular trunks, whence it spreads rapidly into the bases of the pyramids, and especially into the cortex. As it increases, the epithelium undergoes fatty degeneration, and the urine-tubes gradually become obliterated.

Interstitial Nephritis.


Interstitial or chronic nephritis, or, as it is sometimes called, granular degeneration of the kidney, is that more chronic change which

¹ "The Anatomical Changes of the Kidney and other Organs in Scarlatina of Man," by Dr. Klein.—*Trans. Path. Soc. Lond.*, 1877. Vol. xxviii.

occurs in the most chronic form of Bright's disease, chronic Bright's disease with contracted kidney. The process, like the corresponding one in the liver, is characterized by a gradual increase of the connective tissue of the organ, and by atrophy of the tubal structures. Sometimes, as has been seen, this is merely an advanced stage of tubal nephritis. More commonly, however, the process appears to be very chronic from its commencement, and not to be preceded by any marked vascular phenomena or by any active changes in the tubular epithelium.

The first change in the kidney appears to consist in a cellular infiltration of the intertubular connective tissue. In the earliest stages of the process this tissue becomes infiltrated with small round cells. (Fig. 106.) If the growth is rapid, these may be exceedingly nume-

FIG. 106.




Interstitial Nephritis.—The earlier stage of the process. Showing the cellular infiltration of the intertubular connective tissue. The epithelium has fallen out of some of the tubes during the preparation of the section. $\times 200$.

rous, and the intercellular substance soft and amorphous. Usually, however, the process is much slower and more insidious, the cells are not numerous, and the new tissue is more or less developed into a fibrillated structure. This is the condition in which it is usually met with. These changes are almost entirely limited to the cortical portion of the kidney, and although here they are more or less general, the new growth is more abundant in some parts than in others, being usually most so around the Malpighian bodies and in the neighborhood of the capsule, with which it is closely united. In this stage the tubes and their epithelium are unaffected.

In this early stage, the kidney may be somewhat increased in size, the capsule usually separates less readily than in health, and the surface of the organ is slightly granular. On section, the cortical substance is in some cases paler, in others redder than natural. The cut surface also looks obscurely granular. The Malpighian bodies stand out as red points, and the bases of the pyramids and surface of the organ are sometimes hyperæmic. The consistence of the kidney is usually slightly denser and tougher than natural. This, however, will vary with the character of the new growth, as will also the increase in size and the irregularity of the surface. If the new tissue is slowly developed, the size will be but little increased, whereas the increase in consistence and the granular condition will be more marked. If, on the other hand, the process be more rapid, and the cellular infiltration more abundant, there will be a greater increase in the size, the granular character will be slight, and the consistence may be even softer than natural.

The second stage in the process is characterized by the atrophy of the tubular structures. This is probably mainly owing to the pressure exercised by the intertubular growth, and to the cicatricial contraction

FIG. 107.




Interstitial Nephritis.—An advanced stage of the process. Showing the intertubular tissue with the granular and fatty debris which result from the degeneration. $\times 100$.

which it often undergoes. The atrophy consequently is not uniform, but is more marked in some parts than in others. The tubes are now found in many parts diminished in size, or completely obliterated; whilst in others they are irregularly dilated, and filled with degene-

rated epithelial products. Their walls are usually thickened. As the atrophy proceeds the intertubular tissue thus becomes mingled with the granular and fatty débris which results from the retrograde process. (Fig. 107.) The Malpighian bodies become approximated, and the secreting structure throughout large tracts of the kidney is destroyed. (Fig. 108.) The irregular pressure exercised by the

FIG. 108.



Interstitial Nephritis.—A very advanced stage of the process. Showing the large amount of tissue between the tubes of the cortex, and the extensive atrophy of the tubes. The degenerated epithelium which was contained in some of the tubes has fallen out in the preparation of the section. $\times 50$.


new growth also gives rise to the formation of cysts. These originate partly in the Malpighian capsules, and partly in the urine tubes—the latter becoming irregularly dilated.

The small arteries of the kidney also undergo important alterations. These were first described by Dr. Johnson. Dr. Johnson states that the walls of these vessels are thickened, owing to hypertrophy of their longitudinal and circular muscular fibres. This change is well represented in the accompanying drawing. (Fig. 109.) Both the internal longitudinal and the external circular muscular fibres are considerably increased. The external fibrous coat of the vessel is also thickened, and it appears to be continuous with the new intertubular tissue. This thickening of the external coat has been especially insisted upon by Sir W. Gull and Dr. Sutton. I have usually found it associated with the muscular hypertrophy, although the latter is undoubtedly the most prominent structural change.

In this more advanced stage of the disease the kidney is diminished

in size. Its surface is more granular, the capsule more thickened and adherent, and it cannot be removed without tearing the kidney substance. The superficial vessels are seen unduly marked in the depressions between the granulations. The cortex is tough and fibrous, of a

FIG. 109.



Arteries from Contracted Kidney of advanced Chronic Bright's Disease.—*a.* Longitudinal section, showing the great thickening of the internal longitudinal and external circular muscular coat, also of the outer fibrous coat. *b.* Transverse section of another vessel less diseased. Here is seen the thickening of the circular muscular and external fibrous coat. $\times 200$.

reddish, yellowish gray, or buff color, mottled with yellow streaks and patches; and usually numerous small cysts are distributed throughout it. Calcareous deposits are also often seen as white streaks between the tubes of the pyramids.

CHAPTER XL.

INFLAMMATION OF THE BRAIN AND SPINAL CORD.

INFLAMMATORY processes in the nervous centres are probably much less frequent than was formerly supposed. Many of those morbid changes in the brain and spinal cord which are attended by softening, and which were at one time regarded as the result of inflammation, are now known to owe their origin to simple interference with the vascular supply, such as results from thrombosis, embolism, or degenerative changes in the walls of the bloodvessels. (See "Fatty Degeneration of the Brain.")

The injury which determines the occurrence of acute inflammations in the brain or cord is most frequently some external violence—a blow, simple concussion, or fracture of the osseous framework. In other cases it is diseased bone, as in the inflammation of the brain which so often results from disease of the petrous portion of the temporal bone. Lastly, in a comparatively few number of cases, these inflammations are pyæmic.

The inflammatory process is almost invariably limited to small portions of the cerebral or spinal substance. The earliest change consists in a more or less intense and localized hyperæmia, which is frequently attended by rupture and minute extravasations of blood. The nervous tissue then becomes infiltrated with young cells and considerably softened, and it presents a uniform red or mottled color. This red softened tissue gradually acquires a brownish or brownish-yellow color, owing to changes in the hæmoglobin. The nerve-fibres become disintegrated, and the nerve-cells and cells of the neuroglia undergo fatty degeneration, thus forming the so-called “inflammatory,” or “exudation corpuscles.” (See Fig. 11.)

In many cases the accumulation of young cells is sufficient to give rise to the formation of an abscess, and a yellowish or reddish purulent liquid gradually takes the place of the original softened mass. The tissue surrounding the abscess is also hyperæmic, softened, and infiltrated with cells. The cellular infiltration may gradually extend, and thus the abscess increase in size until it opens either externally or into the ventricles. In other cases the abscess becomes limited and encapsuled by the formation of connective tissue from the neuroglia, and this tissue often forms a delicate network traversing the cavity. When the products of these acute inflammations have thus become encapsuled they may gradually dry up into caseous or calcareous masses, or the absorption may be more complete, so as to leave little more than a cicatrix. Respecting the source from which the young cells are derived—they are probably entirely emigrants. The nerve-cells themselves, and the cells of the neuroglia, appear to undergo no active changes in inflammation.

Suppurative inflammations are much more common in the brain than in the spinal cord. In the latter the inflammatory process is rarely of sufficient intensity to produce abscess; it causes merely softening of the nervous tissue, and, when not secondary to a meningitis, usually involves primarily and principally the central gray matter (myelitis).

INFLAMMATORY SOFTENING OF THE BRAIN AND CORD.—Conditions of softening of the cerebral or spinal substance resulting from inflammation, other than those which have been above described, probably rarely occur as *primary* lesions. Most varieties of softening which were formerly described as inflammatory—either from the red color of the softened tissue or from the acuteness of the process—result, as already stated, from thrombosis or embolism (see “Embolism in the Brain”), or are simply passive degenerative changes in which more or less extravasation of blood has taken place into the softened tissue. (See “Fatty Degeneration of the Brain.”) Inflammation and consequent softening of the nervous tissue, however, sometimes occurs as a *secondary* process. It takes place especially around clots of blood or other morbid products within the brain or spinal cord, and results from the injurious influence which these substances exercise upon the immediately adjacent structures. Such a result is not infrequent in cases of cerebral hemorrhage. The nerve-tissue immediately surrounding the clot becomes the seat of an inflammatory process, and it is found after death softened, hyperæmic, and infiltrated with young cells. Inflammation of the superficial portions of the cerebral and spinal substance may also occur as the result of meningitis.


Sclerosis of the Brain and Spinal Cord.

The term “sclerosis” is applied to certain changes in the nervous centres which are characterized by an increase of the connective tissue (neuroglia), and by atrophy and degeneration of the proper nervous elements. The affected portions are, for the most part, increased in consistence; sometimes, however, they are softer than natural. They are usually of a grayish color; hence the change has been called “gray degeneration.” The gray color appears to depend upon the loss of the white sheath of the nerve-fibres.

Respecting the nature of the change—it is probably in many cases the result of a chronic inflammatory process, the new growth of connective tissue leading to the destruction of the nerve-elements. In other cases, however, the atrophy of the nerve-tissue appears to precede the interstitial increase, so that the process would rather be regarded as a passive degeneration. This is probably the case in the secondary degenerations of the cord.

When the diseased portions of the brain and cord are examined microscopically, the cut ends of the nerve-fibres are seen in transverse

FIG. 110.



Sclerosis of Spinal Cord.—
From a case of Progressive Mus-
cular Atrophy. A transverse sec-
tion. Showing the new tissue
between the cut ends of the nerve-
fibres. $\times 200$.

section to be separated by a granular, finely nucleated, and often partially fibrillated tissue, which evidently originates from the neuroglia. (Fig. 110.) The nerve-fibres themselves are atrophied, and more or less fatty debris is found in the altered tissue. In many cases the nucleated character of the interstitial growth is wanting. The walls of the bloodvessels are also stated to be thickened, and their nuclei to be increased.

Sclerosis is more common in the spinal cord than in the brain. In the latter it occurs most frequently as small patches of induration limited to certain portions of the brain. These must be distinguished from the cicatrices left after the absorption of blood or of inflammatory products. Sclerosis of the cord may in the same way be localized and more or less disseminated; more commonly, however, it implicates only particular nervous tracts. It occurs thus in the posterior columns of the cord in locomotor ataxy, and in the anterior cornua in progressive muscular atrophy, etc. In the localized and disseminated forms of sclerosis the cellular character of the new growth is much more marked than it is in the simple uniform sclerosis which involves particular nerve-tracts.

CHAPTER XLI.

INFLAMMATION OF THE LUNGS.

IN the lungs, inflammatory processes comprise the three following principal varieties:—*Croupous*, *broncho-* or *catarrhal*, and *chronic* or *interstitial* pneumonia. Of these, the former occurs as an independent affection, whereas the two latter are usually the result of some antecedent bronchial or pulmonary inflammation.

Croupous Pneumonia.

Croupous, *exudative*, or *lobar* pneumonia, is that form of pulmonary inflammation which is met with in the disease known as acute primary

pneumonia. This is now regarded as a *general* disease of which the pneumonic consolidation is the prominent local lesion.

The local process is characterized by intense inflammatory hyperæmia of the lung, and by the exudation of a large amount of coagulable material into the pulmonary tissue. It is termed “croupous” by the Germans, from the supposed resemblance of the histological process to that of croup. The term “lobar” is applied to it because it almost invariably affects an extensive portion of the lung. The process is commonly described as consisting of three stages—1st, that of *engorgement*; 2d, that of *red hepatization*; and 3d, that of *gray hepatization*.

FIG. 141.



Croupous Pneumonia—Red Hepatization.—Showing the fibrinous coagulum in one of the pulmonary alveoli, inclosing within its meshes numerous leucocytes, which are already commencing to undergo fatty metamorphosis. A few leucocytes are also seen on the alveolar walls, and the alveolar epithelium is swollen and granular. $\times 200$.

In the *first* stage, that of *engorgement*, the lung becomes exceedingly vascular, the changes in the bloodvessels and circulation being such


as have been already described as characteristic of inflammation. The organ is of a dark red color, its specific gravity and absolute weight are increased, its elasticity is diminished, it is less crepitant and more friable than natural, and pits upon pressure. Its cut surface yields a reddish, frothy, tenacious liquid.

In the *second* stage, that of *red hepatization*, there is an exudation of liquor sanguinis and migration of blood-corpuscles into the pulmonary tissue. Some of the vessels may also rupture, and thus small extravasations occur. The exuded liquids coagulate within the air-vesicles and terminal bronchioles, the coagulum inclosing numerous white and some red blood-corpuscles. (Fig. 111.) It is stated by some German pathologists that the coagulum is in part produced by certain changes in the epithelium like those believed to occur in croup. (See "Croupous Inflammation of Mucous Membranes.") The lung is now much heavier than in the preceding stage, and is increased in size, so as to be often marked by the ribs. It is quite solid, sinks in water, and cannot be artificially inflated. It does not crepitate under the fingers, but is remarkably friable, breaking down readily with a soft granular fracture. The cut surface has a markedly granular appearance, seen especially when the tissue is torn. This is owing to the plugs of coagulated exudation matter which fill the alveoli. The color is of a dark reddish-brown, often here and there passing into gray. This admixture with gray sometimes gives a marbled appearance. Throughout this stage there appears to be but little alteration either in the alveolar walls or in the alveolar epithelium. On the former are often seen a few leucocytes, and the latter is usually swollen and granular. (Fig. 111.) The pleura covering the solid lung always participates more or less in the inflammatory process. It is opaque, hyperæmic, and coated with lymph.

The *third* stage, that of *gray hepatization*, is characterized by a continuance in the emigration of leucocytes, and by cell-proliferation. The white blood-corpuscles continue to escape from the vessels, and thus their number within the alveoli gradually increases. The epithelial cells lining the alveolar walls, which were only swollen and granular in the earlier stage of the process, now undergo more active changes. Their nuclei enlarge and multiply, and ultimately numerous new cells are produced. The pulmonary alveoli thus become completely filled with young cell-forms, so that the fibrinous exudation is no longer visible as an independent material, as it is in the stage of red hepatization. (See Fig. 111.) The fibrinous material now disin-

tegrates, and the young cells rapidly undergo retrogressive fatty changes, so that, as usually seen, the alveoli are filled with granular elements, which in many parts have lost their distinctive outlines. (Fig. 112.) The alveolar walls themselves, with few exceptions, remain throughout the process unaltered, although occasionally, when this stage is unusually advanced, they may be found, here and there,

FIG. 112.



Croupous Pneumonia—Gray Hepatization.—Showing the large accumulation of cellular elements within one of the pulmonary alveoli, which in some parts have undergone such extensive fatty degeneration that their distinctive outlines are no longer visible. $\times 200$.

partially destroyed. The weight, density, and friability of the lung now become even greater than in the stage of red hepatization, although the granular aspect of the cut surface is much less marked. The tissue is now quite soft and pulpy, and a puriform liquid exudes from its cut surface. The most prominent feature, however, is the alteration which takes place in the color of the organ. This gradually changes from a dark reddish-brown to a gray or yellowish-white. This is owing partly to the pressure exercised upon the bloodvessels by the exuded substances and newly-formed cells, and partly to the fatty degeneration which the latter have undergone. The stage of gray hepatization when far advanced has been termed suppuration of the lung.

Although these three stages of the pneumonic process have been described as succeeding one another in orderly succession, it must be

remembered that each stage does not occur simultaneously throughout the whole of the affected area of the lung. The changes advance irregularly, so that whilst one portion of the lung is in the stage of red hepatization, another may be in the gray stage—hence the mottled marbled appearance of the consolidation. The rapidity with which the several stages succeed one another is also subject to marked variations. In some cases the pneumonic consolidation very rapidly becomes gray, whilst in others the time occupied in the transition is much longer.

The pneumonic process may terminate in the four following ways:—

1st. *In Resolution*.—The gradual return of the lung to its normal condition is the natural and much the most frequent termination of croupous pneumonia. This is effected by the fatty and mucoid degeneration of the inflammatory products which have accumulated within the alveoli, which thus become so altered that they can be removed by absorption. Granular pigment is also mixed with the softened matters, and appears in the expectoration. This is probably partly derived from the extravasated blood, and is partly that which normally exists in the interlobular connective tissue. Where this process of liquefaction and disintegration is taking place in the lung the granular appearance of its cut surface is completely lost. It is of a yellowish-gray color, and a tenacious puriform liquid can be expressed from its substance. As the softened matters become absorbed, the circulation is gradually restored, and the organ ultimately attains its normal characters.

2d. *In Abscess*.—The formation of an abscess is a very rare result of simple pneumonia. Such a result appears to be favored by a bad constitution, and by any circumstances which tend to impair the general health either before or during the disease. The abscess is more common in the upper than in the lower lobes. Circumscribed gangrene of the lung may also occasionally terminate in abscess. This takes place by the evacuation of the gangrenous tissue through the bronchi, and the formation of a pyogenic membrane from the walls of the cavity, which generates pus. The cavity may ultimately close by granulation and cicatrization. The abscesses of primary origin are usually single, and thus differ from those due to pyæmia.

3d. *In Gangrene*.—Almost as rare as the preceding is the occurrence of gangrene. This is most common in chronic drunkards and in those of debilitated constitution. Two conditions appear to be principally concerned in bringing about this result—one is the interference

with the supply of blood by the extensive formation of coagula in the pulmonary and bronchial vessels, together with considerable hemorrhage into the pulmonary tissue; the other is the septic influence of altered inflammatory products. The gangrene is usually limited to a small area of the pneumonic lung, and is either diffuse or limited by a zone of inflamed tissue.

4th. *In Chronic Pneumonia*.—If the inflammatory process does not subside, and the exuded substances are not absorbed, the alveolar walls gradually become involved. These become thickened by a new growth of fibro-nucleated tissue, and thus is produced more or less fibroid induration of the organ. (See “Interstitial Pneumonia.”) This termination of croupous pneumonia is comparatively rare.

A croupous pneumonia plays a prominent part in some cases of pulmonary phthisis. (See “Pulmonary Phthisis.”)

Broncho- or Catarrhal Pneumonia.

Broncho-, catarrhal, or lobular pneumonia is inflammation of the lung-tissue associated with, and usually secondary to, inflammation of the bronchial mucous membrane. In the earlier stage, the pulmonary inflammation is commonly limited to scattered groups of air-vesicles, hence the term *lobular* which is applied to it. As the process proceeds, the inflammatory nodules may gradually coalesce, so as to produce larger tracts of consolidation. The inflammatory products which fill the alveoli consist principally of cells derived from the epithelium of the alveoli and from the bronchial mucous membrane; exudation and emigration play a much less prominent part in the process than they do in croupous pneumonia. Owing to this preponderance of epithelial products, and to the association of the pulmonary with the bronchial inflammation, the process has been termed *catarrhal pneumonia*.

PATHOLOGY.—The pneumonic process, as already stated, is invariably associated with bronchial catarrh. In some cases, it would appear that the injury which produces the bronchial inflammation produces at the same time inflammation of the air-vesicles, but much more frequently the bronchitis precedes the pneumonia, and gives rise to it in a manner to be hereafter described. Whatever causes inflammation of the bronchial mucous membrane may thus be a cause of broncho-pneumonia. Simple bronchitis, especially in childhood and old age, and also the specific bronchitis associated with measles and

whooping-cough, are the most frequent precursors of the disease. All conditions which tend to impair the general health favor the occurrence of the pneumonia. They do so by rendering the bronchial mucous membrane abnormally liable to become inflamed, and also by diminishing the power of the respiratory muscles, and thus aiding in the production of pulmonary collapse.

Inflammation of the bronchial mucous membrane may give rise to broncho-pneumonia in two ways: 1st, by causing in the first place collapse of the lung-tissue; and 2d, by the direct extension of the inflammation from the bronchi to the air-vesicles. The pneumonic process being the result of the bronchitis, almost invariably involves both lungs.

1. *Broncho-pneumonia consecutive to Collapse.*—Collapse of the lung-tissue greatly favors the occurrence of broncho-pneumonia, and usually the pneumonic process is principally confined to those portions of the lung in which collapse has taken place. There are two circumstances principally concerned in the production of the collapse which is consecutive to bronchitis—the narrowing of the bronchial tubes, and the weakness of the inspiratory power. The collapse thus induced is especially frequent in the posterior and inferior portions of the lungs—those portions in which normally the inflation of the lung is least complete. Commencing here, the process may gradually extend upwards till large areas of the lungs become involved. In other cases, owing to a more irregular distribution of the bronchial obstruction, the collapse is limited to small isolated portions of the lung. These portions vary in size from a hemp-seed to a walnut. They are commonly more or less wedge-shaped, with their apices toward the bronchus with which they communicate, and the lung-tissue around them usually presents various degrees of congestion and emphysema.

The tendency of the pneumonic process to occur in the collapsed portions of the lung is due partly to the hyperæmia which is induced by the collapse, and partly to the irritation of inhaled bronchial secretion. Collapse of the lung-tissue invariably induces more or less congestion. This is owing to the absence of the expansion and contraction of the air-vesicles which normally aid the pulmonary circulation, and also to the impediment to the blood-flow resulting from imperfect aëration. This congestion is quickly followed by œdema, and the bluish-purple collapsed portions of the lung become darker in color, less resistant, and more friable in consistence. In lung-tissue

thus altered a quasi-inflammatory process characterized mainly by epithelial activity is prone to supervene.

Another circumstance which often appears to play a prominent part in the causation of the pneumonic process, is the presence within the alveoli of the inflammatory products of the bronchial mucous membrane. These products are frequently found in scattered groups of air-vesicles, and they are evidently inhaled. (See Fig. 113.) They are found both in the air-containing and in the collapsed portions of the lung, but especially in the latter, the presence of collapse necessarily interfering with their removal by expectoration or absorption. These inhaled products are often found filling small groups of alveoli *without any evidence of subsequent inflammation*, and there can be no doubt that many of the patches of consolidation which are usually described as pneumonic are really non-inflammatory in their nature, and are thus produced. At the same time, owing to the irritation of the inhaled secretion, it tends to induce inflammatory changes within the alveoli, and these changes are frequently largely owing to its presence.


2. *Broncho-pneumonia independent of Collapse*.—Although the pneumonic process is usually consecutive to collapse, it may occur independently. This may be owing either to the direct extension of the inflammation from the bronchi to the air-vesicles, or to the influence of inhaled inflammatory products. In other cases, it is possible that the injury which causes the bronchitis causes at the same time inflammation of the pulmonary alveoli.

HISTOLOGY, ETC.—The appearances presented by the lungs after death vary. The bronchi are always more or less inflamed, and contain thick mucus. The lung-tissue exhibits, associated in various degrees, collapse, congestion, œdema, emphysema, and pneumonic consolidation. The bluish, non-crepitant, depressed portions of collapse, which become darker and more friable with age, are usually most abundant in the lower lobes and margins of the lungs. The collapse sometimes involves the whole of one lobe, but more commonly it is limited to smaller areas. When scattered and limited in its distribution, there is usually more or less emphysema of the intervening portions of the lung.

Those portions of the lung in which the pneumonic process has supervened most commonly appear as scattered nodules of consolidation, varying in size from a small pea to a hazel-nut. These are ill-defined, and pass insensibly into the surrounding tissue, which is

variously altered by congestion, collapse, and emphysema. They are of a reddish-gray color, slightly elevated, smooth, or very faintly granular, and soft and friable in consistence. As they increase in size they may become confluent. In a more advanced stage, the nodular and more diffuse consolidation becomes paler, firmer, drier, and somewhat resembles in appearance ordinary gray hepatization.


FIG. 113.



Broncho-Pneumonia.—From a child, aged four, with capillary bronchitis. A section of one of the patches of consolidation. Showing the stuffing of the alveoli with what appears in the main to be inhaled bronchial secretion. $\times 200$.

When examined microscopically, this consolidation is seen to consist of an accumulation within the alveoli of a gelatinous mucoid-looking substance, small cells resembling leucocytes, and epithelial elements. In many cases much of this accumulation is precisely similar to that

FIG. 114.



Catarrhal Pneumonia.—From a case of acute phthisis. Showing the large epithelial cells which fill the alveoli. $\times 200$.

contained in the smaller bronchi, and it is evidently the inflammatory and richly cellular bronchial secretion which has been inhaled. (Fig. 113.) At the same time it is in the highest degree probable that it is often partly the result of exudation and emigration from the pulmonary capillaries; for, as shown by Cohnheim, blood-stasis so injures the walls of the bloodvessels that the blood-corpuscles and liquor sanguinis readily permeate them. (See

“Embolism.”) Associated with this material are large epithelial elements, the offspring of the alveolar epithelium. These vary considerably in number. In some cases, and in some portions of the consolidation, they may be very few, whilst in others they may constitute the predominant change. (Fig. 114.) These differences probably depend upon how far the inhalation of bronchial secretion constitutes a part of the process.

The subsequent changes which take place in the lungs vary. When the disease does not end in death, resolution is the most common termination. The contents of the alveoli undergo fatty metamorphosis, and are removed by expectoration and absorption, the lung gradually regaining its normal character. This process, however, is less readily effected than in croupous pneumonia, and it often occupies such a lengthened period that some thickening of the bronchial and alveolar walls and dilatation of the smaller bronchi remain. In chronic cases this fibroid thickening is much more marked, and considerable irregularly distributed pigmented induration and bronchial dilatation may be produced. (See “Chronic Pneumonia.”) In these chronic forms the contents of the alveoli sometimes caseate and then become encapsuled, or in quite exceptional cases lead to disintegration.

HYPOSTATIC PNEUMONIA.—Allusion must be made here to a form of lung-consolidation which is often described as pneumonic, but which, in reality, is for the most part non-inflammatory in its nature. This is the so-called hypostatic pneumonia. This condition is met with at the bases and most dependent portions of the lungs in the course of both chronic and acute diseases, and also in the aged and debilitated. It consists in the main of collapse, hyperæmia, and œdema of the lung-tissue, resulting from weak inspiratory power, feeble circulation, and gravitation. The consolidation thus mechanically induced is increased by more or less exudation of liquor sanguinis and blood-corpuscles into the alveoli, which exudation is partly due to the damage to the walls of the capillaries caused by the blood-stasis (Cohnheim). This condition of hypostasis is, as has been already stated, especially favorable to proliferation of the alveolar epithelium; and this proliferation often occurs to a greater or less extent, and so tends to increase the consolidation.


Interstitial or Chronic Pneumonia.

Interstitial or chronic pneumonia is characterized by a gradual increase in the connective tissue of the lung, which leads to an indura-

tion of the pulmonary texture, and to progressive obliteration of the alveolar cavities. It is commonly associated with catarrh and dilatation of the bronchi, and often with ulceration of the bronchial walls, and excavation of the indurated lung.

PATHOLOGY.—It is exceedingly doubtful if interstitial pneumonia is ever a primary and independent affection. It probably in all cases owes its origin to some antecedent inflammation of the pulmonary or bronchial textures, or of the pleura. It may be stated generally that all inflammatory processes in the lung which become chronic lead to an increase of the connective tissue elements, and consequently to a fibroid induration of the organs, and in this respect, therefore, these processes resemble similar ones in other parts—*e. g.*, in the liver, kidney, and mucous membranes. In the lungs by far the most common cause of such induration is phthisis, in all cases of which, excepting in those which are the most acute, there is more or less fibroid growth. The most chronic cases of phthisis—those in which the fibrosis is the most marked—are, it must be admitted, somewhat closely allied to some forms of interstitial pneumonia. The two diseases differ, however, in this respect: that whereas the pulmonary consolidation of phthisis tends to undergo molecular death and disintegration, that of

FIG. 115.



Chronic Bronchitis.—Showing the new growth of fibro-nucleated tissue around the bronchus *b*, and the way in which this tissue is invading the walls of the adjacent alveoli. *v*. A divided bloodvessel. $\times 100$, reduced $\frac{1}{4}$.

interstitial pneumonia exhibits no such tendency, but any destruction and excavation of the indurated lung which may take place are due to secondary inflammation and ulceration commencing in the bronchial walls. In considering the pathology of interstitial pneumonia, there-

fore, it is necessary to exclude in the first place the pulmonary fibrosis of chronic phthisis. (See "Pulmonary Phthisis.") Interstitial pneumonia must also be separated from that form of pulmonary induration which is produced by long-continued mechanical congestion (see "Brown Induration of the Lung") and from those more localized indurations due to chronic bronchitis (Fig. 115), and to syphilis.

There appear to be four conditions which may give rise to interstitial pneumonia. These are as follows:—

1. *Croupous Pneumonia*.—The pulmonary consolidation of acute croupous pneumonia in almost all cases undergoes complete resolution. This resolution is usually effected rapidly, but occasionally it is more protracted. When protracted, the hepatized lung tends to become slightly indurated, owing mainly to thickening of the walls of the alveoli. This indurated hepatization differs but little in its physical characters from ordinary red and gray hepatization; it is simply somewhat firmer, more resistant, and less granular. In very exceptional cases this small amount of induration, commencing in the alveolar walls, may gradually increase, so as ultimately to give rise to that extensive fibrosis of the lung which constitutes what is usually known as interstitial pneumonia.

2. *Broncho-Pneumonia*.—This appears to be a somewhat more frequent cause than the preceding. The greater liability of this form of pneumonia to lead to pulmonary induration, is to be accounted for, partly by its longer duration and greater tendency to become chronic, and partly by the existence of bronchial dilatation with which it is so frequently associated. That bronchial dilatation is favorable to an indurative pneumonic process has been insisted upon by Dr. Wilson Fox.¹ The existence of this dilatation favors the persistence of the catarrhal and pneumonic process. The removal of secretion is rendered difficult, and the retained secretion tends to keep up and increase the irritative process, both in the dilated bronchi and also in the pulmonary alveoli, and this persistence of the bronchial and pulmonary inflammation leads to fibroid thickening of the bronchial and alveolar walls. In this way areas of fibroid induration are produced, which, as the process proceeds, may ultimately involve large portions of the lung. The progressive tendency of the process is probably partly to be explained by the fact that pulmonary fibrosis is a cause of bronchial dilatation, so that fibrosis once established, by inducing further dila-

¹ Reynolds' "System of Medicine," vol. iii. Article, Chronic Pneumonia.

tation of the bronchi favors the extension of the bronchial and pulmonary induration (Wilson Fox).

Under this head may also be included those cases of induration and ulceration of the lung which result from obstruction of a main bronchus—such as is produced by the pressure of an aneurism. Here the retained bronchial secretion sets up inflammatory changes in the bronchial and alveolar wall, which gradually lead to induration and ulceration of the lung.¹

3. *Pleurisy*.—This, in very exceptional cases, leads to the development of an interstitial pneumonia. It appears to be in those cases of pleurisy which are more or less chronic, and in which the effusion remains long unabsorbed, that such a result is most liable to occur. The induration of the lung thus induced, is often, however, exceedingly partial, consisting merely in some increase of the interlobular connective tissue, originating and extending inwards as dense bands from the thickened visceral pleura. In other cases pleurisy probably gives rise to a much more general fibrosis.

4. *The Inhalation of solid irritating Particles*.—This, which occurs in miners, potters, stonemasons, grinders, etc., is the cause of the fibrosis of the lungs so common amongst these workmen. The continuous irritation of the inhaled particles induces a bronchial and alveolar inflammation, and ultimately a progressive fibrosis, with dilatation and ulceration of the bronchi.


HISTOLOGY, ETC.—The appearances presented by the lung when the fibrosis is extensive and general are very characteristic. The organ is diminished in size; the tissue is smooth, dense, firm—in parts almost cartilaginous in consistence; and it is irregularly mottled with black pigment. The alveolar structure of the lung is in most parts completely destroyed, and on section the dilated bronchi are seen as numerous large openings scattered over its surface. The dilated bronchi frequently become the seats of secondary inflammatory processes, which may lead to ulceration and ultimately to extensive excavation of the indurated tissue; but there is a complete absence of any of those caseous changes which are so characteristic of ordinary phthisis. This secondary inflammation of the dilated bronchi is probably induced by the irritating and often putrid secretion which they contain, and which is only with great difficulty completely removed by

¹ See case by Dr. Irvine, "Trans. Path. Soc. Lond.," vol. xxviii. p. 63.

expectoration. The pleura is almost invariably considerably thickened and adherent.

The histological changes may be described generally as consisting in the development of a fibro-nucleated tissue from the walls of the alveoli, from those of the bronchi, and from the interlobular connective tissue; which new growth, as it increases, and from its tendency to contract, gradually replaces and obliterates the alveolar structure. The character of these changes, however, varies somewhat according to the inflammatory antecedents in which they originate. When the result of a croupous pneumonia, the primary, and usually the principal, change takes place in the walls of the alveoli (Fig. 116), although

FIG. 116.




Interstitial Pneumonia.—From a case of so-called “cirrhosis” of the lung, in which the disease was unilateral. The bronchi were much dilated, and there was a complete absence of any caseous change. The drawing shows the new fibro-nucleated growth, both in the alveolar walls and in the interlobular tissue, also the pigmentation. At *a* a divided vessel is seen. $\times 100$.¹

ultimately the interlobular tissue is involved. The alveolar walls become thickened by the growth of a small-celled tissue, in which, associated with the lymphoid cells, there are sometimes elongated fusiform cells such as are found in embryonic tissue which is undergoing fibroid development. The new growth in its earlier stages usually contains new bloodvessels, but later the tissue contracts, and these become to a great extent destroyed. The alveolar cavities which are not obliterated, are either empty, or contain exudation products or a few epithelial cells. In addition to the growth in the alveolar walls, I have met with three cases in which intra-alveolar

¹ When this specimen is examined with a higher magnifying power a delicate reticulum can be seen between the cellular elements.

exudation products were undergoing fibroid development.¹ There was nothing peculiar in the microscopical characters of the lungs, but

FIG. 117.



Chronic Pneumonia.—Vascularization and fibroid development of intra-alveolar exudation products. Bloodvessels are seen in the exudation products, which bloodvessels communicate with those in the alveolar walls. The alveolar walls are also thickened by a fibro-nucleated growth. $\times 100$, and reduced $\frac{1}{2}$.

FIG. 118.



Chronic Pneumonia.—A portion of the intra-alveolar exudation products (Fig. 117) more highly magnified. Showing the elongated spindle cells, the fibrillation, and the bloodvessels containing blood corpuscles. $\times 250$.

the alveoli were found filled with a fibrinous meshwork and leucocytes somewhat similar to those met with in red hepatization. (See Fig. 111.) They differed, however, in this respect—that many of the cells were long and spindle-shaped, and bloodvessels were distributed amongst them, which bloodvessels communicated with those in the alveolar walls. (Figs. 117 and 118.) The alveolar

walls were also thickened by a fibro-nucleated growth. It was therefore perfectly obvious that in these lungs the products of a previous acute croupous pneumonia were becoming

vascularized and undergoing development into a fibroid structure, and that this intra-alveolar change was the principal cause of the fibroid induration of the organs.

When the fibrosis is secondary to an ordinary broncho-pneumonia

¹ For one of these specimens I am indebted to Dr. Goodhart, who records the case in the "Trans. Path. Soc. Lond.," vol. xxv. p. 33.

or to that induced by the inhalation of irritating solid particles, the new growth also originates principally from the alveolar walls. Here, however, the growth in the earlier stages is less uniform and the peribronchial and interlobular connective tissue play a more prominent part in the process.

CHAPTER XLII.

PULMONARY PHTHISIS.

By pulmonary phthisis is understood a disease of the lungs which is characterized by progressive consolidation of the pulmonary texture, and by the subsequent softening and disintegration of much of the consolidated tissue. Respecting the nature of the morbid processes which lead to this consolidation and disintegration of the lungs, various opinions have from time to time been held by pathologists, and this diversity of opinion exists to some extent even at the present day. According to the older views, which were based upon the teaching of Laennec, phthisis was regarded in all cases as a *tuberculous* disease. Tubercle was looked upon as a specific non-inflammatory growth which was characterized by the caseous degeneration which it invariably underwent (see "Acute Tuberculosis"), and this caseous metamorphosis was held to be such a distinguishing peculiarity of the growth that all caseous masses came to be regarded as tuberculous, and phthisis, in which caseation plays such a prominent part, was consequently regarded as a tuberculous disease. The various consolidations of the pulmonary tissue were described as "infiltrated tubercle," and tubercle in some form or other was regarded as so essential a constituent of the disease, that "phthisis" and "pulmonary tuberculosis" came to be synonymous terms.

These older views respecting the nature of phthisis have undergone various modifications during recent years. When the application of the term "tubercle" became limited by Virchow and his followers to the gray granulation, it was evident that such views were no longer tenable, and at the present time many, in accordance with the advocacy of the late Professor Niemeyer, regard tubercle as only an occa-

sional element in the causation of the disease. In considering how far tubercle plays a part in the production of phthisis, it must be borne in mind that this growth is now known to be an inflammatory one. The miliary lesions which are commonly known as tubercular, are the anatomical results of chronic inflammatory processes limited to small circumscribed areas; and although they consist in the main of a small-celled lymphoid tissue, they present certain differences in their histological characters, according to the structure of the organ in which they originate. (See "Acute Tuberculosis.") In the lungs it has been seen that they consist not only of that lymphoid tissue and branched cells which is so characteristic of tubercular lesions, but also of accumulations of epithelium within the pulmonary alveoli. (See "Tuberculosis of the Lungs.")

HISTOLOGY OF PHTHISIS.—The histological changes in the lungs which occur in pulmonary phthisis are similar to those which are met with in these organs in acute miliary tuberculosis. They differ mainly in this respect—that whilst in the latter disease these changes are usually limited to small areas (hence the miliary character of the lesions), in the former they commonly involve much wider tracts of tissue.

In studying pulmonary phthisis, it will be advisable in the first place to describe the various structural changes which are met with in the lungs, together with the more important alterations which they produce in the physical characters of the organs; and subsequently to examine into the nature of the morbid processes upon which these changes depend, and to draw some general conclusions respecting the pathology of the disease.

The structural changes met with in the lungs in phthisis are mainly of four kinds:—1st. *An accumulation of epithelial cells within the pulmonary alveoli*; 2d. *The presence within the alveoli of a fibrinous exudation and leucocytes*; 3d. *A cellular infiltration and thickening of the alveolar walls, together with, in most cases, a similar change in the walls of the terminal bronchioles*; and 4th. *An increase in the interlobular connective tissue*. These four kinds of morbid change are very constantly associated, although in very different degrees; and some of them are more prominent and characteristic than others. The preponderance of one or other of them produces those variations in the physical characters of the lungs which are met with in the different stages, and in the different varieties of the disease. These various structural changes must now be considered separately.

1st. *An accumulation of epithelial cells within the pulmonary alveoli.*—This is one of the most frequent changes met with in phthisis,

FIG. 119.



Acute Phthisis.—Showing one of the alveoli filled with epithelial elements, and marked cellular infiltration of the alveolar wall. $\times 200$.

and is precisely similar to that which has been already described as occurring in cases of true catarrhal pneumonia. (See Fig. 114.) The alveoli are found filled with large nucleated elements, which are the offspring of the epithelial cells normally lining the alveolar walls. (Fig. 119.) In some acute cases of phthisis this alveolar accumulation may constitute almost the only morbid change, and although there is always some cellular infiltration of the alveolar walls, the great bulk of the pulmonary consolidation is due to the stuffing of the alveolar cavities with catarrhal products. (Fig. 120.) In some parts, those in which the change is the most recent, the large cells which fill the alveoli and the alveolar walls will be found but little altered, but in the greater portion of the consolidated tissue the cells will be seen in various stages of retrogressive metamorphosis, and the alveolar walls de-

FIG. 120.




Section of Lung from a Case of Acute Phthisis.—Showing that the consolidation consists almost exclusively of products accumulated *within* the alveoli. In some parts a free space is seen between the alveolar walls and their contents: this is simply due to the shrinking of the latter caused by the hardening of the specimen. $\times 50$.

stroyed; whilst in those tracts of tissue in which the process is most advanced all trace of structure is lost, and nothing is seen but a granular débris. These changes are precisely analogous to those met with in many of the larger nodular lesions of acute tuberculosis. (See Figs. 73 and 74.)

2d. *The presence within the alveoli of a fibrinous exudation and leucocytes.*—This is less frequent than the preceding. (Fig. 121.)

FIG. 121.



Acute Phthisis.—Showing one of the alveoli filled with fibrinous exudation and leucocytes, and some cellular infiltration of the alveolar wall. $\times 200$.


The exudation products are similar to those which fill the alveoli in ordinary croupous pneumonia. (See Fig. 111.) The coagulum, however, is usually not so abundant, neither is the fibrillation quite so distinct. In the most acute forms of phthisis this may constitute the principal cause of the pulmonary consolidation, but more commonly it is associated with more or less epithelial proliferation.

The appearances presented by the lungs in those cases in which the pulmonary consolidation is mainly due to the *intra-alveolar* changes above described are very characteristic. The consolidated tissue is quite soft and friable, breaking down very readily under the finger, and there is a complete absence of any induration. The consolidation, although frequently almost uniform, sometimes presents a somewhat lobulated outline, indicating the implication of different groups of the pulmonary lobules. The color varies from a reddish to a yellowish-gray, and scattered through the consolidated mass are often

small portions of a more decidedly yellow tint. These latter correspond with those parts in which the retrogressive changes are the most advanced, and they are even softer in consistence than the surrounding tissue. In many parts the consolidated tissue will be found broken down, so as to form cavities of various sizes. These always possess irregular walls, which are quite soft and friable, like the solid tissue which surrounds them.

3d. *A cellular infiltration and thickening of the alveolar walls, together with, in most cases, a similar change in the walls of the terminal bronchioles.*—This is very constantly associated with the former intra-alveolar changes, and it must be regarded as the most characteristic phthisical lesion, although its extent varies very considerably in different cases. The change is precisely similar to that which has been already described as occurring in acute miliary tuberculosis. In its earlier stages a few small lymphoid cells are seen infiltrating the alveolar septa, which are thus slightly thickened. (See Figs. 119 and 121.) As the change proceeds the number of these cells increases, and from them an imperfect fibro-nucleated structure is developed, which in some parts very closely resembles adenoid tissue. (Fig. 122.)

FIG. 122.



Section of Lung from a case of somewhat Chronic Phthisis.—Showing the thickening of the alveolar walls by a fibro-nucleated adenoid-like tissue; together with an accumulation of epithelial cells within the alveolar artery. The latter are undergoing retrogressive changes. $\times 200$.

This structure contains no new bloodvessels. As this new tissue develops in the alveolar walls it gradually obliterates and replaces the alveolar cavities, so that whilst in some portions the thickened alveoli may be found still containing epithelial elements, exudation products,

or even giant cells, in others large tracts will be seen consisting almost entirely of the small-celled growth. The development of this new non-vascular tissue in the alveolar walls leads to the partial, or even complete, obliteration of the pulmonary capillaries, which, as will be seen subsequently, constitutes an important element in the causation of the retrograde changes.

The changes which may subsequently take place in this alveolar growth vary. The infiltrated septa may rapidly break down before any marked thickening or development of new tissue has had time to occur; whilst in other less acute cases, there is a considerable development of the imperfect fibro-nucleated tissue, which, although it may remain as a more or less permanent structure, usually owing to insufficient vascular supply, undergoes in its turn retrogressive metamorphosis. These two kinds of change are very often found taking place simultaneously in different portions of the consolidated lung. In those portions in which the new tissue is undergoing degeneration, it, together with the cells which may be contained within the alveoli, will be seen to have become converted into a structureless granular débris, whilst perhaps in immediate vicinity to these more degenerated portions will be found a more permanent fibro-nucleated structure.

Respecting the alteration which the growth of this small-celled tissue produces in the physical characters of the lungs, it may be stated, generally, that it usually leads to more or less induration of the pulmonary texture. The extent of this induration, however, will vary according to the characters of the new tissue. If the tissue be almost entirely cellular, such as is the case when it is very rapidly developed, it will produce but little, if any, induration of the pulmonary consolidation, which, consisting mainly of the intra-alveolar accumulations, will be soft and friable in consistence, much resembling that which has been already described. When, on the other hand, as is more frequently the case, there is any considerable development of the imperfect fibro-nucleated growth, or its reticulum is dense and abundant, there will be a corresponding induration of the consolidated tissue. In many cases these changes produce uniform tracts of indurated consolidation of a grayish color mottled with black pigment, in which there may be scattered here and there yellowish patches corresponding to those portions which have undergone retrogressive fatty changes.

4th. *An increase in the interlobular connective tissue.*—This is met with, to a greater or less extent, in all the more chronic forms of phthisis. This tissue, which surrounds the bronchi and bloodvessels,

and contributes to the formation of the alveoli, is found not only increased in amount, but also altered in character. In the earlier stages of its development, when it contains numerous small cells, although many parts of it may resemble the growth in the alveolar walls, its structure is more like that met with as the result of chronic indurative processes in other organs. It has a much greater tendency to become developed into a fibroid tissue than the alveolar growth, and is rarely the seat of those retrograde changes which are so frequent in the tissue originating in the alveolar walls. As usually met with, it consists either of wavy fibres or of a more or less reticulated structure, with a varying number of round, spindle-shaped or branched cells. (Fig. 123.) Associated with it, in most cases, are granules of black pig-

FIG. 123.



Chronic Phthisis.—Showing the new interlobular fibroid growth surrounding and encapsulating a degenerated and caseous portion of the consolidated lung. $\times 50$, reduced $\frac{1}{2}$.

ment. These differences in the pathological tendencies and structure of the alveolar and interlobular growths are mainly owing to differences in the amount of their vascular supply. Whereas in the former the vessels become obliterated in the manner already described, in the latter this obliteration is much less complete or entirely wanting. In the most chronic cases of phthisis this interlobular growth may constitute the predominant structural change, and large tracts of the pul-

monary texture may be found completely replaced by it. (See "Interstitial Pneumonia.")

An increase in the interlobular connective tissue in phthisis—inasmuch as the new tissue has so marked a tendency to become dense and fibroid—leads to extensive induration of the pulmonary texture; and further, owing to the contraction which the tissue tends to undergo, its growth ultimately produces a corresponding contraction of the diseased lung. In all those cases of phthisis in which there is either a marked thickening of the alveolar walls, or an increase in the interlobular connective tissue, any cavities which may exist in the consolidated and indurated tissue are characterized by the tough and fibroid character of their walls, these presenting a marked contrast to the soft friable tissue which surrounds the cavities in those cases in which the pulmonary consolidation is mainly due to intra-alveolar changes.

Changes in the Bronchi.—Allusion must now be made to certain changes in the bronchi. These tubes are invariably more or less involved in pulmonary phthisis. Some catarrh of the bronchi is constantly present in phthisical lungs. The catarrh is sometimes general, but much more commonly it is limited, and more strictly confined to such portions of the lung as are becoming, or have already become, consolidated. In many cases there is a marked tendency of this bronchial catarrh to lead to extensive cellular infiltration of the deeper structures of the bronchial wall. This is especially the case in the scrofulous. (See "Scrofulous Inflammation," Fig. 63.) This cellular infiltration sometimes leads to the production of small ulcers. These have thickened opaque edges, and when once formed they tend to increase. In addition to these changes in the bronchial mucous membrane, there is often a cellular infiltration of the peri-bronchial tissue, and here small nodules of new growth are frequently met with—especially around the smallest bronchi. The development of these nodules is probably due to the transmission of infective substances by means of the lymphatics from the bronchial mucous membrane (tuberculosis).

PATHOLOGY OF PHTHISIS.—Having thus briefly described the various structural changes met with in the lungs in phthisis, it remains to consider the nature of the morbid processes upon which they depend. In the first place, it is evident that these changes are analogous to those which have been seen to occur in the several forms of pulmonary inflammation. The fibrinous exudation and leucocytes, and the accumulation of epithelial cells within the alveoli in croupous and catarrhal

pneumonia, with, in the more chronic cases, the ultimate infiltration of the alveolar walls; and the increase in the interlobular connective tissue which characterizes the interstitial process, closely resemble the phthisical lesions. These considerations, together with those derived from the study of the etiology of the disease, are sufficient to justify the conclusion that the morbid processes which lead to the consolidation and subsequent disintegration of the lung come within the category of *inflammation*, and that the differences in the histological changes to which they give rise are mainly due to differences in the intensity, in the duration, and in the mode of origin of the inflammatory process.

In considering the causes of these differences in the histological changes in the lungs, it is important to bear in mind what has been already stated respecting the variations in the character of the textural alterations in inflammation which are produced by differences in the intensity and duration of the inflammatory process. (See "*Varieties of Inflammation.*") When studying the process of inflammation, it was seen that the most intense forms of the process were characterized by abundant fibrinous exudation and emigration; whereas in inflammations of less intensity, the textural changes played a more prominent part. These textural changes also varied according to the intensity of the inflammation. In the least severe and most chronic forms, these changes tended to be limited to the elements immediately adjacent to the bloodvessels and lymphatics, whereas in inflammations of somewhat greater intensity more distant elements became involved. Further, whereas in the former case these changes usually resulted in the formation of a small-celled tissue which tended to become fibroid, in the latter, the more distant elements—being in most cases incapable of further development—tended to undergo retrogressive changes. In the lungs, the truth of these propositions was borne out by the differences which were seen to exist in the histological characters of the lesions in the various forms of pulmonary inflammation, and also in acute tuberculosis.

If the pathology of these inflammatory processes in the lungs be kept in view, the explanation of the differences in the histological characters of the lesions in pulmonary phthisis becomes tolerably evident. In those cases in which the inflammatory processes are of slight intensity and of long duration, the most marked structural change will consist in the development of a small-celled growth in the alveolar walls and in the interlobular tissue—a growth which tends,

more or less, to become developed into a fibroid structure; whereas in those cases in which the inflammation is of greater intensity fibrinous exudation and emigration, and proliferation of the alveolar epithelium, will constitute more prominent parts of the process.

The intensity of the inflammatory process not only determines to a great extent the histological characters of the pulmonary consolidation, but also the subsequent changes which take place in it. In those cases of phthisis in which the intensity of the inflammatory process is considerable, not only do the epithelium and exudation products which have accumulated within the alveoli quickly degenerate and break down, but any small-celled tissue which may have been developed in the alveolar walls or around the terminal bronchioles also softens and dies, and thus the vitality of large tracts of the pulmonary consolidation may become destroyed. In those cases, on the other hand, in which the process is less intense, the small-celled growth produced in the alveolar and bronchial walls is more permanent, and there is an increase in the interlobular connective tissue. It is these two kinds of change, the one tending towards death, and the other towards progressive development, which produce the caseation and softening on the one hand, and the induration on the other, which, associated in such various degrees, make up the diverse physical characters of the phthisical lung.

These various secondary changes which may take place in the pulmonary consolidation of phthisis must be considered more fully. They are of three kinds—resolution, development into an imperfect fibroid tissue, and retrograde metamorphosis.

Resolution.—Much of that consolidation of the lung which is the most rapidly induced, and which is consequently owing to the presence of intra-alveolar exudation matter, may become absorbed. The resolution of the consolidation may thus be complete, or after the absorption of the intra-alveolar products there may remain more or less infiltration of the alveolar walls.

Fibroid Development.—This, as has been seen, may take place in the growth in the alveolar walls, and also in the new interlobular tissue. The tissue which originates in the walls of the alveoli, however, being for the most part destitute of bloodvessels, is incapable of forming a very mature structure, although it may become developed into an imperfect adenoid-like tissue, which may remain for some time permanent, and so contribute to the induration of the lung. In the new interlobular tissue, there is not the same interference with the

vascular supply, and hence this forms a much more fully developed and permanent structure, and it is the principal source of the pulmonary fibrosis. The extent of this fibrosis is, for the most part, in direct proportion to the chronicity of the disease.

Retrograde Metamorphosis.—It is this kind of change which leads to that caseation, softening, and disintegration which is so characteristic of phthisis, and which distinguishes phthisical from other forms of pneumonic consolidation. A retrograde change in the inflammatory products is an invariable accompaniment of acute non-phthisical pneumonia. Much of the exudation matter and epithelium which fill the alveoli undergoes fatty and mucoid changes, and as the circulation becomes restored in the pulmonary capillaries, the degenerated products are absorbed, and the lung remains intact. In phthisical consolidation, however, this removal of the inflammatory products does not take place. The contents of the alveoli degenerate, but the degenerated products are not absorbed, and the consolidated lung undergoes a rapid or gradual process of disintegration.

In studying the causes of this retrograde metamorphosis, which constitutes so essential a feature of the disease, it is undoubtedly principally due to conditions interfering with the circulation. Of these conditions, that which occupies the most prominent place is that cellular infiltration of the walls of the alveoli and smaller bronchi which is such a constant though very variable factor in phthisis. It has been seen that this infiltration is especially characteristic of scrofulous inflammations, and that it occurs in a modified form in those who are not markedly scrofulous, and also in all pulmonary inflammations which become chronic. When the infiltration is marked, and especially when rapidly induced, the effect of the pressure which the young cells exercise upon the pulmonary capillaries is to obstruct the circulation, and so not only to prevent the absorption of any intra-alveolar products, but also to lead to necrotic changes.

There are two other conditions which, although of less importance than the preceding, also tend to interfere with the circulation, and so to cause necrosis. These are the pressure which is exercised upon the pulmonary capillaries by the inflammatory products which have accumulated within the alveoli, and that tendency to stagnation of the blood-stream which is an invariable accompaniment of every acute inflammation. The operation of these conditions obtains in the most acute forms of phthisis.

In addition to the interference with the circulation, an important

element in the causation of the retrograde changes of phthisis is probably that inherent weakness of the lungs (usually inherited), which not only renders them especially susceptible to injury, but also, when injured, renders them abnormally incapable of recovering themselves from the inflammatory process which has been induced.

In many cases of phthisis also, especially in the more chronic forms, secondary inflammation and ulceration of the pulmonary consolidation, resulting from the injurious influence of retained secretions and inflammatory products, constitutes an important factor in the causation of the destruction of the lung.

ETIOLOGY OF PHTHISIS.—In studying the etiology of phthisis it will be necessary to consider, in the first place, that inherent pulmonary weakness which exists so frequently in this disease; secondly, to examine into the several methods by which injuries may be inflicted on the lung in such a way as to set up an inflammatory process; and lastly, to point out in what way the occurrence and progress of the disease are influenced by the general health of the individual.

Inherent pulmonary weakness, usually inherited but often acquired, exists in a large proportion of the cases of phthisis. This weakness not only renders the lungs abnormally susceptible to the various kinds of injurious irritation, but also makes them less capable than strong lungs of recovering themselves from the effects of the inflammatory process which the injury has produced. Inherent pulmonary weakness is most frequently, but by no means always, a part of that general constitutional state known as scrofula, in which the mucous membranes generally, and especially that of the respiratory organs, are so abnormally liable to become inflamed.

Inflammatory process may be induced in the lungs by—1st, injuries inflicted upon the surface of the body; 2d, by injuries inflicted through the medium of the bronchi; and 3d, by infection.

By injuries inflicted upon the surface of the body I mean that general chilling of the surface which results from exposure to cold. That this is a frequent cause of bronchial and pulmonary inflammation is well known.


A more important agent in the production of phthisis than the preceding is the injurious irritation of the lungs through the medium of the bronchi. The bronchial mucous membrane, communicating as it does with the external air, is especially liable to injury, and the extreme frequency with which the development of phthisis is preceded or accompanied by successive attacks of bronchial catarrh is a well-

established clinical fact, the important bearing of which upon the successful treatment of the disease it would be difficult to over-estimate. In considering bronchial catarrh as a cause of phthisis it is important to bear in mind how greatly the occurrence and character of this catarrh are influenced by the existence of scrofula.

In other cases injuries inflicted through the medium of the bronchi appear to exercise their influence directly upon the inner surface of the air-vesicles without producing any inflammation of the bronchial mucous membrane. This is probably a very important way in which such injuries cause phthisical consolidation of the lung.

The prominent position which these injuries inflicted through the medium of the bronchi occupies in the causation of phthisis is shown by the fact that the lung-consolidation in this disease is, for the most part, *lobulated* in its distribution. Groups of air-vesicles communicating with the terminal bronchioles become implicated, and thus are produced lobular lesions. (Fig. 124.) This lobulated distribution of

FIG. 124.



Acute Phthisis.—A transverse section of a terminal bronchus (air-passage) and the surrounding alveoli. Showing the lobulated character of the pulmonary consolidation. *b*, cavity of bronchus containing a little mucus. *v*, a bloodvessel. $\times 50$, reduced $\frac{1}{2}$.

the phthisical consolidation is exceedingly characteristic, and even in those acute cases, in which, owing to the rapid and extensive implication of the lung, the consolidation may to the naked eye appear almost uniform (like a croupous pneumonia), the microscope will usually reveal a lobular character.

The last method by which injuries may be inflicted upon the lungs is by means of infection. This infection is precisely similar to that

which has been already described as causing general tuberculosis. By it is understood the property possessed by certain inflammatory products—especially by those which have become caseous—of inducing secondary inflammatory processes in other parts. Limited tuberculosis differs from the general disease only in the extent to which the infective materials are distributed. Whereas in the general affection the dissemination of the infective substances takes place principally by means of the bloodvessels, in the more limited processes the dissemination is effected by means of the lymphatic vessels or serous canals. In the lungs, limited infective processes are exceedingly common in phthisis, and they give rise to many of the miliary nodules of induration which are so frequently met with in phthisical lungs. The source of infection in these cases is usually some caseous induration already existing in the lung, so that the infective process usually supervenes upon pre-existing phthisis.

There remains still one other factor which has a most important influence upon the causation of phthisis—the *general health of the individual*. That the development and progress of phthisis are greatly influenced by the state of the general health is well known. That this should be so will be readily understood if the inflammatory nature of the disease be kept in view, inasmuch as, in all cases, the susceptibility of a living tissue to injury is, *cæteris paribus*, greater, and the power of recovering itself from the effect of the inflammation less, the lower the standard of the general health.

VARIETIES OF PHTHISIS.—In conclusion, allusion must be made to the different varieties of pulmonary phthisis. How far is a subdivision of the disease into different *pathological* varieties admissible? Although the variations observed in the physical characters of phthisical lungs and also in the clinical history of the disease, have led to its subdivision into different kinds, it appears to me that there are no pathological grounds for any such subdivision. The variations observed in the clinical history of phthisis and in the physical characters of phthisical lungs are, I think, mainly to be ascribed to variations in the duration and intensity of the inflammatory processes which give rise to the pulmonary consolidation, and also to the parts which primary pulmonary inflammation, bronchial inflammation, and infection play respectively in the causation of the disease.

CHAPTER XLIII.

CHANGES IN THE BLOOD AND CIRCULATION.

Hyperæmia.

HYPERÆMIA, or congestion, is excess of blood in the more or less dilated vessels of a part. Whatever increases the pressure of the blood, or diminishes the resistance of the vessels, may be a cause of hyperæmia. Hyperæmia is *active* or *arterial*, and *mechanical* or *venous*. These two varieties must be considered separately.

Active Hyperæmia.

Active hyperæmia is an excess of blood in the arteries of a part, with, in most cases, an acceleration of the flow.

CAUSES.—The causes of active hyperæmia may be divided into those which increase blood-pressure, and those which diminish arterial resistance.

1. *Increased Blood-pressure*.—This occurs most commonly from interruption of the main current of blood in any particular part owing to which increased pressure is thrown upon the collateral vessels. These vessels thus become dilated, the amount of blood in them is increased, and the flow is accelerated. This, which is known as *collateral hyperæmia*, is seen after the obstruction of the main current from any cause, as from the ligature of the vessel, or from its occlusion by a thrombus or embolus. (See “Embolism.”)

General obstruction in the capillaries of a part will in the same way cause a compensatory hyperæmia. This is exemplified by the application of external cold causing contraction of the superficial capillaries and congestion of internal organs; and by obstruction of the capillaries in one part of an organ causing hyperæmia of the parts adjacent.

2. *Diminished Arterial Resistance*.—This is much the most frequent cause of active hyperæmia. It may arise from—

a. *Relaxation or paralysis of the wall of the vessel*.—The relaxation of the muscular coat of the vessels, and their consequent dilatation,

may be owing to—1st, external warmth; 2d, direct paralysis of the vaso-motor nerve; 3d, indirect paralysis of the vaso-motor nerve; and 4th, changes in the walls of the vessel resulting from injury independently of the nervous centres. External warmth is a common cause of hyperæmia. This is seen in the effect of warm baths, fomentations, etc.

The effects of direct paralysis of the vaso-motor nerves are seen in the active congestion of the head and neck which follows pressure upon the sympathetic in the neck, as by an aneurism; and in the unilateral congestion which results from experimental sections or disease of one-half of the spinal cord. Some emotional conditions also are attended by paralysis of the vascular nerves and consequently by active hyperæmia: this is seen in blushing. Certain substances again taken internally produce vaso-motor paralysis, as the nitrite of amyl, alcohol, tobacco, etc.

Indirect or reflex paralysis of the vaso-motor nerves is most frequently due to irritation of sensory nerves. The dilating action of the irritated sensory nerve is usually confined to the region supplied by it. Of the numerous examples of hyperæmia from this cause may be mentioned that which follows injury of the conjunctiva, the hyperæmias associated with facial neuralgias, and the priapism that often results from irritation of the urethra. The hyperæmia of the skin, caused by friction and irritating substances, is also due to the same cause.

Hyperæmia from the direct injury of the walls of the vessel independently of the nervous system, has already been described when speaking of vascular dilatation as the earliest change in the process of inflammation. In inflammation, the irritation is so severe as to cause not only dilatation of the vessel and accelerated blood-flow, but also a subsequent retardation of the circulation and exudation of liquor sanguinis and blood-corpuscles. (See "Inflammation.") If the injury be less intense or less prolonged in its action it produces simply dilatation of the vessels and increased rapidity of flow—*i. e.*, active hyperæmia.

β. *Sudden removal of external pressure.*—The sudden removal of external pressure from vessels is followed by their dilatation, and consequently by hyperæmia. As examples of hyperæmia from this cause may be mentioned that which results from dry cupping, and from the sudden removal of ascitic fluid, and of the fluid from a hydrocele.

γ. *Atony of the walls of the vessels from mal-nutrition.*—This is a much less important cause of hyperæmia. Fatty degeneration of the muscular and internal coats of the smaller arteries may, however, in some cases lead to their dilatation, and thus be a cause of active hyperæmia.

RESULTS.—The results of active hyperæmia are principally such as might be expected to follow from an increase in the amount of the arterial blood, and in the rapidity of its flow, in any particular organ or tissue. There is increased redness and pulsation, a sensation of throbbing being often experienced by the patient. There is also some increase in bulk. The temperature at the same time undergoes a marked elevation, sometimes as much as 3° Cent. Serous effusions, hemorrhage, and thrombosis—so common as results of mechanical hyperæmia—are rarely met with. If the hyperæmia be of long duration, the small arteries become permanently enlarged, their walls gradually thicken, and the peri-vascular connective tissue may increase. Function may, or may not, be interfered with. It is in the nervous centres that functional changes are most marked. They include great excitability, paræsthesiæ of sight and hearing, convulsions, etc. In the skin, kidneys, and other secreting organs, the secretions are usually increased.

Mechanical Hyperæmia.

In mechanical hyperæmia the excess of blood is principally in the veins, and the flow, instead of being accelerated, is retarded.

CAUSES.—The causes of mechanical hyperæmia are such as interfere with the return of the blood by the veins, either by directly impeding its exit from any vein or system of veins, or by diminishing the normal circulating forces. They are—

1. *A Direct Impediment to the Return of Blood by the Veins.*—This is the most fertile cause of mechanical hyperæmia. Any obstruction to the return of blood by the veins is followed by distension and impeded flow behind the obstruction. The congestion of some of the abdominal viscera which results from the obstruction to the portal circulation in cirrhosis of the liver; that of the lung in mitral constriction and regurgitation; that of the systemic circulation in insufficiency of the tricuspid valve; and that of the lower extremities from the pressure of the gravid uterus on the iliac veins, are a few of the numerous familiar examples of mechanical hyperæmia from this cause.

2. *Gravitation*.—This becomes an important auxiliary in the production of hyperæmia in disease, especially when it is associated with diminished cardiac power. The effect of gravitation in determining congestion of the most dependent parts is exemplified in chronic exhaustive and in many of the acute febrile diseases, in which the nutrition generally becomes impaired, the heart's power weakened, and in which the patient is unable frequently to change his position. The integuments of the back, and the posterior portions of the lungs, are the parts which are thus most frequently affected.

3. *Increased Local Resistance*.—This results from diseased conditions of the arterial walls, owing to which they either lose their elasticity and contractility and thus their power of equalizing and regulating the blood-flow, or become considerably enlarged. In either case the circulation will be impeded, there will be an accumulation of blood and retardation of flow in the veins beyond, and the veins themselves will gradually become relaxed and dilated. Such conditions may arise from simple atony of the arterial walls, or from atheromatous, fatty, or calcareous changes. They are most common in advanced life. The part they play in the production of senile gangrene has been already alluded to. (See "Senile Gangrene.")

4. *Diminished Cardiac Power*.—This is one of the most important causes of mechanical hyperæmia, especially when it is associated with any of the preceding ones. The motor power of the heart becomes impaired in many of the chronic exhausting diseases, also in the acute febrile diseases, as in typhus and typhoid fever, and in all those conditions of degeneration and softening of its structure which lead to the dilatation of its cavities. In whichever of these ways the *vis a tergo* is diminished, it will tend to produce venous hyperæmia.

RESULTS.—Long-continued mechanical hyperæmia leads to impairment of vitality and function. The tissues gradually atrophy and undergo retrogressive changes, although from the amount of serosity and blood which they contain their size and absolute weight may be increased. Their temperature becomes lowered. The most important results, however, of this variety of hyperæmia are *transudation of serum, hemorrhage, fibroid induration, thrombosis, and gangrene*.

1. *Transudation of Serum*.—This gives rise to œdema and dropsical effusions. Its occurrence is greatly favored by the stretching of the walls of the vessels, and by the damage which their structure sustains. The transuded serum usually differs from blood-serum in being of lower specific gravity, and in containing more water and less

of the solid constituents. The greater the pressure, the more nearly does the transuded liquid resemble the liquor sanguinis, and the greater is the amount of albumen which it contains. If the pressure be very great, it may yield a fibrinous coagulum. In mucous membranes mechanical hyperæmia gives rise to chronic catarrh.

2. *Hæmorrhage*.—This is another result of mechanical hyperæmia. It usually occurs only when the obstruction to the venous current is very great. Those vessels which are the least supported are the first to give way. The hæmorrhage into the stomach in cirrhosis of the liver, and into the lung in mitral disease, are familiar examples of hæmorrhage from this cause.

Not only does blood escape from the vessels by rupture of their walls in mechanical hyperæmia, but the red blood-corpuscles pass through the walls of the capillaries into the surrounding tissues without rupture taking place. This passage of the red corpuscles through the capillary walls, which was discovered by Cohnheim, may be observed in the web of the frog's foot after ligature of the femoral vein. The corpuscles in passing through the vessel become constricted in their centre, so as to assume an hour-glass shape. This emigration only occurs when the obstruction is considerable.

3. *Fibroid Induration*.—This, which is due to a gradual increase in the connective tissue around the bloodvessels, is one of the most important results of long-continued mechanical hyperæmia. The interstitial growth leads to atrophy of the other structures, and thus to impairment of the functions of the organ. In the stomach, it produces atrophy of the glandular structures; in the kidney, compression of the urine-tubes; in the liver, obstruction to the portal circulation; in the heart, diminution in motor power. The alterations which this change produces in the physical characters of the organs—viz., a hardness and induration associated with abnormal redness or pigmentation due to the excess of blood, are exceedingly characteristic.


4. *Thrombosis*.—This, as a result of mechanical obstruction, will be described in the following chapter.

5. *Gangrene*.—This only occurs from mechanical hyperæmia when the obstruction is very general and complete. It has been already described under the head of "Gangrene."

MECHANICAL HYPERÆMIA OF THE LIVER—NUTMEG LIVER.—Long-continued mechanical hyperæmia of the liver gives rise to the condition known as *Nutmeg Liver*. This is the condition which so frequently results from disease of the heart. The change is cha-

racterized by a large accumulation of blood in the hepatic veins, by dilatation and thickening of these veins, by atrophy of the hepatic cells in the central portions of the acini, and by an increase in the interlobular connective tissue.

FIG. 125.



Nutmeg Liver.—Showing the destruction of the liver-cells and the pigmentation in the central portions of the acinus, together with the new growth of connective tissue at the periphery. V. Hepatic vein. P. Portal canal. $\times 50$. (When the specimen is more highly magnified, the central hepatic vein and its tributaries are seen to be filled with red blood-corpuscles, and their walls thickened. The intercellular network which immediately surrounds it is also thickened, and the peripheral connective tissue growth contains numerous nuclei.)

The impediment to the return of blood by the hepatic vein leads to atrophy of the cells in the central portions of the acini and also to the formation of granular pigment, so that when examined microscopically, these portions of the acini are seen to consist of broken-down cells and granules of pigment. (Fig. 125.) The veins here are found much dilated and filled with red blood-corpuscles. Their walls are thickened, and there often appears to be also more or less thickening of the intercellular network which immediately surrounds the central vein. Owing to this thickening of the central veins and of the adjacent intercellular network, and to the destruction of the liver-cells, the most central portions of the acini, in advanced stages of the disease, may present a very fibrous appearance. At the periphe-


ral parts of the acini the new interlobular growth is seen insinuating itself between the almost unaltered liver-cells. This new interlobular growth is usually distinctly nucleated, but for the most part, less so than that met with in cirrhosis. Its cellular character has been especially insisted upon by Dr. Wickham Legg.

In the earlier stages of this affection the liver is often considerably increased in size from the large amount of blood which it contains. On section, it presents a peculiar mottled appearance, the centre of the lobules being of a dark red color, whilst the peripheral portions are of a yellowish-white. This latter appearance is often increased

by the presence of more or less fat in the peripheral liver-cells. Ultimately the organ may undergo a gradual diminution in size. This is due partly to the atrophy of the cells in the central portions of the lobule, and partly to the pressure of the interlobular growth. The interlobular growth tends to cause obstruction to the portal circulation, as in cirrhosis.

MECHANICAL HYPERÆMIA OF THE LUNGS.—In the lungs long-continued mechanical hyperæmia produces that peculiar induration and pigmentation of the organs which is known as *Brown Induration*. This condition most frequently results from stenosis and insufficiency of the mitral orifice. The alterations produced in the pulmonary texture consist in the first place of elongation and dilatation of the pulmonary capillaries, so that even in uninjected preparations the alveolar walls appear abnormally tortuous. The epithelial cells lining the alveoli also become swollen and probably multiply, and they are seen in large numbers, filled with dark brown pigment, covering the alveolar walls. (Fig. 126.) They frequently accumulate within the

FIG. 126.



Brown Induration of the Lung.—Showing the abnormal number of swollen pigmented epithelial cells covering the alveolar walls, the increase of connective tissue around the bloodvessel, *a*, and the large quantity of pigment. *b*. The alveolar cavity. $\times 200$.

alveolar cavities. These changes are followed by an increase in the interlobular connective tissue, by the formation of large quantities of brownish-black pigment, and often by a thickening of the alveolar walls. Sometimes the pulmonary capillaries rupture, and blood is extravasated into the lung-tissue.

Lungs in which these changes are at all advanced present a more

or less uniform brownish-red tint, mottled with brown or blackish-colored specks and streaks. They are heavier and tougher than natural, less crepitant, and upon squeezing them the pulmonary tissue is found to be denser and thicker than that of a healthy lung.

POST-MORTEM APPEARANCES OF HYPERÆMIA.—The post-mortem appearances presented by hyperæmic organs and tissues vary considerably. Very frequently parts which were hyperæmic during life show no signs of it after death. If the blood does not coagulate rapidly it passes on into the veins, and thus the recognition of arterial and capillary hyperæmia very often becomes impossible. The effect of gravitation must also be taken into account in estimating hyperæmia. After death the blood naturally gravitates to the most dependent parts: this is seen in the post-mortem congestion of the posterior portions of the lungs, and of the most dependent portions of the various coils of the intestine. The uniform redness of post-mortem staining, again, must not be confounded with the redness of hyperæmia. In capillary and arterial hyperæmia, the color is red, and the injection often presents the appearance of a capilliform network. If very intense it may to the naked eye appear uniform, but a lens will always discover its capillary nature. When the veins are the seat of the hyperæmia the injection is called ramiform, and the color is dark blue.

The anatomical peculiarities in the distribution of the bloodvessels will, however, materially affect the appearance of the hyperæmia. In the intestines it is often punctiform, being situated in the vessels of the villi; so also in the kidney, when its seat is the Malpighian corpuscles. A punctiform appearance may also be produced by minute extravasations of blood. If the hyperæmia is of long-standing, the tissue becomes pigmented. This is often well seen in the stomach and intestines, also in the lungs.

CHAPTER XLIV.

THROMBOSIS.

THROMBOSIS is a coagulation of the blood within the vessels during life. This coagulation is owing to changes in the walls of the vessel itself, or to impeded bloodflow. The coagulum is called a *thrombus*. It may form in the heart, in the arteries, in the capillaries, or in the veins. It is much the most common in the last-named vessels.

Thrombi must be distinguished from the coagula that form after death, and also from those formed in the last moments of life which are so commonly found in the cardiac cavities. Post-mortem coagula are soft, and are often divisible into two layers, colored and uncolored; they do not adhere to the walls of the vessel, and rarely completely fill its cavity. The clots formed in the heart just before death constitute a connecting link between post-mortem coagula and thrombi. They are more or less decolorized, and are firmer in consistence and more fibrinous than post-mortem clots. They are not firmly adherent to the cardiac walls, but are often so entangled amongst the columnæ carneæ, chordæ tendineæ, and papillary muscles, that they cannot be quite readily separated. They appear to be the result of the mechanical defibrination of the blood by the cardiac contractions a little while before death; the contractions not being sufficiently strong to empty the cavities, some of the blood remains behind, and becomes "whipped up" and defibrinated. These clots are most common in the right cardiac cavities, and they often extend some way into the pulmonary artery, from which, however, they can very readily be removed. They are met with most frequently in those cases in which the death-struggle has been prolonged, and in which there has been a gradual loss of power in the cardiac contractions. The proportion of fibrogenous substance in the blood will also materially influence the extent of their formation.


A *thrombus*, or ante-mortem clot, is firmer, dryer, and more fibrinous than either of the preceding, and it is adherent to the walls of the vessel. Its characters, however, vary with its age, and with the circumstances under which it originates. When freshly-formed it is

of a dark red color and soft gelatinous consistence, closely resembling the post-mortem clot. It gradually becomes paler, dryer, less elastic, and more friable. If it is rapidly produced—as when the circulation is suddenly arrested in a portion of a vessel by the application of a ligature—the coagulum at once completely fills the vessel, and as it becomes firmer it maintains a more or less uniform structure. Thrombi, however, which have undergone a slow and gradual formation, are rarely thus uniform in structure, but are made up of numerous concentric layers, and so present a *stratified* appearance. This is owing to the coagulation taking place gradually upon the inner surface of the vessel, and to the white corpuscles adhering to the successive layers of coagulum. These corpuscles, from the property which characterizes them of adhering to one another and to the sides of the vessels, and especially to any porous substances with which they may come in contact, cohere with the fibrinous layers as they are deposited; so that ultimately the thrombus is made up of layers of fibrinous coagulum and white corpuscles more or less alternately and concentrically arranged. This gives to it a stratified appearance.

The thrombus may completely or only partially fill the cavity of the vessel. In most cases, however, when coagulation has commenced, it proceeds until the vessel is obstructed, and when once this has occurred, the formation of the thrombus continues to extend in the course of the vessel until it meets with a current of blood strong enough to arrest its progress. Its ultimate extent will thus mainly depend upon the vessel in which it is formed, upon the size and situation of the collateral branches, and upon the force of the circulating current. The direction in which the coagulation principally extends, whether in the arteries or veins, is consequently backwards, from vessels of smaller to those of larger calibre; the formation of the thrombus continuing until it meets with a current sufficiently strong to restore the circulation, which in many cases is as far as the entrance of the next large collateral vessel. The end of the thrombus next the heart is rounded and conical in shape. (See Fig. 129, c.)

The thrombus when once formed either becomes *organized* or *softens*.


Organization.—This is most frequent in the uniform non-stratified thrombi, and especially in those occurring in arteries. The process consists in the gradual transformation of the thrombus into connective tissue. A thrombus which is undergoing a process of organization gradually diminishes in size, it becomes more and more decolorized, firmer and more fibrous in consistence, its union with the wall of the



vessel becomes more intimate, and ultimately it becomes converted into a fibro-cellular cord. These changes are probably principally owing to the white blood-corpuscles which it contains.

Soon after the formation of the thrombus the number of young cells


FIG. 127.



Section of an Arterial Thrombus Thirty-seven Days Old.—a. New bloodvessels. b. Leucocytes and anastomosing cells. (Rindfleisch.)

in it becomes greatly increased. In a somewhat more advanced stage the red blood-corpuscles almost entirely disappear, the clot becomes firm and homogeneous, and in addition to the small round elements, numerous anastomosing spindle-shaped cells make their appearance. (Fig. 127.) The intercellular material then becomes fibrillated, the walls of the vessel become infiltrated with cells, and numerous new bloodvessels are formed which intersect the thrombus in all directions. These vessels communicate with the cavity of the thrombosed vessel, and with its vaso vasorum. (Fig. 128.) The vascular fibrillated structure into which the thrombus has become organized gradually undergoes a process of atrophy and contraction, the new vessels disappear, and ulti-

FIG. 128.



Longitudinal Section of the Ligatured End of the Crural Artery of a Dog, Fifty Days after the Application of the Ligature.—Showing the newly-formed vessels in the thrombus and their communication with the vaso vasorum. Th. Thrombus. M. Muscular coat. Z. External coat and vaso vasorum. × 20. (O. Weber.)

mately it becomes converted into a fibro-cellular cord. In some cases it becomes calcified, and thus forms a phlebolite.

Respecting the source from which the large number of new cells which make their appearance in the thrombus are derived—it appears to be probable that they originate partly from the white blood-corpuscles and partly from the endothelial and connective-tissue cells belonging to the walls of the vessel. Recent researches tend to show that the endothelium takes a prominent part in the process. Whether, however, they are all of them the offspring of these elements, or whether some of them may not be leucocytes which have penetrated from without, is unknown. From these small cells the elongated connective-tissue cells are produced. The exact source from which the fibrillated intercellular material is derived—whether the intercellular coagulum itself fibrillates, or whether it disappears and the fibres are derived from the protoplasm of the cells—is uncertain.

Softening.—If the thrombus does not become organized, it usually undergoes a process of softening and liquefaction. This is most common in stratified thrombi, and especially in those occurring in the veins and heart. It is probable also that in some cases the thrombus may become completely absorbed; a partial absorption is quite frequent. The softening usually commences in the centre of the clot, and gradually extends towards the circumference. The thrombus breaks down into a soft pulpy material, which sometimes has the appearance of pounded cooked meat, and in other cases to the naked eye looks much like pus. Under the microscope, however, it is seen to consist merely of albuminous granules, molecular fat, and more or less altered red and white blood-corpuscles. These changes in the clot are owing simply to its disintegration; but it is probable that in some quite exceptional cases the thrombus may suppurate, and that owing to the proliferation of the white blood-corpuscles which it contains it may become converted into true pus. Pus, it must be remembered, may also enter the softened thrombus from without. The whole of the thrombus may thus become softened, or the process may be limited to the more central portions, whilst the external layers become organized. Very frequently, as the older portions of the clot are becoming disintegrated and softened, fresh coagulation takes place at its extremities.

Thrombi may also become perforated by openings through which the circulation is re-established. This may occur in organized or partially softened thrombi, and constitutes what is known as *canalization* of the thrombus.

CAUSES.—The causes of thrombosis are of two kinds—those which lead to a retardation of the blood-flow, and those in which there is some abnormal condition of the walls of the vessels or of the blood.

1. *Thrombosis from Retardation of the Blood-flow* may result from—

a. Interruption or narrowing of the vessel.—This occurs after the application of a ligature. Coagulation commences at the point of contact, and extends as far as the first large collateral branches, thus permanently closing the vessel. The laceration of the inner coat of the artery by the ligature, and the folds into which it is thrown, contributes very materially to the adhesion of the clot. The pressure exercised by tumors, cicatricial tissue, extravasations of blood, and the closure of a vessel by the impaction of an embolus, may in the same way by impeding or arresting the circulation cause thrombosis. General obstruction in the capillaries of a part also causes coagulation in the adjacent veins.

β. Solution of the continuity of the vessel.—The formation of a thrombus after the division or tearing of a vessel constitutes the means by which hemorrhage is immediately arrested—there must be either thrombosis or continuous hemorrhage. In the arteries, the severed end of the vessel contracts and retracts within its sheath, coagulation commences around it and extends upwards as far as the first large collateral branch. In the veins hemorrhage is frequently arrested by the valves, and the formation of a thrombus will evidently depend upon the relative situations of the valves and collateral vessels. The hemorrhage from the uterus after the separation of the placenta is arrested either by uterine contraction or by thrombosis.

γ. Dilatation of the vessels, or of the heart.—The most familiar example of thrombosis from this cause is that which occurs in an aneurism. The greater the amount of dilatation the greater is the retardation of the blood-flow. The coagulation commences at the sides of the vessel, and may extend until it completely fills the cavity. Coagulation from the same cause is not uncommon in the dilated plexuses of the prostate gland. In the heart, thrombosis is most frequent in the auricles. It usually commences in the auricular appendix, where there is very little propulsive power, and it may gradually extend into the auricular cavity. It is also met with in the ventricles, commonly commencing here between the columnæ carneæ.

δ. Diminished cardiac power.—This is a common cause of thrombosis in the veins. The coagulation commences just behind the flaps

of the valves, from which it gradually extends into the cavity of the vessel. This appears to be owing to the force of the current not being sufficiently strong to completely open the valves, and the blood consequently stagnates and coagulates behind them. The crural and iliac veins, the venous plexuses of the back, and the cerebral sinuses are the situations in which thrombosis from this cause is most frequently met with. It occurs in the course of many chronic exhausting diseases in which the cardiac power becomes diminished, and is especially frequent in phthisis, cancer, etc. This state of the blood, which often has an abnormal tendency to coagulate, together with the quiescent condition of the patient, materially aids in causing the coagulation.

2. *Thrombosis from Abnormal Conditions of the Vessels or of the Blood.*

a. *Causes in the vessels.*—Any abnormal condition of the walls of a vessel may be a cause of thrombosis. When the wall becomes abnormal it acts as a foreign body and the blood coagulates upon it, and it may continue to do so until the cavity of the vessel becomes filled with coagulum. Thrombi produced in this way are consequently stratified. The walls of a vessel may become altered as the result of inflammatory processes, and inflammation was formerly regarded as the main, if not the only, cause of thrombosis; hence thrombosis in veins is frequently termed “phlebitis” even at the present day. Inflammation of veins, as already stated, is certainly rare as a *primary* condition, although it not unfrequently results from the formation of a thrombus. When occurring primarily, inflammatory processes, both in the arteries and the veins, have their seat in the external and middle coats or in the deeper layers of the intima. They never commence in the lining membrane of the vessel: this only becomes affected secondarily. The vitality of the lining membrane becomes impaired as the result of the inflammatory process, and when this has occurred it acts as a foreign body, and thus there is a tendency for the blood to coagulate upon its surface. In other cases the lining membrane is completely destroyed, and the subjacent diseased tissues thus come into contact with the circulating blood, and in the same way cause the formation of a thrombus. Such inflammatory changes occur in the arteries, constituting the condition known as “atheroma,” which, in the smaller vessels, may be a cause of thrombosis. In the heart they constitute endocarditis; and here also, as has been seen, coagulation may take place upon the abnormal surface of the inflammatory vegetations. (See “Endocarditis.”)

The walls of a vessel may also become altered, and thus thrombosis result, from inflammation or gangrene of the tissues in which it is situated. The vitality of the vessel becomes destroyed and the blood coagulates within it; and by this means the occurrence of hemorrhage is frequently prevented. The projection of new formations, as sarcoma, into the cavity of vessels, may in the same way cause the formation of a thrombus.

β. Causes in the blood.—There can be no doubt that certain conditions of the blood favor coagulation, and consequently tend to promote the occurrence of thrombosis. The phenomenon of blood-coagulation has been shown by Professor A. Schmidt to depend upon the white blood-corpuscles, and it is well known that, under certain circumstances, an increase in the number of these corpuscles in the blood favors the formation of thrombi. Such an increase is frequently met with in inflammatory processes which involve largely the lymphatic organs (leucocytosis). It is probable that other conditions of the blood, such as a loss of its serum, and alterations produced by fevers and by septic poisons, may also favor coagulation. (See “Septicæmia.”) An increased tendency of the blood to coagulate has long been known to exist in many acute inflammatory diseases, and in the latter months of pregnancy.

An increased tendency of the blood to coagulate, to whatever circumstances it may be due, is, however, probably never sufficient in itself to determine the formation of a thrombus; it can hence only be regarded as a predisposing cause. It is especially in those conditions in which the circulation is impeded from diminished cardiac power, that it becomes an important agent in producing thrombosis.

RESULTS.—The results of thrombosis comprise certain changes in the walls of the vessels, more or less obstruction to the circulation, and embolism. These must be considered separately.

1. *Changes in the vessels.*—More or less alteration in the wall of the vessel is an invariable consequence of the formation of a thrombus. When the thrombus undergoes a process of organization, it becomes, as already described, intimately united with the vascular wall. The latter in the first place becomes infiltrated with cells and considerably thickened, but ultimately, together with the thrombus, gradually atrophies. It is when the thrombus softens and becomes disintegrated that the most important changes take place in the vessel. These changes are of an acute inflammatory nature, and appear to result from the irritating influence of the softened thrombus. They are

most frequently observed in the veins, where softening is most liable to occur.

The walls of a vein within which a thrombus is softening are considerably thickened, so that the vessel more resembles an artery. The inner surface has lost its translucency, and is of a dead opaque color. The vasa vasorum are hyperæmic. Under the microscope, the cells of the intima and of the middle and external coats are found to be considerably increased in number, and numerous leucocytes are seen infiltrating the different textures. In some cases small collections of pus are seen in the external coats. Similar changes are observed in the arteries.

In these acute inflammatory changes which are produced in veins by thrombi, proliferation of the endothelium of the vessel often appears to play a prominent part. As already stated, in inflammation of vessels not resulting from thrombosis, the endothelium undergoes no active changes.

2. *Obstruction to the circulation.*—The consequences of the obstruction to the circulation which results from the formation of a thrombus will depend upon the rapidity of its formation, the nature and size of the vessel obstructed, the situation and number of the collateral branches, and the force of the circulating current. When a thrombus forms in a vein of small size and there are numerous collateral vessels, as in the prostatic or uterine plexuses, the circulation is but little interfered with, and no symptoms of obstruction result. If, however, the main trunk of a large vein becomes obliterated, as that of the femoral or iliac veins, the obstruction is followed by hyperæmia, the extent and duration of which will depend upon the facility with which the circulation can be restored by the collateral vessels. Thrombosis in the above-named veins frequently occurs, as already stated, in the latter stages of many chronic debilitating diseases, especially in phthisis; also in the puerperal state, where it gives rise to the condition known as *phlegmasia dolens*. The formation of a thrombus here is followed by œdema and swelling of the limb, which becomes tense, elastic, and painful. In the early stage there may be some cyanosis, but this is usually quickly followed by a pallid whiteness of the surface. There is often more or less tenderness in the course of the vein, which feels enlarged, hard, and knotted, owing to the secondary inflammatory changes in its walls. At the same time there is sometimes swelling and tenderness of the lymphatics, which may be seen as red lines traversing the limb. Diffuse inflammation of the

skin and subcutaneous cellular tissue occasionally occurs. These changes are owing partly to the mechanical impediment to the circulation, and partly to the obstruction of lymphatics and to the secondary inflammatory processes in the vein and tissues which ensue. The circulation is usually ultimately restored; but if the impediment has been of long duration, the tissues become thickened, and the limb is left in a hard, indurated, and somewhat enlarged condition.

The formation of a thrombus in an artery is followed in the first place by anæmia of the parts supplied by it; the ultimate result will depend upon the facility with which the circulation can be restored by the collateral vessels. If the circulation be quickly re-established, as is usually the case, the vitality of the tissues may not become impaired; but if not, the part may undergo a process of molecular disintegration and softening, the softened tissue often being surrounded by a zone of hyperæmia which results from the attempt to establish a collateral circulation.

3. *Embolism*.—Portions of the thrombus may be carried away by the circulation, thus constituting embolism. This, which is the most important result of thrombosis, will be considered in the following chapter.

CHAPTER XLV.

EMBOLISM.

EMBOLISM is the arrest of solid substances circulating in the blood in vessels which are too small to allow them to pass. The solid substances are termed *emboli*. These are very various in their nature.

By far the most frequent source of emboli are thrombi, portions of which are carried from the seat of their formation by the circulation, and become arrested in distant vessels—thus constituting embolism. A thrombus may give rise to emboli in various ways. It may soften and break down, and if the lumen of the vessel be thus restored, its fragments become distributed by the blood-current. In those cases in which the thrombus does not fill the vessel, portions of it may readily be carried away by the blood passing over it. Perhaps, however, the

most frequent way in which a thrombus gives rise to embolism is by its conical end being broken off by the current of blood from a collateral vessel. The formation of a thrombus, as already described, usually ceases opposite the entrance of a large collateral vessel, and if its conical end project a little way into the cavity of this vessel it may be readily broken off by the blood-current. (Fig. 129.) It is especially venous thrombi which give rise to embolism: the veins of

the leg, the iliac, hypogastric, and jugular veins being amongst the most common sources. Emboli from cardiac thrombi are also exceedingly common, whilst those from arterial are the least frequent. In embolism originating in thrombosis some sudden movement or exertion often determines the separation of the embolus.

Emboli may, however, originate independently of thrombi; vegetations and calcareous or atheromatous masses separated from the valves of the heart, or from the inner surface of arteries; portions of new growths, as carcinoma, which, having perforated the vessels, have been carried away by the current; parasites which have made their way into the interior of vessels; pigment granules, and other substances, may all constitute emboli.

The emboli become arrested in the first vessels they meet with which are too small to allow them to pass: the size of the vessel will consequently depend upon the size of the embolus. They are often so minute that they pass into and become impacted in the smallest capillaries. The seat of impaction is usually at the bifurcation of the vessel, or where, from the giving off of branches, the calibre is diminishing rapidly. (See Fig. 130.) Thus emboli originating in the systemic veins or in the right cardiac cavities will most commonly become arrested in the vessels of the lungs; those originating in the arteries, the left cardiac cavities, or the pulmonary veins—in the systemic arteries and capillaries, especially in those of the spleen, kidneys, and brain; and those originating in the portal venous system—in the hepatic branches of the portal vein. In some cases, however, the smallest emboli may pass through the capil-

FIG. 129.



A Thrombus in the Saphenous Vein.—Showing the projection of the conical end of the thrombus into the femoral vessel. *S.* Saphenous vein. *T.* Thrombus. *C.* Conical end projecting into femoral vein. At *v v*, opposite the valves, the thrombus is softened. (Virchow.)


laries of the lungs and become arrested in those of the kidneys, spleen, or other organs. Thus, with the exception of emboli originating in the portal vessels, the seat of arrest is the arteries or capillaries.

The emboli are usually carried in the direction of the main current; hence those carried by the aortic stream more commonly pass into the thoracic aorta than into the carotid and subclavian vessels, and into the left carotid and renal artery than into the corresponding arteries of the opposite side. Gravitation also influences the direction in which they are carried, especially those of large size, which move somewhat more slowly than the blood-stream. Owing to this they are more common in the lower lobes and posterior parts of the lungs than in the superior and anterior portions of these organs.

The embolus, when arrested, may either completely or only partially fill the cavity of the vessel. If, as is frequently the case, the arrest takes place at a point of bifurcation, the embolus may partially fill both branches, allowing a small stream of blood to pass. This may break off portions of it, and so cause secondary emboli, which becomes impacted in more distant vessels. The amount of obstruction which immediately follows the arrest will partly depend upon the nature of the embolus itself. If the embolus be from a soft, recently formed thrombus, it will adapt itself to the cavity of the vessel, and so completely occlude it. If, on the other hand, it is irregular in shape and firm in consistence, as when derived from a calcified cardiac vegetation, it may not fill the vessel, but allow a small current of blood to pass it.

The arrest of the embolus and the consequent obstruction to the circulation, is followed by the formation of thrombi behind and in front of it, which extend as far as the entrance of the first large collateral vessels. (Fig. 130.) If the embolus does not completely fill the vessel, coagulum is deposited in successive layers upon its surface until the occlusion of the vessel is complete, and then the secondary thrombus extends, as in the former case, until it meets with a current of blood strong enough to arrest its progress. If the embolus is a portion of a soft thrombus, it will in most cases be im-

FIG. 130.



Embolus impacted at the Bifurcation of a Branch of the Pulmonary Artery.—Showing the formation of thrombi behind and in front of it, and the extension of these as far as the entrance of the next collateral vessels. E. Embolus. t t'. Thrombi. (Virchow.)

possible to distinguish it from the secondary thrombus which surrounds it. If, however, it is a calcareous mass, or a portion of an old thrombus, it may usually be distinguished from the more recent secondary coagulum.

Emboli may, in rare cases, become absorbed. They may also, when derived from thrombi, become organized or soften. The changes in the secondary thrombi are similar to those already described as occurring in the primary—comprising adhesions to the wall of the vessel, softening, and organization.

RESULTS.—The results of embolism are of two kinds, those depending upon the mechanical obstruction to the circulation, and those produced by the irritating or infective properties of the emboli themselves. (See “Pyæmia.”)

Changes in the Obstructed Vessels.—The first series of changes are those occurring in the walls of the vessel within which the embolus becomes arrested. These changes depend upon the mechanical and chemical properties of the embolus. If the embolus possesses no infective properties, being derived from a source where no putrefactive changes are taking place, it, together with the thrombus which it causes to form around and beyond it, usually becomes organized or reabsorbed, and the walls of the vessel become more or less thickened. Or, if the embolus has rough surfaces, as when derived from a calcareous vegetation in the heart, it may produce some inflammation of the vessel. When, on the other hand, the embolus is impregnated with unhealthy pus or with other putrid inflammatory products, it is very liable to cause gangrenous inflammation and sloughing of the walls of the vessel within which it is impacted, which may extend for some distance into the surrounding tissues.

The vessels also undergo important changes, owing to the arrest of the circulation. These changes have been investigated by Cohnheim, and have already been alluded to in the chapter on “Inflammation.” Cohnheim found that the exclusion of the blood from a vascular region for a sufficient length of time produced so much impairment of the vitality of the vascular walls, that when the blood was again allowed to circulate, the vessels dilated, the blood-flow was retarded, and the liquor sanguinis and red and white blood-corpuscles escaped into the surrounding tissues. An arrest of the circulation for a still longer period was followed by capillary hemorrhages, and ultimately by necrosis of the vessel.

Allusion must be made here to embolism as a cause of aneurism.

That aneurisms, especially of the cerebral arteries in young people, are often due to embolism is now pretty generally admitted by pathologists. With regard to the mode in which the embolus causes dilatation of the artery, although this may differ in different cases, an injurious influence of the embolus upon the walls of the artery, and a consequent inflammatory softening of the vessel, is probably the most common condition.

Changes in the Organ.—Important changes take place in the organ or tissue, the vessels of which have become plugged by the embolus. The first effect of the plugging of a vessel by an embolus, is the arrest of the circulation through it, and if the vessel be the main nutrient or functional artery, this is followed by the sudden cessation of the function and nutrition of the part. Thus, plugging of one of the larger arteries in the brain is followed by sudden loss of consciousness and paralysis (apoplexy); plugging of the pulmonary artery, by sudden asphyxia; and of the coronary arteries, by sudden paralysis of the heart. The subsequent changes will depend upon the structure of the organ, the arrangement of its vessels, the facility with which a collateral circulation can be established, and the nature of the embolus. If the circulation be quickly restored by the collateral vessels, a sufficient arterial anastomosis being established behind the obstructed artery, the part recovers itself without undergoing any structural change, and its nutrition and functions are restored. If, however, this is not the case, the walls of the obstructed vessels become so much damaged that exudation and emigration take place, and the vessels and surrounding tissues may ultimately lose their vitality. The interference with the circulation is partly due to the thrombosis which occurs around the impacted embolus. The organs in which the arrangement of the bloodvessels is such that the circulation cannot be readily restored by the anastomosing vessels, and consequently those in which necrotic changes most commonly result from embolism, are the spleen, the kidneys, the lungs, the brain, and the retina. Those arteries with which anastomoses do not exist are called by Cohnheim "terminal" arteries.

The area of tissue from which the supply of blood has been cut off is usually surrounded by a zone of intense hyperæmia, which results from the stress which is thrown upon the collateral vessels. This zone of hyperæmia is very characteristic, and indicates at once the nature of the lesion. When, owing to the absence of sufficient arterial anastomosis (when terminal arteries are blocked), the circulation can-

not be restored, the vessels in the embolic area become so much altered that exudation, emigration, and hemorrhage take place, and thus is produced a tract of tissue within which the circulation has become arrested, and which is more or less extensively infiltrated with blood. This is known as a *hemorrhagic infarct*.

These hemorrhagic infarcts are very frequently met with, especially in the lungs, spleen, and kidneys. They are blackish-red, firm masses of consolidation, and owing to the distribution of the bloodvessels they are usually wedge-shaped, the apex of the cone being towards the

centre of the organ. The more complete the obstruction and the more vascular the tissue, the greater is the amount of infarction, and the more rapid is the softening and disintegration that ensues. (Fig. 131.)




Diagram of a Hemorrhagic Infarct.—*a*. Artery obliterated by an embolus (*e*). *v*. Vein filled with a secondary thrombus (*th*). 1. Centre of infarct which is becoming disintegrated. 2. Area of extravasation. 3. Area of collateral hyperæmia. (O. Weber.)

This vascular engorgement of the embolic area was formerly supposed to be due to the increased stress which is thrown on the collateral vessels. The investigations of Cohnheim, however, show that it is really mainly owing to the impairment of the vitality of the walls of the bloodvessels, and the consequent exudation, emigration, and ultimate necrosis of the vascular walls. When the force of the blood-stream in the artery is annihilated by the impaction of the embolus there is a backward pressure and regurgitation from the veins into the

capillaries, so that there is produced considerable venous engorgement of the last-named vessels. There is thus a substitution in the capillaries and small arteries of venous for arterial blood, and owing to this the vitality of these vessels becomes impaired, and hemorrhage results. The infarction consequently does not occur immediately after the impaction of the embolus, but only after the lapse of a certain time.

The subsequent changes which take place in the infarct depend upon its size, upon the extent to which the circulation in it is interfered with, and upon the nature of the embolus which caused the infarction. If the infarct is small and the embolus possesses no infective properties, the coagulated blood may gradually become decolorized,

and the mass undergo a gradual process of absorption. The infarct then changes from a dark red to a brown or yellow tint, its more external portions become organized into connective tissue, and the whole gradually contracts, until ultimately a cicatrix may be all that remains to indicate the change. If, however, the infarction is considerable, the molecular disintegration and softening may be so extensive as to convert the mass into a pulpy granular material. This may subsequently dry up and become encapsuled. In all these secondary changes which take place in the infarct, its most external portions are surrounded by a red zone of hyperæmic tissue. This is exceedingly characteristic.

If an embolus possesses irritating or infective properties, as when it is derived from a part where putrefactive inflammatory changes are going on, it sets up inflammatory processes both in the vessel within which it becomes impacted, and also in the surrounding tissues. These inflammatory changes frequently lead to the formation of abscesses, which are known as *embolic abscesses*. (See "Pyæmia.") The formation of such abscesses may be associated with more or less infarction of the embolic area.

Thrombosis and Embolism of the Brain.

Thrombosis and embolism are the most common causes of *cerebral softenings*.

Softenings from Thrombosis.—This is commonly the result of atheromatous or calcareous changes in the cerebral arteries. Such changes cause a diminution in the lumen or a roughening of the lining membrane of the vessels, impair their elasticity and contractility, and so favor the occurrence of thrombosis. As a result of the interference with the supply of blood, the cerebral substance undergoes a more or less rapid process of fatty degeneration and softening, such as has been already described. (See "Fatty Degeneration of the Brain.") The softened portions when recent, and when the obstruction is rapidly induced, are often of a reddish color, although with age they gradually become decolorized. In the more gradually induced obstructions the color of the softened tissue is usually white.

Softening from Embolism.—The softening resulting from embolism is, for the most part, entirely dependent upon the obstruction to the circulation caused by the embolus and by the resulting thrombosis.

It is rapidly induced and is often attended by the extravasation of blood, when it constitutes one form of acute red softening. If the interference with the circulation be slight, there may be no extravasation of blood, and the process of disintegration may be more gradual, so that the softened portions are white in color, and the condition then more resembles the chronic white softening already described as resulting from degeneration of the cerebral bloodvessels. (See "Fatty Degeneration of the Brain.") The softened tissue will also be white in color when one of the large vessels is obstructed, so that a large portion of one hemisphere loses its vitality. The vessel most frequently blocked is the middle cerebral artery, in some part of its course; and in the majority of cases it is that of the left side. In almost all cases in which softening of the cerebral substance results from embolism, the embolus is arrested in one of the vessels beyond the circle of Willis, because here the circulation cannot be readily restored by the collateral vessels.

When the interference with the circulation is attended by vascular engorgement and extravasation of blood, the softened portion, in the early stage, is either of a uniform dark-red color, or presents numerous red hemorrhagic points. The softening is most marked in the centre, whilst the hyperæmia and redness may extend for some distance around it. Under the microscope, the softened portion is seen to consist of broken-down nerve-fibres, altered blood-corpuscles, granules of fat, and the large granular corpuscles which result from the fatty degeneration of the nerve-cells and of the cells of the neuroglia. (See Fig. 11.) The surrounding capillaries are dilated and filled with coagula, and the granular corpuscles envelop their walls. In a more advanced stage all trace of nervous structure is lost, the softened mass becomes decolorized, and passes from a dark-red color to a chocolate, brown, yellow, or even white. It may liquefy and form a cyst; more commonly, however, it gradually dries up, and a process of repair takes place by the growth of the surrounding neuroglia, which forms a fibrous network in the place of the softened tissue. This contracts, and ultimately a cicatrix with hæmatoidin crystals may be all that remains.

Red softening from embolism is often very difficult to distinguish in the post-mortem room from that which results from thrombosis.

CHAPTER XLVI.

LEUKÆMIA.

LEUKÆMIA, or leucocythæmia, is a disease characterized by a large increase in the number of the white corpuscles in the blood, which increase is associated with a new formation of lymphatic tissue in the spleen, in the lymphatic glands, and sometimes in other organs. Respecting the relation which subsists between the new growth of lymphatic tissue and the increase in the number of the white blood-corpuscles it is probable, as stated by Virchow, that the former is a cause of the latter, and that the increase is mainly due to a larger number of lymphatic elements entering the blood than in health. At the same time it is difficult to admit this as the only cause of the increase, as new growths of lymphatic tissue may take place in these organs without the production of any leukæmia. This has been seen to occur in Hodgkin's disease. It is possible that the power of the white blood-corpuscles and lymph-corpuscles to form red corpuscles is diminished, or that the white corpuscles may increase by multiplication in the blood. The pathology of the disease, however, is still obscure.

The new formation of lymphatic tissue occurs in the spleen or in the lymphatic glands; and subsequently the liver, the kidneys, the intestines, the lungs, and other organs may become infiltrated with lymphatic elements. The growth is a *continuous* one, the new elements many of them enter the blood, the remainder, not undergoing any retrogressive changes, give rise to a general increase in the size of the organs in which they are generated.

As the new growth may occur in the spleen or in the lymphatic glands, Virchow has distinguished two forms of leukæmia—the splenic and the lymphatic. In the former the white blood-cells are somewhat larger and more granular than normal (Fig. 132); whilst in the latter they are smaller. In most cases of the disease, however, the spleen is the organ principally involved. Often the spleen and the lymphatic glands are simultaneously affected. A purely lymphatic leukæmia is exceedingly rare. The coagulating power of the blood in leukæmia

FIG. 132.



Leukæmic Blood.—From a young man aged twenty-four, with enormous enlargement of the spleen. $\times 260$.

is much diminished, and when the liquid is allowed to stand the white corpuscles form a creamy layer upon its surface.

In the *spleen*—which must thus be regarded as much the most important organ in the production of leukæmia—the change is characterized mainly by increased growth. The organ becomes enlarged, and often enormously so. Its cut surface often presents an almost natural appearance. The trabeculæ, however, are

usually thickened, so as to appear as white lines scattered through the organ; and the Malpighian corpuscles are commonly rather large and prominent. The capsule is in many cases natural, in others more or less thickened. Sometimes the cut surface is mottled with pale wedge-shaped patches which appear to be of embolic origin. Under the microscope the splenic tissue is seen to be but little altered, the principal change consisting in an over-fulness of the bloodvessels with leukæmic blood.

In the *lymphatic glands* the process is similar to that in the spleen, and the glands become enlarged and vascular.

In some cases this excessive development of lymphatic structures takes place in other parts. The follicles of the intestine, and the medulla of bone, are those most commonly involved. In the intestines, the follicles may become so much enlarged as to form distinct firm tumors; and the medulla of bone is, according to Neumann, in exceptional cases the principal seat of the diseased process.

In the course of the disease a new growth of lymphatic tissue or an infiltration with lymphatic elements usually takes place in non-lymphatic structures, principally in the liver and kidneys, less frequently in the lungs and muscle. The new growth in these organs sometimes forms distinct tumors, but much more commonly exists as an infiltration. How far these lymphoid growths are the result of a hyperplasia of the cells in the interstitial tissue of the organ in which they are situated, and how far an emigration of the leucocytes, which exist in such large numbers in the blood, takes part in their formation, is unknown. The latter, however, is probably the most important factor in the process.

The organ which is the most common seat of these lymphoid growths is the *liver*. Here the vessels throughout the organ become enlarged and distended with white blood, and the infiltration commencing in

the interlobular tissue, gradually extends into the lobules themselves. In the earlier stages, the lobules are seen to be clearly mapped out by a grayish-white interlobular infiltration. As this increases the liver-cells become compressed and atrophy, until ultimately the lobules may be entirely replaced by it. Associated with this infiltration there is often a formation of small, round, whitish nodules, somewhat resembling gray tubercles. Owing to these changes, the liver becomes very considerably increased in size.

In the *kidneys*, which are also frequently affected, the change is similar to that in the liver. Here also it consists for the most part in an infiltration, with which may be associated the formation of roundish nodules and masses.

CHAPTER XLVII.

THE PREPARATION AND MOUNTING OF SPECIMENS.

IN this, the concluding chapter, it is proposed to describe very briefly the more common methods of preparing and mounting specimens for microscopical examination. It is not intended to give an account of those various and complex methods of investigation which are required for minute histological research, but merely to describe those more easy ones which suffice in the great majority of cases for the recognition of the structural alterations met with in disease. In order to be as brief as possible, those methods only will be described which I believe to be the most readily available and at the same time to yield the most satisfactory results. The subject will be treated under the following heads: 1st. *The Examination of Specimens whilst fresh*; 2d. *The Hardening of Specimens*; 3d. *The Making of Sections*; 4th. *Staining*; 5th. *Preservation and Mounting*.

The Examination of Specimens whilst Fresh.

Very much may be learned from the examination of specimens in their fresh state before they have been submitted to any hardening process. The examination may be conducted either by making thin

sections of the tissue with a Valentin's knife, or by cutting off a minute portion and tearing it to pieces with fine needles. The specimen prepared in either of these ways should be examined in glycerine. Thin membranous structures may be examined by simply spreading them out in a drop of water placed upon the glass slide, and then covering them with glycerine.

The method of making sections of organs whilst fresh with a Valentin's knife is an exceedingly easy and useful way of at once ascertaining in the post-mortem room the existence of the more marked structural changes. The knife should be well wetted with water, and then drawn *rapidly* through the substance of the organ. The section thus obtained is gently washed in water before being examined in glycerine. A rough examination of the liver, kidneys, and lungs is in this way readily made.

The muscular fibres of the heart and tumors may be examined by the process of teasing. This process, however, is usually more readily effected after the specimen has been kept in Müller's fluid or in a solution of bichromate of potash in the manner to be hereafter described. In the examination of tumors, much may be learned by gently scraping the freshly-cut surface with a knife and examining in glycerine the elements which are thus separated.

The Hardening of Specimens.

In order to make a complete and satisfactory examination of diseased structures, it is in almost all cases necessary to submit them to a process of hardening. By this process they are so altered that fine sections of them can be made, and the sections are also rendered more fit for permanent preservation than are those obtained from tissues in their fresh state. Numerous substances have been employed as hardening agents. Those, however, which are the most generally valuable are *Müller's fluid*, *chromic acid*, *bichromate of potash*, and *alcohol*.

Müller's Fluid.—Of all hardening agents this is much the most satisfactory and the one most generally applicable. It is prepared in the following way:—

Take of —

Potassium Bichromate	2 parts.
Sodium Sulphate	1 part.
Water	100 parts.
Dissolve.	

The tissue should be kept in this solution for from one to two weeks, and then placed in common alcohol for two or three days, after which it is ready for making sections.

The advantages of Müller's fluid are that it alters the macroscopical characters of tissues less than other hardening agents, and its penetrating powers are so great that it is not necessary to cut the tissue up into small pieces, but the whole organ may be placed in it. The tissue may be kept in it longer without injury than in other aqueous solutions.

Müller's fluid is also valuable for the maceration of tissues which are to be examined by teasing—as tumors, muscle, etc. Its action tends to facilitate the isolation of the elements.

Chromic Acid.—This is much less generally applicable than the preceding. It should be used in aqueous solution—strength one-sixth per cent. It is important to employ the solution in large quantities—from six to ten fluidounces for each specimen. The tissue must be cut into small pieces—not much larger than a hazel-nut—before it is placed in the solution. The hardening is usually completed in from five to eight days. It is hastened by renewing the solution. If kept too long in the solution, the tissue becomes brittle. All tissues which have been hardened in chromic acid should be placed in common alcohol for about twenty-four hours before sections are made. Chromic acid is preferable for hardening skin and mucous membranes.

Bichromate of Potash.—This agent is more gradual in its action than chromic acid, and produces less shrinking of the tissue. It is especially valuable for the hardening of kidney, bloodvessels, nerve-tissue, ovary, and some other structures. It is often advisable to commence the hardening with bichromate of potash, and then to complete it in chromic acid or common alcohol. The bichromate is also valuable for the maceration of tissues which are to be examined by teasing. The bichromate should be used in aqueous solution—strength one to two per cent. It penetrates more readily than the chromic acid, and consequently it is not necessary to use quite such small pieces of the tissue. The tissue must be kept in the solution from two to four weeks.

Alcohol.—This is most valuable as an agent for the completion of the hardening of tissues which have been previously placed in Müller's fluid, chromic acid, or in bichromate of potash. The tissues should be kept in it one or two days. Some tissues, however, appear to harden better if they are at once placed in alcohol; amongst these are

the lymphatic glands, spleen, and testis. Common (methylic) alcohol answers for all these purposes. Lastly, alcohol must be employed for the hardening of all tissues which have been injected. Absolute alcohol is often necessary for the completion of hardening of many tissues.

The Softening of Tissues.—Some tissues, as osseous structures, tumors which have become calcified, etc., require to be softened before sections can be made. This may be effected by placing them in the one-sixth per cent. chromic acid solution, to each six ounces of which three or more drops of concentrated hydrochloric acid have been added.

The Making of Sections.

Although various instruments have been contrived for making sections, the one which answers best is a common razor, or a similar blade firmly fixed in a wooden handle. The usefulness of this instrument is much increased if the lower surface of the blade be ground perfectly flat, whilst the upper is made slightly concave.

Hardened tissues from which sections are to be made must always have first been placed in common alcohol, in the manner already described. In making the section the blade of the instrument must be kept well wetted with common alcohol, and the section when cut must be placed at once in the same liquid.

Imbedding.—Hardened tissues which are too small or too delicate to be held in the hand for the purpose of making sections, must be imbedded. The object of imbedding is to fix the tissue in some substance which can not only be readily held, but which can also be easily cut with the section-knife. Various substances are used for this purpose. Of these the one which is most generally applicable is a mixture of wax and olive oil. Equal quantities of pure white wax and olive oil should be warmed together and well mixed, and the mass when cold may be kept to be used as required. When a specimen is to be imbedded, a small case of paper should be made somewhat larger than the specimen. The edges of the paper can be secured with a little mucilage or with a pin. The mixture of wax and oil is then melted and the paper case filled with it. The specimen to be imbedded—which must have been removed from common alcohol—is now to be slightly dried on blotting-paper and immersed in the melted compound. The process of immersion requires a little care. The specimen should be placed upon the point of a fine needle, and as soon as the wax

commences to solidify at the sides of the case, it must be dipped beneath the melted liquid so as to be completely covered by it. The needle must then be removed by giving it a gentle twist so as not to disturb the specimen. When the wax is cool the paper can be removed, and the mass placed in common alcohol. Sections can be made from it when convenient.

Staining.

When the section has been made and placed in common alcohol, in the manner described, it is usually advisable to stain it. The object of staining is to impart to the different portions of the tissue different shades of color, and so that its structure can be more readily recognized. The substances which are the most generally applicable for this purpose are logwood and carmine. Of these logwood is the preferable, inasmuch as it not only differentiates most tissues more completely, but it is a much more agreeable color to work with.

Staining with Logwood.—The following is the method for preparing the logwood solution recommended by Dr. Klein:—

Take of—

Extract. Hæmatox.	6 grammes.
Alumen	18 grammes.
Mix thoroughly in a mortar.	

Add gradually whilst stirring—

Distilled Water	28 cub. cent.
	Filter.

To the filtrate add—

Spirit. Rect.	1 drachm.
-----------------------	-----------

The solution must be kept in a stoppered bottle for a few days before being used.

For staining, several (5–10) drops are to be added to half a watch-glass of distilled water. The thus diluted solution should be filtered before being used. Sections which have been removed from common alcohol, are to be placed in the dilute solution for from ten to twenty minutes, and then to be washed in distilled water.

Gradual Staining with Weak Carmine.—In staining with carmine two methods may be employed: the one consists in using a strong solution, so as to stain rapidly; the other in staining very gradually

with a much less concentrated liquid. Of these, the method of gradual staining with weak solutions yields the most satisfactory results, inasmuch as by this means the differentiation of the tissue is rendered more complete than when the coloration is rapidly induced. The process of rapid staining with strong carmine is, however, the most convenient, and it yields, in most cases, sufficiently satisfactory results.

The following is the method for gradual staining recommended by Dr. Klein:—

Take of—

Powdered Carmine 2 grammes.
Rub it up with a few drops of Distilled Water.

Add—

Liquor Ammoniaë fort. 4 cub. cent.
Distilled Water 48 cub. cent.

Mix together with a glass rod, and filter.
The solution to be kept in a stoppered bottle.

For staining, take one drop of the above solution, and add to it from nine to twelve drops of distilled water. Sections, which have been removed from common alcohol, to be placed in this from sixteen to twenty-four hours.

Rapid Staining with Strong Carmine.—To stain rapidly, the sections may be placed in the above strong solution of carmine without the addition of water. In this case, the strong solution of carmine should have been kept exposed to the air for a sufficient length of time to allow the excess of ammonia to escape. It is consequently advisable to have two bottles of strong carmine solution—one for rapid, and the other for gradual, staining.

The time required for rapid staining is from thirty seconds to five minutes. This, however, will vary very considerably according to the nature of the tissue to be stained. A knowledge of the amount of time which yields the best results will be readily learnt by a little practice.

The sections when removed from the carmine staining solution must be washed for a few seconds in distilled water containing one-quarter per cent. of acetic acid.

Preservation and Mounting.

When the section has been stained, it is ready to be prepared for mounting permanently. The two substances in which tissues are most commonly mounted are glycerine and Dammar varnish. Of these, the latter should be employed in all cases in which the preservation of the section is intended to be permanent. Many tissues, however, yield more satisfactory results when examined in glycerine; but they cannot be preserved in this liquid for any great length of time without undergoing some alteration.

The methods of mounting in Dammar varnish and glycerine must be described separately.

Mounting in Dammar Varnish.—When the section is to be mounted in Dammar, it having been already stained and washed, it must first be placed in *absolute* alcohol for about a quarter of an hour. From the alcohol it must be transferred to oil of cloves, the superfluous alcohol having been first removed from the section by blotting-paper. In the oil of cloves it becomes, in a few seconds, quite transparent. The superfluous oil must be removed by blotting-paper, and the section is then ready for mounting in the Dammar varnish. If more convenient, it may be kept for one or two days in the oil of cloves before it is mounted in the Dammar. In order to make the specimen still more secure, a layer of the varnish may be placed round the edge of the covering glass a few days after the section has been mounted.

The following is the method for preparing the Dammar varnish:—

Take of—

Gum Dammar in powder $\frac{1}{2}$ oz., and dissolve it in Turpentine $1\frac{1}{2}$ oz.

Filter.

Gum Mastic $\frac{1}{2}$ oz., and dissolve it in Chloroform 2 oz. *Filter.*

Mix the two filtered solutions, and again filter.

Mounting in Glycerine.—When the section is to be mounted in glycerine no further preparation is necessary. It is simply washed in distilled water and at once transferred to a drop of strong glycerine upon the slide. The edges of the covering glass must then be fastened down with some cementing liquid. The Dammar varnish answers exceedingly well for this purpose.

PRINCIPAL WORKS REFERRED TO.

- ARNOTT, HENRY.—The Histology and Diagnosis of Cancer.
- BILLROTH, THEODOR.—Die allgemeine chirurgische Pathologie und Therapie.
- BUHL, LUDWIG.—Lungenentzündung, Tuberkulose, und Schwindsucht.
- COHNHEIM.—*Ueber Entzündung und Eiterung.* Virchow's Archiv, vol. xl.
Ueber das Verhalten der fixen Bindegewebskörperchen bei der Entzündung.
 Virchow's Archiv, vol. xlv.
 Neue Untersuchungen über die Entzündung. 1873.
- CORNIL ET RANVIER.—Manuel d'Histologie Pathologique.
- FOERSTER, AUGUST.—Handbuch der pathologischen Anatomie.
- FOX, WILSON.—On the Artificial Production of Tubercle.
- HÜTER, C.—*Die septikämischen und pyämischen Fieber.* Handbuch der allgm. u. spec. Chirurgie. Von Pitha und Billroth.
- JONES AND SIEVEKING.—Manual of Pathological Anatomy. 2d Edit. Edited by Dr. Payne.
- KLEBS.—Handbuch der pathologischen Anatomie.
- KLEIN.—The Anatomy of the Lymphatic System.
- KÜHNE, W.—Lehrbuch der physiologischen Chemie.
- LISTER, J.—*On the Early Stages of Inflammation.* Philosophical Trans., 1859.
- LÜCKE.—*Die Geschwülste.* Handbuch der allgm. u. spec. Chirurgie. Von Pitha und Billroth.
- NIEMEYER, F.—Text-book of Practical Medicine. Translated by Drs. Humphreys and Hackley.
- PAGET, SIR J.—Lectures on Surgical Pathology. Edited by Prof. Turner. Pathological Soc. Lond. Transactions.
- RECKLINGHAUSEN.—*Ueber Eiter und Bindegewebskörperchen.* Virchow's Archiv, vol. xxviii.
- RINDFLEISCH, E.—Lehrbuch der pathologischen Gewebelehre.
- RYNECK.—*Zur Kenntniss der Stase des Blutes in den Gefässen entzündeter Theile.* Rollet's Untersuch. aus dem Institute für Phys. u. Histol. in Graz.
- SANDERSON, J. BURDON.—*The Process of Inflammation.* Holmes's System of Surgery, vol. v.
The Inoculation of Tubercle. Appendix of Reports of Medical Officer to Privy Council, for 1868 and 1869.
Recent Researches on Artificial Tuberculosis in their Relation to the Pathology of Phthisis. 1869.
- SCHÜPPEL, OSCAR.—*Untersuchungen über Lymphdrüsen-Tuberkulose.*

STRICKER, S.—*Various papers by, in his*—Studien aus dem Institute für experimentelle Pathologie in Wien. 1869.

Manual of Human and Comparative Histology, vol. i. Edited by Prof. Stricker; translated by Mr. Power.

UHLE UND WAGNER.—Handbuch der allgemeinen Pathologie.

VIRCHOW, RUDOLF.—Die Cellular Pathologie.

Die krankhaften Geschwülste.

Gesammelte Abhandlungen.

Handbuch der speciellen Pathologie und Therapie. Band I.

WAGNER, E.—Manual of General Pathology. Translated by Drs. John Van Duyn and E. C. Seguin. 6th Edition.

WEBER, O.—*Die Gewebserkrankungen*. Handbuch der allgm. u. spec. Chirurgie. Von Pitha und Billroth.

WILKS AND MOXON.—Lectures on Pathological Anatomy. 2d Edition.

INDEX.

ABSCESS, 174
 in connective tissue, 217
 of bone, 219
 of brain, 260
 of kidneys, 248
 of liver, 245
 of lungs, 266
 of lymphatic glands, 233
 Acute miliary tuberculosis, 181
 Adenomata, 131
 of mammary gland, 132
 of mucous membranes, 134
 Adeno-myxoma, 132
 Adeno-sarcoma, 132
 Adipose tissue, atrophy of, 33
 growth of, 43
 Alimentary canal, amyloid degeneration of, 77
 Amyloid degeneration, 67
 of alimentary canal, 77
 of kidneys, 73
 of liver, 71
 of lymphatic glands, 77
 of spleen, 76
 substance, nature of, 68
 reactions of, 68
 source of, 71
 Angiomata, 128
 Aneurism, from arteritis, 225
 from embolism, 310
 Arteries, atheroma of, 223
 calcification of, 83
 fatty degeneration of, 55
 inflammation of, 223
 in chronic Bright's disease, 259
 in syphilis, 282
 Atheroma, 224
 Atrophy, 33
 causes of, 35
 numerical, 34
 physical characters of, 38
 simple, 33
 of adipose tissue, 34

BACTERIA, 201
 Blood, as cause of disease, 18
 alteration of, in malignant growths, 98

Blood—
 ante-mortem coagulation of. *See* "Thrombosis."
 post-mortem coagulation of, 299
 Blood-corpuscles, as source of new formations, 101
 as source of pus, 175
 emigration of, in inflammation, 167
 exudation of, in mechanical hyperæmia, 295
 Blood-cysts, 146
 Bloodvessels, amyloid degeneration of, 69, 73
 calcification of, 83
 fatty degeneration of, 55
 inflammation of, 223
 Bone, atrophy of, 39
 caries of, 220
 inflammation of, 218
 necrosis of, 220
 sclerosis of, 219
 Brain, abscess of, 260
 embolism in, 313
 fatty degeneration of, 57
 inflammation of, 259
 inflammatory softening of, 260
 softening of, from embolism, 313
 red softening of, 59, 313
 sclerosis of, 261
 thrombosis of, 313
 tubercle of, 190, 181
 white softening of, 58
 yellow softening of, 58
 Brown atrophy of heart, 62
 induration of lungs, 297

CACHEXIA, cancerous, 97
 Calcareous degeneration, 80
 of arteries, 83

Cancer, 147
 adenoid, 159
 bloodvessels of, 149
 cells of, 148
 clinical characters of, 159
 colloid, 154
 development of, 149
 encephaloid, 153
 epithelial, 156

Cancer—

- lymphatics of, 149
- melanotic, 151
- osteoid, 151
- scirrhous, 152
- secondary changes in, 151
- stroma of, 149
- structure of, 148
- varieties of, 151

Capillaries, fatty degeneration of, 56

Caries, 220

Carmine for staining, 322

Cartilage, erosion of, 213

inflammation of, 212

Caseation, 54

of products of scrofulous inflammation, 179

of tubercle, 187

Caseous masses, pathological significance of, 54

Carcinomata, 147

Catarrh, 237

Cell, as seat of nutrition, 20

definition of, 24

limiting membrane of, 22

nucleus of, 22, 23

protoplasm of, 21, 23

Cells, constitution of, 20

genesis of, 23

"indifferent," 102

multiplication of, 24

physiology of, 22

Cell-wall, nature of, 23

Cerebral softenings, 57, 313

Cicatrical tissue, 216

Clots, ante-mortem, 299

post-mortem, 299

Cloudy swelling, 170

Colloid cancer, 154

degeneration, 63

of muscle, 66

Condylomata, 130

Congestion. *See* "Hyperæmia."

Connective tissue, fatty infiltration of, 44

inflammation of, 214

suppuration of, 216

Cornea, inflammation of, 213

Corpora amylacea, 78

Corpuscles, exudation, 52

Croup, 239

Cystic-sarcoma, 137

Cysts, 161

classification of, 163

modes of origin of, 161

secondary changes in, 162

structure of, 162

DAMMAR varnish for mounting, 323

Degeneration, 41

amyloid, 67

calcareous, 80

causes of, 42

Degeneration—

colloid, 64

fatty, 43

mucoid, 63

pigmentary, 84

Diphtheria, 240

Disease, definition of, 17

Dysentery, 241

EMBOLISM, 307

in brain, 313

results of, 310

Emigration of white blood-corpuscles in

inflammation, 167

Emphysema, 40

Encephaloid cancer, 153

Enchondromata, 113

Endocarditis, 226

acute, 228

chronic, 229

ulcerative, 228

Epithelioma, 156

cylindrical, 159

Epulis, 111, 145

Exostoses, 118

Exudation corpuscles, 52

Exudation in inflammation, cause of, 173

of liquor sanguinis in inflammation,

168

in mechanical hyperæmia, 294

FAT, absorption of, 53

source of, in fatty degeneration, 43

infiltration, 43, 47

Fatty degeneration, 43

causes of, 43

of arteries, 55

of brain, 57

of capillaries, 56

of heart, 60

of kidneys, 62

of muscle, 59

infiltration, 47

of connective tissue, 44

of heart, 48

of liver, 49

of muscle, 47

Fibroid induration, as result of inflam-

mation, 177

of mechanical hyperæmia,

295

of syphilis, 206

of heart, 230

Fibromata, 109

Fibroplastic tumor.* *See* "Sarcomata,

spindle-celled."

GANGRENE, 26

causes of, 30

dry, 27

Gangrene—
 evidences of, 26
 moist, 27
 senile, 32

Genesis of cells, 23

Giant cells, 185

Glioma, 142

Gluge, corpuscles of, 52

Gray degeneration, 261

Gummata, 207

HÆMATOIDIN, 85

Hæmorrhagic infarct, 312

Heart, brown atrophy of, 62

changes in pyrexia, 92

fatty degeneration of, 60

infiltration of, 47

fibroid induration of, 230

inflammation of, 226

Heterology, 104

Hodgkin's disease, 124

Homology, 103

Horns, 130

Hyperæmia, 291

active, 291

mechanical, 293

of liver, 295

of lungs, 297

post-mortem appearances of, 298

Hyperplasia, 94

Hypertrophy, 93

INFARCT, 312

Infiltrations, 42

Infiltration, fatty, 47

Inflammation, 164

acute, 177

alterations in nutrition in, 169

as cause of gangrene, 31

causes of blood-stasis in, 172

changes in the bloodvessels and circulation in, 165

chronic, 177

croupous, 239

emigration of red blood-corpuscles in, 168

of white blood-corpuscles in, 167

exudation of liquor sanguinis in, 168

fibrinous, 239

formation of pus in, 175

idiopathic, 178

infective, 178

scrofulous, 179

specific, 178

stasis in, 167

traumatic, 178

varieties of, 176

of arteries, 223

of bloodvessels, 223

of bone, 218

Inflammation—

of brain and spinal cord, 259

of cartilage, 212

of common connective tissue, 214

of cornea, 213

of heart, 226

of kidney, 248

of liver, 245

of lungs, 262

of lymphatic structures, 232

of mucous membranes, 237

of serous membranes, 242

of veins, 226

Inflammatory new formations, nature of, 95

fever. *See* "Surgical Fever."

stasis, 167

Interstitial hepatitis, 245

nephritis, 249, 255

pneumonia, 271

Intestine, amyloid degeneration of, 77

tuberculosis of, 193

typhoid ulceration of, 235

KIDNEY, abscess of, 249

amyloid degeneration of, 73

fatty degeneration of, 62

inflammation of, 248

interstitial nephritis, 255

leukæmic growth in, 317

surgical, 249

scarlatinal nephritis, 254

suppurative nephritis, 249

tubal nephritis, 251

LARDACEOUS degeneration. *See*
"Amyloid Degeneration."

Leukæmia, 125, 315

Leukæmic growths in kidney, 317

in liver, 316

in lymphatic glands, 316

in spleen, 316

Lipomata, 112

Liver, abscess of, 245

acute inflammation of, 245

acute yellow atrophy of, 248

amyloid degeneration of, 71

changes in pyrexia, 91

cirrhosis of, 245

fatty infiltration of, 49

leukæmic growth in, 316

nutmeg, 295

syphilitic growths in, 211

Logwood for staining, 321

Lungs, abscess of, 266

broncho-pneumonia, 267

brown induration of, 297

catarrhal pneumonia, 267

cirrhosis of. *See* "Interstitial Pneumonia."

croupous pneumonia, 262

HENRY C. LEA'S SON & CO.'S

(LATE HENRY C. LEA)

CLASSIFIED CATALOGUE

OF

MEDICAL AND SURGICAL PUBLICATIONS.

In asking the attention of the profession to the works advertised in the following pages, the publishers would state that no pains are spared to secure a continuance of the confidence earned for the publications of the house by their careful selection and accuracy and finish of execution.

The large number of inquiries received from the profession for a finer class of bindings than is usually placed on medical books has induced us to put certain of our standard publications in half Russia, and that the growing taste may be encouraged, the prices have been fixed at so small an advance over the cost of sheep, as to place it within the means of all to possess a library that shall have attractions as well for the eye as for the mind of the reading practitioner.

The printed prices are those at which books can generally be supplied by book-sellers throughout the United States, who can readily procure for their customers any works not kept in stock. Where access to bookstores is not convenient, books will be sent by mail post-paid on receipt of the price, and as the limit of mailable weight has been removed, no difficulty will be experienced in obtaining through the post-office any work in this catalogue. No risks, however, are assumed either on the money or on the books, and no publications but our own are supplied, so that gentlemen will in most cases find it more convenient to deal with the nearest bookseller.

An ILLUSTRATED CATALOGUE, of 64 octavo pages, handsomely printed, will be forwarded by mail, post-paid, on receipt of ten cents.

HENRY C. LEA'S SON & CO.

Nos. 706 and 708 Sansom St., PHILADELPHIA, June, 1880.

INCREASED INDUCEMENT FOR SUBSCRIBERS TO THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES.

**TWO MEDICAL JOURNALS, containing nearly 2000 LARGE PAGES,
Free of Postage, for FIVE DOLLARS Per Annum.**

TERMS FOR 1880.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, published quarterly (1150 pages per annum), with	}	Five Dollars
THE MEDICAL NEWS AND ABSTRACT, monthly (768 pp. per annum),	}	per annum,
		in advance.

SEPARATE SUBSCRIPTIONS TO

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES, when not paid for in advance, Five Dollars.

THE MEDICAL NEWS AND ABSTRACT, free of postage, in advance, Two Dollars, and a half.

* * Advance paying subscribers can obtain at the close of the year cloth covers, gilt-lettered, for each volume of the Journal (two annually), and of the News and Abstract (one annually), free by mail, by remitting ten cents for each cover.

It will thus be seen that for the moderate sum of FIVE DOLLARS in advance, the subscriber will receive, free of postage, the equivalent of three or four large octavo volumes, stored with the choicest matter, original and selected, that can be furnished by the medical literature of both hemispheres. Thus taken together, the "JOURNAL," and the "NEWS AND ABSTRACT" combine the advantages of the elaborate preparation that can be devoted to the Quarterly with the prompt conveyance of intelligence by the Monthly; while the whole being under a single editorial supervision, the subscriber is secured against the duplication of matter inevitable when periodicals from different sources are taken together.

The periodicals thus offered at this unprecedented rate are universally known for

(For "THE OBSTETRICAL JOURNAL," see p. 24).

their high professional standing.

I.

THE AMERICAN JOURNAL OF THE MEDICAL SCIENCES,

EDITED BY I. MINIS HAYS, M.D.,

for more than half a century has maintained its position in the front rank of the medical literature of the world. Cordially supported by the profession of America, it circulates wherever the language is read, and is universally regarded as a national exponent of American medicine—a position to which it is entitled by the distinguished names from every section of the Union which are to be found among its collaborators.* It is issued quarterly, in January, April, July, and October, each number containing about three hundred octavo pages, appropriately illustrated wherever necessary. A large portion of this space is devoted to Original Communications, embracing papers from the most eminent members of the profession throughout the country.

Following this is the REVIEW DEPARTMENT, containing extended reviews by competent writers of prominent new works and topics of the day, together with numerous elaborate Analytical and Bibliographical Notices, giving a fairly complete survey of medical literature.

Then follows the QUARTERLY SUMMARY OF IMPROVEMENTS AND DISCOVERIES IN THE MEDICAL SCIENCES, classified and arranged under different heads, and furnishing a digest of medical progress, abroad and at home.

Thus during the year 1879 the "JOURNAL" contained 67 Original Communications, mostly elaborate in character, 147 Reviews and Bibliographical Notices, and 215 articles in the Quarterly Summaries, illustrated with 70 wood engravings.

That the efforts thus made to maintain the high reputation of the "JOURNAL" are successful, is shown by the position accorded to it in both America and Europe as a leading organ of medical progress:—

This is universally acknowledged as the leading American Journal, and has been conducted by Dr. Hays alone until 1869, when his son was associated with him. We quite agree with the critic, that this journal is second to none in the language, and cheerfully accord to it the first place, for nowhere shall we find more able and more impartial criticism, and nowhere such a repertory of able original articles. Indeed, now that the "British and Foreign Medico-Chirurgical Review" has terminated its career, the American Journal stands without a rival.—*London Med. Times and Gazette*, Nov. 24, 1877.

The best medical journal on the continent.—*Boston Med. and Surg. Journal*, April, 1879.

The present number of the American Journal is an exceedingly good one, and gives every promise of maintaining the well-earned reputation of the review. Our venerable contemporary has our best wishes, and we can only express the hope that it may continue its work with as much vigor and excellence for the next fifty years as it has exhibited in the past.—*London Lancet*, Nov. 24, 1877.

The Philadelphia Medical and Physical Journal issued its first number in 1820, and, after a brilliant career, was succeeded in 1827 by the American Journal of the Medical Sciences, a periodical of world-wide reputation; the ablest and one of the oldest periodicals in the world—a journal which has an unsullied record.—*Gross's History of American Med. Literature*, 1876.

The best medical journal ever published in Europe or America.—*Va. Med. Monthly*, May, 1879.

It is universally acknowledged to be the leading American medical journal, and, in our opinion, is second to none in the language.—*Boston Med. and Surg. Journal*, Oct. 1877.

This is the medical journal of our country to which the American physician abroad will point with the greatest satisfaction, as reflecting the state of medical culture in his country. For a great many years it has been the medium through which our ablest writers have made known their discoveries and observations.—*Address of L. P. Yandell, M.D., before International Med. Congress*, Sept. 1876.

And that it was specifically included in the award of a medal of merit to the Publishers in the Vienna Exhibition in 1873.

The subscription price of the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES" has never been raised during its long career. It is still FIVE DOLLARS per annum; and when paid for in advance, the subscriber receives in addition the "MEDICAL NEWS AND ABSTRACT," making in all nearly 2000 large octavo pages per annum, free of postage.

II.

THE MEDICAL NEWS AND ABSTRACT.

Thirty-seven years ago the "MEDICAL NEWS" was commenced as a monthly to convey to the subscribers of the "AMERICAN JOURNAL" the clinical instruction and

* Communications are invited from gentlemen in all parts of the country. Elaborate articles inserted by the Editor are paid for by the publishers.

current information which could not be accommodated in the Quarterly. It consisted of sixteen pages of such matter, together with sixteen more known as the Library Department and devoted to the publishing of books. With the increased progress of science, however, this was found insufficient, and some years since another periodical, known as the "MONTHLY ABSTRACT," was started, and was furnished at a moderate price to subscribers to the "AMERICAN JOURNAL." These two monthlies will hereafter be consolidated, under the title of "THE MEDICAL NEWS AND ABSTRACT," and will be furnished *free of charge* in connection with the "AMERICAN JOURNAL."

The "NEWS AND ABSTRACT" will consist of 64 pages monthly, in a neat cover. It will contain a CLINICAL DEPARTMENT in which will be continued the series of ORIGINAL AMERICAN CLINICAL LECTURES, by gentlemen of the highest reputation throughout the United States, together with a choice selection of foreign Lectures and Hospital Notes and Gleanings. Then will follow the MONTHLY ABSTRACT, systematically arranged and classified, and presenting five or six hundred articles yearly; and each number will conclude with a NEWS DEPARTMENT, giving current professional intelligence, domestic and foreign, the whole fully indexed at the close of each volume, rendering it of permanent value for reference.

As stated above, the subscription price to the "NEWS AND ABSTRACT" will be Two Dollars and a Half per annum, invariably in advance, at which rate it will rank as one of the cheapest medical periodicals in the country. But it will also be furnished, free of all charge, in commutation with the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES," to all who remit FIVE DOLLARS in advance, thus giving to the subscriber, for that very moderate sum, a complete record of medical progress throughout the world, in the compass of about two thousand large octavo pages.

In this effort to furnish so large an amount of practical information at a price so unprecedentedly low, and thus place it within the reach of every member of the profession, the publishers confidently anticipate the friendly aid of all who feel an interest in the dissemination of sound medical literature. They trust, especially, that the subscribers to the "AMERICAN MEDICAL JOURNAL," will call the attention of their acquaintances to the advantages thus offered, and that they will be sustained in the endeavor to permanently establish medical periodical literature on a footing of cheapness never heretofore attempted.

PREMIUM FOR OBTAINING NEW SUBSCRIBERS TO THE "JOURNAL."

Any gentleman who will remit the amount for two subscriptions for 1880, one of which at least must be for a *new subscriber*, will receive as a PREMIUM, free by mail, a copy of any one of the following recent works:—

- "BARNES'S MANUAL OF MIDWIFERY" (see p. 24),
- "TILBURY FOX'S EPITOME OF DISEASES OF THE SKIN," new edition (see p. 18),
- "FOTHERGILL'S ANTAGONISM OF MEDICINES" (see p. 16),
- "HOLDEN'S LANDMARKS, MEDICAL AND SURGICAL" (see p. 6),
- "BROWNE ON THE USE OF THE OPHTHALMOSCOPE" (see p. 29),
- "FLINT'S ESSAYS ON CONSERVATIVE MEDICINE" (see p. 15),
- "STURGES'S CLINICAL MEDICINE" (see p. 14),
- "SWAYNE'S OBSTETRIC APHORISMS," new edition (see p. 21),
- "TANNER'S CLINICAL MANUAL" (see p. 5),
- "WEST ON NERVOUS DISORDERS OF CHILDREN" (see p. 20).

* * Gentlemen desiring to avail themselves of the advantages thus offered will do well to forward their subscriptions at an early day, in order to insure the receipt of complete sets for the year 1880.

☞ The safest mode of remittance is by bank check or postal money order, drawn to the order of the undersigned. Where these are not accessible, remittances for the "JOURNAL" may be made at the risk of the publishers, by forwarding in REGISTERED letters. Address,

HENRY C. LEA'S SON & CO., Nos. 706 and 708 Sansom St., Phila., Pa.

DUNGLISON (ROBLEY), M.D.,*Late Professor of Institutes of Medicine in Jefferson Medical College, Philadelphia.*

MEDICAL LEXICON; A DICTIONARY OF MEDICAL SCIENCE: Containing a concise explanation of the various Subjects and Terms of Anatomy, Physiology, Pathology, Hygiene, Therapeutics, Pharmacology, Pharmacy, Surgery, Obstetrics, Medical Jurisprudence, and Dentistry. Notices of Climate and of Mineral Waters; Formulæ for Official, Empirical, and Dietetic Preparations; with the Accentuation and Etymology of the Terms, and the French and other Synonyms; so as to constitute a French as well as English Medical Lexicon. A New Edition. Thoroughly Revised, and very greatly Modified and Augmented. By RICHARD J. DUNGLISON, M.D. In one very large and handsome royal octavo volume of over 1100 pages. Cloth, \$6 50; leather, raised bands, \$7 50; half Russia, \$8. (*Just Issued.*)

The object of the author from the outset has not been to make the work a mere lexicon or dictionary of terms, but to afford, underneath, a condensed view of its various medical relations, and thus to render the work an epitome of the existing condition of medical science. Starting with this view, the immense demand which has existed for the work has enabled him, in repeated revisions, to augment its completeness and usefulness, until at length it has attained the position of a recognized and standard authority wherever the language is spoken.

Special pains have been taken in the preparation of the present edition to maintain this enviable reputation. During the ten years which have elapsed since the last revision, the additions to the nomenclature of the medical sciences have been greater than perhaps in any similar period of the past, and up to the time of his death the author labored assiduously to incorporate everything requiring the attention of the student or practitioner. Since then, the editor has been equally industrious, so that the additions to the vocabulary are more numerous than in any previous revision. Especial attention has been bestowed on the accentuation, which will be found marked on every word. The typographical arrangement has been much improved, rendering reference much more easy, and every care has been taken with the mechanical execution. The work has been printed on new type, small but exceedingly clear, with an enlarged page, so that the additions have been incorporated with an increase of but little over a hundred pages, and the volume now contains the matter of at least four ordinary octavos.

A book well known to our readers, and of which every American ought to be proud. When the learned author of the work passed away, probably all of us feared lest the book should not maintain its place in the advancing science whose terms it defines. Fortunately, Dr. Richard J. Dunglison, having assisted his father in the revision of several editions of the work, and having been, therefore, trained in the methods and imbued with the spirit of the book, has been able to edit it, not in the patchwork manner so dear to the heart of book editors, so repulsive to the taste of intelligent book readers, but to edit it as a work of the kind should be edited—to carry it on steadily, without jar or interruption, along the grooves of thought it has travelled during its lifetime. To show the magnitude of the task which Dr. Dunglison has assumed and carried through, it is only necessary to state that more than six thousand new subjects have been added in the present edition.—*Phila. Med. Times*, Jan. 3, 1874.

About the first book purchased by the medical student is the Medical Dictionary. The lexicon explanatory of technical terms is simply a *stue qua non*. In a science so extensive, and with such collaterals as medicine, it is as much a necessity also to the practising physician. To meet the wants of students and most physicians, the dictionary must be condensed while comprehensive, and practical while perspicacious. It was because Dunglison's met these indications that it became at once the dictionary of general use wherever medicine was studied in the English language. In no former revision have the alterations and additions been so great. More than six thousand new subjects and terms have been added. The chief terms have been set in black letter, while the derivatives follow in small caps; an arrangement which greatly facilitates reference. We

may safely confirm the hope ventured by the editor "that the work, which possesses for him a filial as well as an individual interest, will be found worthy a continuance of the position so long accorded to it as a standard authority."—*Cincinnati Clinic*, Jan. 10, 1874.

It has the rare merit that it certainly has no rival in the English language for accuracy and extent of references.—*London Medical Gazette*

As a standard work of reference, as one of the best, if not the very best, medical dictionary in the English language, Dunglison's work has been well known for about forty years, and needs no words of praise on our part to recommend it to the members of the medical, and, likewise, of the pharmaceutical profession. The latter especially are in need of such a work, which gives ready and reliable information on thousands of subjects and terms which they are liable to encounter in pursuing their daily avocations, but with which they cannot be expected to be familiar. The work before us fully supplies this want.—*Am. Journ. of Pharm.*, Feb. 1874.

A valuable dictionary of the terms employed in medicine and the allied sciences, and of the relations of the subjects treated under each head. It reflects great credit on its able American author, and well deserves the authority and popularity it has obtained.—*British Med. Journ.*, Oct. 31, 1874.

Few works of this class exhibit a grander monument of patient research and of scientific lore. The extent of the sale of this lexicon is sufficient to testify to its usefulness, and to the great service conferred by Dr. Robley Dunglison on the profession, and indeed on others, by its issue.—*London Lancet*, May 13, 1875.

HOBLYN (RICHARD D.), M.D.

A DICTIONARY OF THE TERMS USED IN MEDICINE AND THE COLLATERAL SCIENCES. Revised, with numerous additions, by ISAAC HAYS, M.D., Editor of the "American Journal of the Medical Sciences." In one large royal 12mo. volume of over 500 double-columned pages; cloth, \$1 50; leather, \$2 00

It is the best book of definitions we have, and ought always to be upon the student's table.—*Southern Med. and Surg. Journal*.

RODWELL (G. F.), F.R.A.S., &c.

A DICTIONARY OF SCIENCE: Comprising Astronomy, Chemistry, Dynamics, Electricity, Heat, Hydrodynamics, Hydrostatics, Light, Magnetism, Mechanics, Meteorology, Pneumatics, Sound, and Statics. Preceded by an Essay on the History of the Physical Sciences. In one handsome octavo volume of 694 pages, and many illustrations: cloth, \$5.

A CENTURY OF AMERICAN MEDICINE, 1776-1876. By Doctors E. H. Clarke, H. J. Bigelow, S. D. Gross, T. G. Thomas, and J. S. Billings. In one very handsome 12mo. volume of about 350 pages: cloth, \$2 25. (*Lately Issued.*)

This work appeared in the pages of the American Journal of the Medical Sciences during the year 1876. As a detailed account of the development of medical science in America, by gentlemen of the highest authority in their respective departments, the profession will no doubt welcome it in a form adapted for preservation and reference.

NEILL (JOHN), M.D., and SMITH (FRANCIS G.), M.D.,

Prof. of the Institutes of Medicine in the Univ. of Penna.

AN ANALYTICAL COMPENDIUM OF THE VARIOUS BRANCHES OF MEDICAL SCIENCE; for the Use and Examination of Students. A new edition, revised and improved. In one very large and handsomely printed royal 12mo. volume, of about one thousand pages, with 374 wood-cuts, cloth, \$4; strongly bound in leather, with raised bands, \$4 75.

HARTSHORNE (HENRY), M.D.,

Professor of Hygiene in the University of Pennsylvania.

A CONSPECTUS OF THE MEDICAL SCIENCES; containing Handbooks on Anatomy, Physiology, Chemistry, Materia Medica, Practical Medicine, Surgery, and Obstetrics. Second Edition, thoroughly revised and improved. In one large royal 12mo. volume of more than 1000 closely printed pages, with 477 illustrations on wood. Cloth, \$4 25; leather, \$5 00. (*Lately Issued.*)

We can say with the strictest truth that it is the best work of the kind with which we are acquainted. It embodies in a condensed form all recent contributions to practical medicine, and is therefore useful to every busy practitioner throughout our country, besides being admirably adapted to the use of students of medicine. The book is faithfully and ably executed.—*Charleston Med. Journ.*, April, 1875.

The work is intended as an aid to the medical student, and as such appears to admirably fulfil its object by its excellent arrangement, the full compilation of facts, the perspicuity and terseness of language, and the clear and instructive illustrations in some parts of the work.—*American Journ. of Pharmacy*, Philadelphia, July, 1874.

The volume will be found useful, not only to students, but to many others who may desire to refresh their memories with the smallest possible expenditure of time.—*N. Y. Med. Journal*, Sept. 1874.

The student will find this the most convenient and useful book of the kind on which he can lay his hand.—*Pacific Med. and Surg. Journ.*, Aug. 1874.

This is the best book of its kind that we have ever examined. It is an honest, accurate, and concise compend of medical sciences, as fairly as possible representing their present condition. The changes and the additions have been so judicious and thorough as to render it, so far as it goes, entirely trust-

worthy. If students must have a conspectus, they will be wise to procure that of Dr. Hartshorne.—*Detroit Rev. of Med. and Pharm.*, Aug. 1874.

The work before us, however, has many redeeming features not possessed by others, and is the best we have seen. Dr. Hartshorne exhibits much skill in condensation. It is well adapted to the physician in active practice, who can give but limited time to the familiarizing of himself with the important changes which have been made since he attended lectures. The manual of physiology has also been improved and gives the most comprehensive view of the latest advances in the science possible in the space devoted to the subject. The mechanical execution of the book leaves nothing to be wished for.—*Peninsular Journal of Medicine*, Sept. 1874.

After carefully looking through this conspectus, we are constrained to say that it is the most complete work, especially in its illustrations, of its kind that we have seen.—*Glasgow Lancet*, Sept. 1874.

The favor with which the first edition of this Compendium was received, was an evidence of its various excellences. The present edition bears evidence of a careful and thorough revision. Dr. Hartshorne possesses a happy faculty of seizing upon the salient points of each subject, and of presenting them in a concise and yet perspicuous manner.—*Leavenworth Med. Herald*, Oct. 1874.

LUDLOW (J. L.), M.D.

A MANUAL OF EXAMINATIONS upon Anatomy, Physiology, Surgery, Practice of Medicine, Obstetrics, Materia Medica, Chemistry, Pharmacy, and Therapeutics. To which is added a Medical Formulary. Third edition, thoroughly revised and greatly extended and enlarged. With 370 illustrations. In one handsome royal 12mo. volume of 816 large pages, cloth, \$3 25; leather, \$3 75.

The arrangement of this volume in the form of question and answer renders it especially suitable for the office examination of students, and for those preparing for graduation.

TANNER (THOMAS HAWKES), M.D., &c.

A MANUAL OF CLINICAL MEDICINE AND PHYSICAL DIAGNOSIS. Third American from the Second London Edition. Revised and Enlarged by TILBURY FOX, M. D., Physician to the Skin Department in University College Hospital, &c. In one neat volume small 12mo., of about 375 pages, cloth, \$1 50.

* * On page 3, it will be seen that this work is offered as a premium for procuring new subscribers to the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES."

GRAY (HENRY), F.R.S.,*Lecturer on Anatomy at St. George's Hospital, London.*

ANATOMY, DESCRIPTIVE AND SURGICAL. The Drawings by H. V. CARTER, M.D., and Dr. WESTMACOTT. The Dissections jointly by the Author and Dr. CARTER. With an Introduction on General Anatomy and Development by T. HOLMES, M.A., Surgeon to St. George's Hospital. A new American, from the eighth enlarged and improved London edition. To which is added "LANDMARKS, MEDICAL AND SURGICAL," by LUTHER HOLDEN, F.R.C.S., author of "Human Osteology," "A Manual of Dissections," etc. In one magnificent imperial octavo volume of 983 pages, with 522 large and elaborate engravings on wood. Cloth, \$6; leather, raised bands, \$7; half Russia, \$7 50. (*Just Ready.*)

The author has endeavored in this work to cover a more extended range of subjects than is customary in the ordinary text-books, by giving not only the details necessary for the student, but also the application of those details in the practice of medicine and surgery, thus rendering it both a guide for the learner, and an admirable work of reference for the active practitioner. The engravings form a special feature in the work, many of them being the size of nature, nearly all original, and having the names of the various parts printed on the body of the cut, in place of figures of reference, with descriptions at the foot. They thus form a complete and splendid series, which will greatly assist the student in obtaining a clear idea of Anatomy, and will also serve to refresh the memory of those who may find in the exigencies of practice the necessity of recalling the details of the dissecting room; while combining, as it does, a complete Atlas of Anatomy, with a thorough treatise on systematic, descriptive, and applied Anatomy, the work will be found of essential use to all physicians who receive students in their offices, relieving both preceptor and pupil of much labor in laying the groundwork of a thorough medical education.

Since the appearance of the last American Edition, the work has received three revisions at the hands of its accomplished editor, Mr. Holmes, who has sedulously introduced whatever has seemed requisite to maintain its reputation as a complete and authoritative standard text-book and work of reference. Still further to increase its usefulness, there has been appended to it the recent work by the distinguished anatomist, Mr. Luther Holden—"Landmarks, Medical and Surgical"—which gives in a clear, condensed, and systematic way, all the information by which the practitioner can determine from the external surface of the body the position of internal parts. Thus complete, the work, it is believed, will furnish all the assistance that can be rendered by type and illustration in anatomical study. No pains have been spared in the typographical execution of the volume, which will be found in all respects superior to former issues. Notwithstanding the increase of size, amounting to over 100 pages and 57 illustrations, it will be kept, as heretofore, at a price rendering it one of the cheapest works ever offered to the American profession.

The recent work of Mr. Holden, which was noticed by us on p. 53 of this volume, has been added as an appendix, so that, altogether, this is the most practical and complete anatomical treatise available to American students and physicians. The former finds in it the necessary guide in making dissections; a very comprehensive chapter on minute anatomy; and about all that can be taught him on general and special anatomy; while the latter, in its treatment of each region from a surgical point of view, and in the valuable edition of Mr. Holden, will find all that will be essential to him in his practice.—*New Remedies*, Aug. 1878.

This work is as near perfection as one could possibly or reasonably expect any book intended as a text-book or a general reference book on anatomy to be. The American publisher deserves the thanks of the profession for appending the recent work of Mr. Holden, "*Landmarks, Medical and Surgical*," which has already been commended as a separate book. The latter work—treating of topographical anatomy—has become an essential to the library of every intelligent practitioner. We know of no book that can take its place, written as it is by a most distinguished anatomist. It would be simply a waste of words to say anything further in praise of Gray's Anatomy, the text-book in almost every medical college in this country, and the daily reference book of every practitioner who has occasion

to consult his books on anatomy. The work is simply indispensable, especially this present American edition.—*Va. Med. Monthly*, Sept. 1878.

The addition of the recent work of Mr. Holden, as an appendix, renders this the most practical and complete treatise available to American students, who find in it a comprehensive chapter on minute anatomy, about all that can be taught on general and special anatomy, while its treatment of each region, from a surgical point of view, in the valuable section by Mr. Holden, is all that will be essential to them in practice.—*Ohio Medical Recorder*, Aug. 1878.

It is difficult to speak in moderate terms of this new edition of "Gray." It seems to be as nearly perfect as it is possible to make a book devoted to any branch of medical science. The labors of the eminent men who have successively revised the eight editions through which it has passed, would seem to leave nothing for future editors to do. The addition of Holden's "*Landmarks*" will make it as indispensable to the practitioner of medicine and surgery as it has been heretofore to the student. As regards completeness, ease of reference, utility, beauty, and cheapness, it has no rival. No student should enter a medical school without it; no physician can afford to have it absent from his library.—*St. Louis Clin. Record*, Sept. 1878.

ALSO FOR SALE SEPARATE—

HOLDEN (LUTHER), F.R.C.S.,*Surgeon to St. Bartholomew's and the Foundling Hospitals.*

LANDMARKS, MEDICAL AND SURGICAL. From the 2d London Ed. In one handsome volume, royal 12mo., of 128 pages: cloth, 88 cents. (*Now Ready.*)

HEATH (CHRISTOPHER), F.R.C.S.,*Teacher of Operative Surgery in University College, London.*

PRACTICAL ANATOMY: A Manual of Dissections. From the Second revised and improved London edition. Edited, with additions, by W. W. KEEN, M.D., Lecturer on Pathological Anatomy in the Jefferson Medical College, Philadelphia. In one handsome royal 12mo. volume of 578 pages, with 247 illustrations. Cloth, \$3 50; leather, \$4 00.

ALLEN (HARRISON), M.D.*Professor of Physiology in the Univ. of Pa.*

A SYSTEM OF HUMAN ANATOMY: INCLUDING ITS MEDICAL and Surgical Relations. For the Use of Practitioners and Students of Medicine. With an Introductory Chapter on Histology. By E. O. SHAKESPEARE, M. D., Ophthalmologist to the Phila. Hosp. In one large and handsome quarto volume, with several hundred original illustrations on lithographic plates, and numerous wood-cuts in the text. (*In Press.*)

In this elaborate work, which has been in active preparation for several years, the author has sought to give, not only the details of descriptive anatomy in a clear and condensed form, but also the practical applications of the science to medicine and surgery. The work thus has claims upon the attention of the general practitioner, as well as of the student, enabling him not only to refresh his recollections of the dissecting room, but also to recognize the significance of all variations from normal conditions. The marked utility of the object thus sought by the author is self-evident, and his long experience and assiduous devotion to its thorough development are a sufficient guarantee of the manner in which his aims have been carried out. No pains have been spared with the illustrations. Those of normal anatomy are from original dissections, drawn on stone by Mr. Hermann Faber, with the name of every part clearly engraved upon the figure, after the manner of "Holden" and "Gray," and in every typographical detail it will be the effort of the publisher to render the volume worthy of the very distinguished position which is anticipated for it.

ELLIS (GEORGE VINER).*Emeritus Professor of Anatomy in University College, London.*

DEMONSTRATIONS OF ANATOMY; Being a Guide to the Knowledge of the Human Body by Dissection. By GEORGE VINER ELLIS, Emeritus Professor of Anatomy in University College, London. From the Eighth and Revised London Edition. In one very handsome octavo volume of over 700 pages, with 256 illustrations. Cloth, \$4.25; leather, \$5.25. (*Just Ready.*)

This work has long been known in England as the leading authority on practical anatomy, and the favorite guide in the dissecting-room, as is attested by the numerous editions through which it has passed. In the last revision, which has just appeared in London, the accomplished author has sought to bring it on a level with the most recent advances of science by making the necessary changes in his account of the microscopic structure of the different organs, as developed by the latest researches in textural anatomy.

Ellis's Demonstrations is the favorite text-book of the English student of anatomy. In passing through eight editions it has been so revised and adapted to the needs of the student that it would seem that it had almost reached perfection in this special line. The descriptions are clear, and the methods of pursuing anatomical investigations are given with such detail that the book is honestly entitled to its name.—*St. Louis Clinical Record*, June, 1879.

The success of this old manual seems to be as well deserved in the present as in the past volumes. The book seems destined to maintain yet for years

its leadership over the English manuals upon dissecting.—*Phila. Med. Times*, May 24, 1879.

As a dissector, or a work to have in hand and studied while one is engaged in dissecting, we regard it as the very best work extant, which is certainly saying a very great deal. As a text-book to be studied in the dissecting-room, it is superior to any of the works upon anatomy.—*Gtinetnatt Med. News*, May 24, 1879.

We most unreservedly recommend it to every practitioner of medicine who can possibly get it.—*Va. Med. Monthly*, June, 1879.

WILSON (ERASMUS), F.R.S.

A SYSTEM OF HUMAN ANATOMY, General and Special. Edited by W. H. GOBZECHT, M.D., Professor of General and Surgical Anatomy in the Medical College of Ohio. Illustrated with three hundred and ninety-seven engravings on wood. In one large and handsome octavo volume, of over 600 large pages; cloth, \$4; leather, \$5.

SMITH (HENRY H.), M.D.,*Prof. of Surgery in the Univ. of Penna., &c.***and HORNER (WILLIAM E.), M.D.,***Late Prof. of Anatomy in the Univ. of Penna.*

AN ANATOMICAL ATLAS; illustrative of the Structure of the Human Body. In one volume, large imperial octavo, cloth, with about six hundred and fifty beautiful figures. \$4 50.

SCHAFER (EDWARD ALBERT), M.D.,*Assistant Professor of Physiology in University College, London.*

A COURSE OF PRACTICAL HISTOLOGY: Being an Introduction to the Use of the Microscope. In one handsome royal 12mo. volume of 304 pages, with numerous illustrations: cloth, \$2 00. (*Just Issued.*)

HORNER'S SPECIAL ANATOMY AND HISTOLOGY. Eighth edition, extensively revised and modified. In 2 vols. 8vo., of over 1000 pages, with 320 wood-cuts: cloth, \$6 00

SHARPEY AND QUAIN'S HUMAN ANATOMY. Revised, by JOSEPH LEIDY, M.D., Prof. of Anat. in Univ. of Penna. In two octavo vols. of about 1300 pages, with 511 illustrations. Cloth, \$4 00.

BELLAMY'S STUDENT'S GUIDE TO SURGICAL ANATOMY: A Text-book for Students preparing

for their Pass Examination. With engravings on wood. In one handsome royal 12mo. volume. Cloth, \$2 25.

CLELAND'S DIRECTORY FOR THE DISSECTION OF THE HUMAN BODY. In one small volume, royal 12mo. of 182 pages: cloth \$1 25.

HARTSHORNE'S HANDBOOK OF ANATOMY AND PHYSIOLOGY. Second edition, revised. In one royal 12mo. vol., with 220 wood-cuts; cloth, \$1 75.

CARPENTER (WILLIAM B.), M.D., F.R.S., F.G.S., F.L.S.,
Registrar to University of London, etc.

PRINCIPLES OF HUMAN PHYSIOLOGY; Edited by HENRY POWER, M.B. Lond., F.R.C.S., Examiner in Natural Sciences, University of Oxford. A new American from the Eighth Revised and Enlarged English Edition, with Notes and Additions, by FRANCIS G. SMITH, M.D., Professor of the Institutes of Medicine in the University of Pennsylvania, etc. In one very large and handsome octavo volume, of 1083 pages, with two plates and 373 engravings on wood; cloth, \$5 50; leather, \$6 50; half Russia, \$7. (Just Issued.)

We have been agreeably surprised to find the volume so complete in regard to the structure and functions of the nervous system in all its relations, a subject that, in many respects, is one of the most difficult of all, in the whole range of physiology, upon which to produce a full and satisfactory treatise of the class to which the one before us belongs. The additions by the American editor give to the work as it is a considerable value beyond that of the last English edition. In conclusion, we can give our cordial recommendation to the work as it now appears. The editors have, with their additions to the only work on physiology in our language that, in the fullest sense of the word, is the production of a philosopher as well as a physiologist, brought it up as fully as could be expected, if not desired, to the standard of our knowledge of its subject at the present day. It will deservedly maintain the place it has always had in the favor of the medical profession.—*Journal of Nervous and Mental Disease*, April, 1877.

Such enormous advances have recently been made in our physiological knowledge, that what was perfectly

new a year or two ago, looks now as if it had been a received and established fact for years. In this encyclopædic way it is unrivalled. Here, as it seems to us, is the great value of the book; one is safe in sending a student to it for information on almost any given subject, perfectly certain of the fullness of information it will convey, and well satisfied of the accuracy with which it will there be found stated.—*London Med. Times and Gazette*, Feb. 17, 1877.

The merits of "Carpenter's Physiology" are so widely known and appreciated that we need only allude briefly to the fact that in the latest edition will be found a comprehensive embodiment of the results of recent physiological investigation. Care has been taken to preserve the practical character of the original work. In fact the entire work has been brought up to date, and bears evidence of the amount of labor that has been bestowed upon it by its distinguished editor, Mr. Henry Power. The American editor has made the latest additions, in order fully to cover the time that has elapsed since the last English edition.—*N. Y. Med. Journal*, Jan. 1877.

FOSTER (MICHAEL), M.D., F.R.S.,
Prof. of Physiology in Cambridge Univ., England.

TEXT-BOOK OF PHYSIOLOGY. A new American, from the third English edition. Edited with notes and additions by EDWARD T. REICHERT, M.D., Demonstrator of Experimental Therapeutics in Univ. of Penna. In one handsome royal 12mo. volume of 1030 pages, with 259 illustrations. Cloth, \$1 50. Leather, \$2 00. (Just Ready.)

AMERICAN EDITOR'S PREFACE.

The high reputation acquired on both sides of the Atlantic by Dr. Foster's "Text-Book on Physiology," as a lucid exposition of functional physiology, in its most modern aspect, has seemed to call for an edition more thoroughly adapted to the wants of the American student. The plan of the author has presupposed an acquaintance with the details of physiological anatomy such as the student is accustomed to look to in his treatises on physiology. The absence of these details has rendered many parts of the work vague, if not altogether incomprehensible, and has therefore proved a serious drawback to the usefulness of the book as an accompaniment to lectures on physiology, as it is usually taught in our schools, and this deficiency the editor has endeavored to supply, by brief notes and the introduction of a large number of illustrations.

The almost limitless amount of material accumulated by modern research has rendered difficult the task of selection and compression, without exceeding the reasonable limits of a convenient text-book. In his selection the editor has been guided by his experience in the wants of students, and has endeavored merely to present, in the most concise form, such facts as would seem to be indispensable to a correct appreciation of the structure and function of the important organs. In accomplishing this, his additions have considerably exceeded his expectations, amounting to about 140 pages, including the illustrations which have been increased in number from 72 to 259. If he shall thus have succeeded in rendering this admirable work better fitted for the wants of the American student, he will feel abundantly rewarded.

Nothing has been omitted from the English edition, and all additions have been distinguished by insertions in brackets [—].

The great popularity of "Foster's Text-book of Physiology," both in England and in this country, renders it unnecessary to say anything further regarding its merit. We shall, therefore, simply point out that, in the American edition, numerous important additions have been made to the text of the third English edition whereby the usefulness of the work as a text-book for students has been greatly increased.—*Phila. Med. and Surg. Reporter*, April 24, 1880.

Foster's Physiology is a most excellent book, not only for the student, but especially for the physician who wishes to revise and increase his physiological knowledge. The style of the author is clear, his arrangement and classification of subjects very satisfactory, and he has presented in this volume a most valuable exposition of physiology as now known.—*The Am. Practitioner*, April, 1880.

Every sentence shows careful and mature thought,

and the student, at the outset, will have to make up his mind that Foster's work requires study, and not simply a cursory reading. The work is well illustrated, and its method clear and logical, and eminently practical, and fully up to the present advanced status of this most important branch of medicine. Foster's work is a credit to his scholarship and research, and will be accepted everywhere by the student and practitioner as a reliable authority. It should be in the hands of every medical student.—*St. Louis Clin. Record*, April, 1880.

In the presentation of the latest and the most serviceable facts the work is peculiarly rich, and the author is also decidedly happy in his freedom from tedium or useless details, and omission of so very little that a book for students needs to teach. The style is scholarly, and the volume a thoroughly welcome one.—*Cincinnati Med. Gaz. and Recorder*, April, 1880.

THE SAME. English Student's Edition, without notes or additions. In one small 12mo. volume of 804 pages with 72 illustrations. Cloth, 75 cents.

DALTON (J. C.), M.D.,

Professor of Physiology in the College of Physicians and Surgeons, New York, &c.

A TREATISE ON HUMAN PHYSIOLOGY. Designed for the use of Students and Practitioners of Medicine. Sixth edition, thoroughly revised and enlarged, with three hundred and sixteen illustrations on wood. In one very beautiful octavo volume, of over 800 pages. Cloth, \$5 50; leather, \$6 50; half Russia, \$7. (Just Issued.)

During the past few years several new works on physiology, and new editions of old works, have appeared, competing for the favor of the medical student, but none will rival this new edition of Dalton. As now enlarged, it will be found also to be, in general, a satisfactory work of reference for the practitioner.—*Chicago Med. Journ. and Examiner*, Jan. 1876.

Prof. Dalton has discussed conflicting theories and conclusions regarding physiological questions with a fairness, a fulness, and a clearness which lend freshness and vigor to the entire book. But his discussions have been so guarded by a refusal of admission to those speculative and theoretical explanations, which at best exist in the minds of observers themselves as only probabilities, that none of his readers need be led into grave errors while making them a study.—*The Medical Record*, Feb 19, 1876.

The revision of this great work has brought it forward with the physiological advances of the day, and renders it, as it has ever been, the finest work for students extant.—*Nashville Journ. of Med. and Surg.*, Jan. 1876.

For clearness and perspicuity, Dalton's Physiology commended itself to the student years ago, and was a pleasant relief from the verbose productions which it supplanted. Physiology has, however, made many advances since then—and while the style has been preserved intact, the work in the present edition has been brought up fully abreast of the times. The new chemical notation and nomenclature have also been introduced into the present edition. Notwithstanding the multiplicity of text-books on physiology, this will lose none

of its old time popularity. The mechanical execution of the work is all that could be desired.—*Peninsular Journal of Medicine*, Dec. 1875.

This popular text-book on physiology comes to us in its sixth edition with the addition of about fifty per cent. of new matter, chiefly in the departments of pathological chemistry and the nervous system, where the principal advances have been realized. With so thorough revision and additions, that keep the work well up to the times, its continued popularity may be confidently predicted, notwithstanding the competition it may encounter. The publisher's work is admirably done.—*St. Louis Med. and Surg. Journ.*, Dec. 1875.

We heartily welcome this, the sixth edition of this admirable text book, than which there are none of equal brevity more valuable. It is cordially recommended by the Professor of Physiology in the University of Louisiana, as by all competent teachers in the United States, and wherever the English language is read, this book has been appreciated. The present edition, with its 316 admirably executed illustrations, has been carefully revised and very much enlarged, although its bulk does not seem perceptibly increased.—*New Orleans Medical and Surgical Journal*, March, 1876.

The present edition is very much superior to every other, not only in that it brings the subject up to the times, but that it does so more fully and satisfactorily than any previous edition. Taken altogether, it remains in our humble opinion, the best text book on physiology in any land or language.—*The Clinic*, Nov. 6, 1875.

GREENE (WILLIAM H.), M.D.,

Demonstrator of Chemistry in Med. Dept., Univ. of Penna.

A MANUAL OF MEDICAL CHEMISTRY. For the Use of Students.

Based upon Bowman's Medical Chemistry. In one royal 12mo. volume of 312 pages. With illustrations. Cloth, \$1 75. (Just Issued.)

It is well written, and gives the latest views on vital chemistry, a subject with which most physicians are not sufficiently familiar. To those who may wish to improve their knowledge in that direction, we can heartily recommend this work as being worthy of a careful perusal.—*Phila. Med. and Surg. Reporter*, April 24, 1880.

The little work before us is one which we think will be studied with pleasure and profit. The descriptions, though brief, are clear, and in most cases sufficient for the purpose. This book will, in nearly all cases, meet general approval.—*Am. Journ. of Pharmacy*, April, 1880.

CLASSEN (ALEXANDER),

Professor in the Royal Polytechnic School, Atz la-Chapelle.

ELEMENTARY QUANTITATIVE ANALYSIS. Translated with

notes and additions by EDGAR F. SMITH, Ph.D., Assistant Prof. of Chemistry in the Towne Scientific School, Univ. of Penna. In one handsome royal 12mo. volume, of 324 pages, with illustrations; cloth, \$2 00. (Just Ready.)

It is probably the best manual of an elementary nature extant, inasmuch as its methods are the best. It teaches by examples, commencing with single determinations, followed by separations, and then

advancing to the analysis of minerals and such products as are met with in applied chemistry. It is an indispensable book for students in chemistry.—*Boston Journ. of Chemistry*, Oct. 1878.

GALLOWAY (ROBERT), F.C.S.,

Prof of Applied Chemistry in the Royal College of Science for Ireland, &c.

A MANUAL OF QUALITATIVE ANALYSIS. From the Fifth London Edition. In one neat royal 12mo. volume, with illustrations; cloth, \$2 75. (Lately Issued.)

REMSEN (IRA), M.D., Ph.D.,

Professor of Chemistry in the Johns Hopkins University, Baltimore.

PRINCIPLES OF THEORETICAL CHEMISTRY, with special reference to the Constitution of Chemical Compounds. In one handsome royal 12mo. vol. of over 232 pages; cloth, \$1 50. (Just Issued.)

BOWMAN'S INTRODUCTION TO PRACTICAL CHEMISTRY, INCLUDING ANALYSIS. Sixth American, from the sixth and revised London edition. With numerous illustrations. In one neat vol., royal 12mo., cloth, \$2 75.

WOHLER AND FITTIG'S OUTLINES OF ORGANIC CHEMISTRY. Translated with additions from the Eighth German Edition. By IRA REMSEN, M.D., Ph.D., Prof. of Chemistry and Physics in Williams

College, Mass. In one volume, royal 12mo. of 550 pp., cloth, \$3.

LEHMANN'S MANUAL OF CHEMICAL PHYSIOLOGY. Translated from the German, with Notes and Additions, by J. CHESTON MORRIS, M.D. With illustrations on wood. In one octavo volume of 336 pages. Cloth, \$2 25.

LEHMANN'S PHYSIOLOGICAL CHEMISTRY. Complete in two large octavo volumes of 1200 pages, with 200 illustrations; cloth, \$6.

FOWNES (GEORGE), Ph.D.

A MANUAL OF ELEMENTARY CHEMISTRY; Theoretical and Practical. Revised and corrected by HENRY WATTS, B.A., F.R.S., author of "A Dictionary of Chemistry," etc. With a colored plate, and one hundred and seventy-seven illustrations. A new American, from the twelfth and enlarged London edition. Edited by ROBERT BRIDGES, M.D. In one large royal 12mo. volume, of over 1000 pages; cloth, \$2 75; leather, \$3 25. (Just Issued.)

Two careful revisions by Mr. Watts, since the appearance of the last American edition of "Fownes," have so enlarged the work that in England it has been divided into two volumes. In reprinting it, by the use of a small and exceedingly clear type, cast for the purpose, it has been found possible to comprise the whole, without omission, in one volume, not unhandy for study and reference. The enlargement of the work has induced the American Editor to confine his additions to the narrowest compass, and he has accordingly inserted only such discoveries as have been announced since the very recent appearance of the work in England, and has added the standards in popular use to the Decimal and Centigrade systems employed in the original.

Among the additions to this edition will be found a very handsome colored plate, representing a number of spectra in the spectroscope. Every care has been taken in the typographical execution to render the volume worthy in every respect of its high reputation and extended use, and though it has been enlarged by more than one hundred and fifty pages, its very moderate price will still maintain it as one of the cheapest volumes accessible to the chemical student.

This work, inorganic and organic, is complete in one convenient volume. In its earliest editions it was fully up to the latest advancements and theories of that time. In its present form, it presents, in a remarkably convenient and satisfactory manner, the principles and leading facts of the chemistry of to-day. Concerning the manner in which the various subjects are treated, much deserves to be said, and mostly, too, in praise of the book. A review of such a work as *Fownes's Chemistry* within the limits of a book-notice for a medical weekly is simply out of the question.—*Cincinnati Lancet and Clinic*, Dec. 14, 1878.

When we state that, in our opinion, the present edition sustains in every respect the high reputation which its predecessors have acquired and enjoyed, we express therewith our full belief in its intrinsic value as a text-book and work of reference.—*Am. Journ. of Pharm.*, Aug. 1878.

The conscientious care which has been bestowed upon it by the American and English editors renders it still, perhaps, the best book for the student and the practitioner who would keep alive the acquisitions of his student days. It has, indeed, reached a some-

what formidable magnitude with its more than a thousand pages, but with less than this no fair representation of chemistry as it now is can be given. The type is small but very clear, and the sections are very lucidly arranged to facilitate study and reference.—*Med. and Surg. Reporter*, Aug. 3, 1878.

The work is too well known to American students to need any extended notice; suffice it to say that the revision by the English editor has been faithfully done, and that Professor Bridges has added some fresh and valuable matter, especially in the inorganic chemistry. The book has always been a favorite in this country, and in its new shape bids fair to retain all its former prestige.—*Boston Jour. of Chemistry*, Aug. 1878.

It will be entirely unnecessary for us to make any remarks relating to the general character of *Fownes's Manual*. For over twenty years it has held the foremost place as a text-book, and the elaborate and thorough revisions which have been made from time to time have little chance for any wide awake rival to step before it.—*Canadian Pharm. Jour.*, Aug. 1878.

As a manual of chemistry it is without a superior in the language.—*Med. Med. Jour.*, Aug. 1878.

ATTFIELD (JOHN), Ph.D.,

Professor of Practical Chemistry to the Pharmaceutical Society of Great Britain, &c.

CHEMISTRY, GENERAL, MEDICAL, AND PHARMACEUTICAL; including the Chemistry of the U. S. Pharmacopœia. A Manual of the General Principles of the Science, and their Application to Medicine and Pharmacy. Eighth edition revised by the author. In one handsome royal 12mo. volume of 700 pages, with illustrations. Cloth, \$2 50; leather, \$3 00. (Just Ready.)

We have repeatedly expressed our favorable opinion of this work, and on the appearance of a new edition of it, little remains for us to say, except that we expect this eighth edition to be as indispensable to us as the seventh and previous editions have been. While the general plan and arrangement have been adhered to, new matter has been added covering the observations made since the former edition. The present differs from the preceding one chiefly in these alterations and in about ten pages of useful tables added in the appendix.—*Am. Jour. of Pharmacy*, May, 1879.

A standard work like Attfield's Chemistry need only be mentioned by its name, without further comments. The present edition contains such alterations and additions as seemed necessary for the demonstration of the latest developments of chemical principles, and the latest applications of chemistry to pharmacy. The author has bestowed arduous labor on the revision, and the extent of the information thus introduced may be estimated from the fact that the index contains three hundred new references relating to additional material.—*Druggists' Circular and Chemical Gazette*, May, 1879.

This very popular and meritorious work has now reached its eighth edition, which fact speaks in the highest terms in commendation of its excellence. It has now become the principal text-book

of chemistry in all the medical colleges in the United States. The present edition contains such alterations and additions as seemed necessary for the demonstration of the latest developments of chemical principles, and the latest applications of chemistry to pharmacy. It is scarcely necessary for us to say that it exhibits chemistry in its present advanced state.—*Cincinnati Medical News*, April, 1879.

The popularity which this work has enjoyed is owing to the original and clear disposition of the facts of the science, the accuracy of the details, and the omission of much which freights many treatises heavily without bringing corresponding instruction to the reader. Dr. Attfield writes for students, and primarily for medical students; he always has an eye to the pharmacopœia and its official preparations; and he is continually putting the matter in the text so that it responds to the questions with which each section is provided. Thus the student learns easily, and can always refresh and test his knowledge.—*Med. and Surg. Reporter*, April 19, '79.

We noticed only about two years and a half ago the publication of the preceding edition, and remarked upon the exceptionally valuable character of the work. The work now includes the whole of the chemistry of the pharmacopœia of the United States, Great Britain, and India.—*New Remedies*, May, 1879.

BLOXAM (C. L.),*Professor of Chemistry in King's College, London.***CHEMISTRY, INORGANIC AND ORGANIC.** From the Second London Edition. In one very handsome octavo volume, of 700 pages, with about 300 illustrations. Cloth, \$4 00; leather, \$5 00. (*Lately Issued.*)

We have in this work a complete and most excellent text-book for the use of schools, and can heartily recommend it as such.—*Boston Med. and Surg. Journ.*, May 28, 1874.

The above is the title of a work which we can most conscientiously recommend to students of chemistry. It is as easy as a work on chemistry could be made, at the same time that it presents a full account of that science as it now stands. We have spoken of the work as admirably adapted to the wants of students; it is quite as well suited to the requirements of practitioners who wish to review their chemistry, or have occasion to refresh their memories on any point relating to it. In a word, it is a book to be read by all who wish to know what is the chemistry of the present day.—*American Practitioner*, Nov. 1873.

It would be difficult for a practical chemist and teacher to find any material fault with this most admirable treatise. The author has given us almost a cyclopaedia within the limits of a convenient volume, and has done so without penning the *useless* paragraphs too commonly making up a great part of the bulk of many cumbersome works. The progressive scientist is not disappointed when he looks for the record of new and valuable processes and discoveries, while the cautious conservative does not find its pages monopolized by uncertain theories and speculations. A peculiar point of excellence is the crystallized form of expression in which great truths are expressed in very short paragraphs. One is surprised at the brief space allotted to an important topic, and yet, after reading it, he feels that little, if any more should have been said. Altogether, it is seldom you see a text-book so nearly faultless.—*Cincinnati Lancet* Nov. 1873.

CLOWES (FRANK), D.Sc., London.*Senior Science-Master at the High School, Newcastle-under-Lyme, etc.***AN ELEMENTARY TREATISE ON PRACTICAL CHEMISTRY AND QUALITATIVE INORGANIC ANALYSIS.** Specially adapted for Use in the Laboratories of Schools and Colleges and by Beginners. From the Second and Revised English Edition, with about fifty illustrations on wood. In one very handsome royal 12mo. volume of 372 pages: cloth, \$2 50. (*Just Issued.*)

It is short, concise, and eminently practical. We therefore heartily commend it to students, and especially to those who are obliged to dispense with a master. Of course, a teacher is in every way desirable, but a good degree of technical skill and practical knowledge can be attained with no other instructor than the very valuable handbook now under consideration.—*St. Louis Clin. Record*, Oct. 1877.

The work is so written and arranged that it can be comprehended by the student without a teacher, and the descriptions and directions for the various work

are so simple, and yet concise, as to be interesting and intelligible. The work is unencumbered with theoretical deductions, dealing wholly with the practical matter, which it is the aim of this comprehensive text-book to impart. The accuracy of the analytical methods are vouched for from the fact that they have all been worked through by the author and the members of his class, from the printed text. We can heartily recommend the work to the student of chemistry as being a reliable and comprehensive one.—*Druggists' Advertiser*, Oct. 15, 1877.

KNAPP'S TECHNOLOGY; or Chemistry Applied to the Arts, and to Manufactures. With American additions by Prof. WALTER R. JOHNSON. In two

very handsome octavo volumes, with 500 wood engravings, cloth, \$6 00.

PARRISH (EDWARD),*Late Professor of Materia Medica in the Philadelphia College of Pharmacy.***A TREATISE ON PHARMACY.** Designed as a Text-Book for the Student, and as a Guide for the Physician and Pharmacist. With many Formulæ and Prescriptions. Fourth Edition, thoroughly revised, by THOMAS S. WIEGAND. In one handsome octavo volume of 977 pages, with 280 illustrations; cloth, \$5 50; leather, \$6 50; half Russia, \$7. (*Lately Issued.*)

Of Dr. Parrish's great work on pharmacy it only remains to be said that the editor has accomplished his work so well as to maintain, in this fourth edition, the high standard of excellence which it had attained in previous editions, under the editorship of its accomplished author. This has not been accomplished without much labor, and many additions and improvements, involving changes in the arrangement of the several parts of the work, and the addition of much new matter. With the modifications thus effected it constitutes, as now presented, a compendium of the science and art indispensable to the pharmacist, and of the utmost value to every practitioner of medicine desirous of familiarizing himself with the pharmaceutical preparation of the articles which he prescribes for his patients.—*Chicago Med. Journ.*, July, 1874.

The work is eminently practical, and has the rare merit of being readable and interesting, while it preserves a strictly scientific character. The whole work reflects the greatest credit on author, editor, and publisher. It will convey some idea of the liberality which has been bestowed upon its production when we mention that there are no less than 280 carefully executed illustrations. In conclusion, we heartily recommend

the work, not only to pharmacists, but also to the multitude of medical practitioners who are obliged to compound their own medicines. It will ever hold an honored place on our own bookshelves.—*Dublin Med. Press and Observer*, Aug. 12, 1874.

We expressed our opinion of a former edition in terms of unqualified praise, and we are in no mood to detract from that opinion in reference to the present edition, the preparation of which has fallen into competent hands. It is a book with which no pharmacist can dispense, and from which no physician can fail to derive much information of value to him in practice.—*Pacific Med. and Surg. Journ.*, June, '74.

Perhaps one, if not the most important book upon pharmacy which has appeared in the English language has emanated from the transatlantic press. "Parrish's Pharmacy" is a well-known work on this side of the water, and the fact shows us that a really useful work never becomes merely local in its fame. Thanks to the judicious editing of Mr. Wiegand, the posthumous edition of "Parrish" has been saved to the public with all the mature experience of its author, and perhaps none the worse for a dash of new blood.—*Lond. Pharm. Journal*, Oct. 17, 1874.

FARQUHARSON (ROBERT), M.D.,

Lecturer on *Materia Medica* at St. Mary's Hospital Medical School.

A GUIDE TO THERAPEUTICS AND MATERIA MEDICA. Second American edition, revised by the Author. Enlarged and adapted to the U. S. Pharmacopœia. By FRANK WOODBURY, M.D. In one neat royal 12mo. volume of 498 pages: cloth, \$2.25. (Just Ready.)

The appearance of a new edition of this convenient and handy book in less than two years may certainly be taken as an indication of its usefulness. Its convenient arrangement, and its terseness, and, at the same time, completeness of the information given, make it a handy book of reference.—*Am. Journ. of Pharmacy*, June, 1879.

This work contains in moderate compass such well-digested facts concerning the physiological and therapeutical action of remedies as are reasonably established up to the present time. By a convenient arrangement the corresponding effects of each article in health and disease are presented in parallel columns, not only rendering reference easier, but also impressing the facts more strongly upon the mind of the reader. The book has been adapted to the wants of the American student, and

copious notes have been introduced, embodying the latest revision of the Pharmacopœia, together with the antidotes to the more prominent poisons, and such of the newer remedial agents as seemed necessary to the completeness of the work. Tables of weights and measures, and a good alphabetical index end the volume.—*Druggists' Circular and Chemical Gazette*, June, 1879.

It is a pleasure to think that the rapidity with which a second edition is demanded may be taken as an indication that the sense of appreciation of the value of reliable information regarding the use of remedies is not entirely overwhelmed in the cultivation of pathological studies, characteristic of the present day. This work certainly merits the success it has so quickly achieved.—*New Remedies*, July, '79.

STILLÉ (ALFRED), M.D.,

Professor of Theory and Practice of Medicine in the University of Penna.

THERAPEUTICS AND MATERIA MEDICA; a Systematic Treatise on the Action and Uses of Medicinal Agents, including their Description and History. Fourth edition, revised and enlarged. In two large and handsome 8vo. vols. of about 2000 pages. Cloth, \$10; leather, \$12; half Russia, \$13. (Lately Issued.)

It is unnecessary to do much more than to announce the appearance of the fourth edition of this well known and excellent work.—*Brit. and For. Med. Chir. Review*, Oct. 1875.

For all who desire a complete work on therapeutics and materia medica for reference, in cases involving medico-legal questions, as well as for information concerning remedial agents, Dr. Stillé's is "*par excellence*" the work. The work being out of print, by the exhaustion of former editions, the author has laid the profession under renewed obligations, by the careful revision, important additions, and timely re-issuing a work not exactly supplemented by any other in the English language, if in any language. The mechanical execution handsomely sustains the well-known skill and good taste of the publisher.—*St. Louis Med. and Surg. Journal*, Dec. 1874.

From the publication of the first edition "Stillé's Therapeutics" has been one of the classics; its absence from our libraries would create a vacuum which could be filled by no other work in the language, and its presence supplies, in the two volumes

of the present edition, a whole cyclopædia of therapeutics.—*Chicago Medical Journal*, Feb. 1876.

The rapid exhaustion of three editions and the universal favor with which the work has been received by the medical profession, are sufficient proof of its excellence as a repository of practical and useful information for the physician. The edition before us fully sustains this verdict, as the work has been carefully revised and in some portions rewritten, bringing it up to the present time by the admission of chloral and croton-chloral nitrite of amyl, bichloride of methylene, methylic ether, lithium compounds, gelsemium, and other remedies.—*Am. Journ. of Pharmacy*, Feb. 1876.

We can hardly admit that it has a rival in the multitude of its citations and the fulness of its research into clinical histories, and we must assign it a place in the physician's library; not, indeed, as fully representing the present state of knowledge in pharmacodynamics, but as by far the most complete treatise upon the clinical and practical side of the question.—*Boston Med. and Surg. Journal*, Nov. 5, 1874.

GRIFFITH (ROBERT E.), M.D.

A UNIVERSAL FORMULARY, Containing the Methods of Preparing and Administering Official and other Medicines. The whole adapted to Physicians and Pharmacutists. Third edition, thoroughly revised, with numerous additions, by JOHN M. MAISON, Professor of Materia Medica in the Philadelphia College of Pharmacy. In one large and handsome octavo volume of about 800 pp., cl., \$4 50; leather, \$5 50. (Lately Issued.)

To the druggist a good formulary is simply indispensable, and perhaps no formulary has been more extensively used than the well-known work before us. Many physicians have to officiate, also, as druggists. This is true especially of the country physician, and a work which shall teach him the means by which to administer or combine his remedies in the most efficacious and pleasant manner, will always hold its place upon his shelf. A formulary of this kind is of benefit also to the city physician in largest practice.—*Cincinnati Clinic*, Feb. 21, 1874.

A more complete formulary than it is in its present form the pharmacist or physician could hardly desire. To the first some such work is indispensable, and it is hardly less essential to the practitioner who compounds his own medicines. Much of what is contained in the introduction ought to be committed to memory by every student of medicine. As a help to physicians it will be found invaluable, and doubtless will make its way into libraries not already supplied with a standard work of the kind.—*The American Practitioner*, Louisville, July, '74.

CHRISTISON'S DISPENSATORY. With copious additions, and 213 large wood engravings. By R. BELLSFIELD GRIFFITH, M.D. One vol. 8vo., pp. 1000, cloth. \$4 00.

CARPENTER'S PRIZE ESSAY ON THE USE OF ALCOHOLIC LIQUORS IN HEALTH AND DISEASE. New edition, with a Preface by D. F. CONDRÉ, M.D., and explanations of scientific words. In one neat 12mo. volume, pp. 178, cloth. 60 cents.

STILLE (ALFRED), M.D., LL.D., and MAISCH (JOHN M.), Ph.D.,
Prof. of Theory and Practice of Medicine and of Clinical Med. in Univ. of Pa. *Prof. of Mat. Med. and Bot. in Phila. Coll. Pharmacy, Secy. to the American Pharmaceutical Association.*

THE NATIONAL DISPENSATORY: Containing the Natural History, Chemistry, Pharmacy, Actions and Uses of Medicines, including those recognized in the Pharmacopœias of the United States, Great Britain, and Germany, with numerous references to the French Codex. Second edition, thoroughly revised, with numerous additions. In one very handsome octavo volume of 1692 pages, with 239 illustrations. Extra cloth, \$6 75; leather, raised bands, \$7 50; half Russia, raised bands and open back, \$8 25. (*Now Ready.*)

PREFACE TO THE SECOND EDITION.

The demand which has exhausted in a few months an unusually large edition of the National Dispensatory is doubly gratifying to the authors, as showing that they were correct in thinking that the want of such a work was felt by the medical and pharmaceutical professions, and that their efforts to supply that want have been acceptable. This appreciation of their labors has stimulated them in the revision to render the volume more worthy of the very marked favor with which it has been received. The first edition of a work of such magnitude must necessarily be more or less imperfect; and though but little that is new and important has been brought to light in the short interval since its publication, yet the length of time during which it was passing through the press rendered the earlier portions more in arrears than the later. The opportunity for a revision has enabled the authors to scrutinize the work as a whole, and to introduce alterations and additions wherever there has seemed to be occasion for improvement or greater completeness. The principal changes to be noted are the introduction of several drugs under separate headings, and of a large number of drugs, chemicals, and pharmaceutical preparations classified as allied drugs and preparations under the heading of more important or better known articles: these additions comprise in part nearly the entire German Pharmacopœia and numerous articles from the French Codex. All new investigations which came to the authors' notice up to the time of publication have received due consideration.

The series of illustrations has undergone a corresponding thorough revision. A number have been added, and still more have been substituted for such as were deemed less satisfactory.

The new matter embraced in the text is equal to nearly one hundred pages of the first edition. Considerable as are these changes as a whole, they have been accommodated by an enlargement of the page without increasing unduly the size of the volume.

While numerous additions have been made to the sections which relate to the physiological action of medicines and their use in the treatment of disease, great care has been taken to make them as concise as was possible without rendering them incomplete or obscure. The doses have been expressed in the terms both of troy weight and of the metrical system, for the purpose of making those who employ the Dispensatory familiar with the latter, and paving the way for its introduction into general use.

The Therapeutical Index has been extended by about 2250 new references, making the total number in the present edition about 6000.

The articles there enumerated as remedies for particular diseases are not only those which, in the authors' opinion, are curative, or even beneficial, but those also which have at any time been employed on the ground of popular belief or professional authority. It is often of as much consequence to be acquainted with the worthlessness of certain medicines or with the narrow limits of their power, as to know the well attested virtues of others and the conditions under which they are displayed. An additional value possessed by such an Index is, that it contains the elements of a natural classification of medicines, founded upon an analysis of the results of experience, which is the only safe guide in the treatment of disease.

This evidence of success, seldom paralleled, shows clearly how well the authors have met the existing needs of the pharmaceutical and medical professions. Gratifying as it must be to them, they have embraced the opportunity offered for a thorough revision of the whole work, striving to embrace within it all that might have been omitted in the former edition, and all that has newly appeared of sufficient importance during the time of its collaboration, and the short interval elapsed since the previous publication. After having gone carefully through the volume we must admit that the authors have labored faithfully, and with success, in maintaining the high character of their work as a compendium meeting the requirements of the day, to which one can safely turn in quest of the latest information concerning everything worthy of notice in connection with Pharmacy, Materia Medica, and Therapeutics.—*Am. Jour. of Pharmacy*, Nov. 1879.

It is with great pleasure that we announce to our readers the appearance of a *second* edition of the National Dispensatory. The total exhaustion of the first edition in the short space of six months, is a sufficient testimony to the value placed upon the work by the profession. It appears that the rapid sale of the first edition must have induced both the editors and the publisher to make preparations for a new edition immediately after the first had been issued, for we find a large amount of new matter added and a good deal of the previous text altered and improved, which proves that the authors do not

intend to let the grass grow under their feet, but to keep the work up to the time.—*New Remedies*, Nov. 1879.

This is a great work by two of the ablest writers on materia medica in America. The authors have produced a work which, for accuracy and comprehensiveness, is unsurpassed by any work on the subject. There is no book in the English language which contains so much valuable information on the various articles of the materia medica. The work has cost the authors years of laborious study, but they have succeeded in producing a dispensatory which is not only national, but will be a lasting memorial of the learning and ability of the authors who produced it.—*Edinburgh Medical Journal*, Nov. 1879.

It is by far more international or universal than any other book of the kind in our language, and more comprehensive in every sense.—*Pacific Med. and Surg. Journ.*, Oct. 1879.

The National Dispensatory is beyond dispute the very best authority. It is throughout complete in all the necessary details, clear and lucid in its explanations, and replete with references to the most recent writings, where further particulars can be obtained, if desired. Its value is greatly enhanced by the extensive indices—a general index of materia medica, etc., and also an index of therapeutics. It would be a work of supererogation to say more about this well-known work. No practicing physician can afford to be without the National Dispensatory.—*Canada Med. and Surg. Journ.*, Feb. 1880.

CORNIL (V.),
Prof. in the Faculty of Med., Paris.

RANVIER (L.),
Prof. in the College of France.

MANUAL OF PATHOLOGICAL HISTOLOGY. Translated, with Notes and Additions, by E. O. SHAKESPEARE, M. D., Pathologist and Ophthalmic Surgeon to Philada. Hospital, Lecturer on Refraction and Operative Ophthalmic Surgery in Univ. of Penna., and by HENRY C. SIMES, M. D., Demonstrator of Pathological Histology in the Univ. of Pa. In one very handsome octavo volume of over 700 pages, with over 350 illustrations. Cloth, \$5 50; leather, \$6 50; half Russia, \$7. (*Just Ready.*)

The work of Cornil and Ranvier is so well known as a lucid and accurate text-book on its important subject, that no apology is needed in presenting a translation of it to the American profession. It is only necessary to say that the labors of Drs. Shakespeare and Simes have been by no means confined to the task of rendering the work into English. As it appeared in France, in successive portions, between 1868 and 1876, a part of it, at least, was somewhat in arrears of the present state of science, while the diffuseness of other portions rendered condensation desirable. The translators have, therefore, sought to bring the work up to the day, and, at the same time, to reduce it in size, as far as practicable, without impairing its completeness. These changes will be found throughout the volume, the most extensive being in the sections devoted to Sarcoma, Carcinoma, Tuberculosis, the Bloodvessels, the Mammas, and the classification of tumors. Corresponding modifications have been made in the very extensive and beautiful series of illustrations, and every care has been taken in the typographical execution to render it one of the most attractive volumes which have issued from the American press.

The translators have done their work very well indeed, rendering it into smooth and excellent English, and in their selection of new material they have in the main used good discrimination. We heartily commend the book as one of the best works on the subject.—*Boston Med. and Surg. Journal*, March 11, 1880.

Their book is not a collection of the work of others, but has been written in the laboratory beside the microscope. It bears the marks of personal knowledge and investigation upon every page, controlled by and controlling the work of others. . . . In short, its translation has made it the best work in pathology attainable in our language, one that every student, certainly, ought to have.—*Archives of Medicine*, April, 1880.

This work, in the original, has for years past occupied a prominent place in the library of French pathologists, as we should naturally be led to believe from the reputation of the distinguished authors. Now that it has been presented to the English student for the first time, it will be perused

with unusual interest. The illustrations are by no means the least valuable part of the work. Indispensable as they are to any work of this nature, in the work before us the artist has succeeded in producing cuts which will prove unusually valuable to the reader. The translation is well done, and gives evidence throughout the volume that it was made by a person thoroughly conversant with the subject.—*N. Y. Med. Gazette*, Feb. 28, 1880.

This will be found an exceedingly interesting and valuable work by all who are engaged in the study of, or take an interest in, histology—normal or morbid. The material which was utilized in its preparation was derived from autopsies and operations in the hospitals of Paris, which are so very rich in it. Nothing was taken for granted, but everything verified by microscopical investigation by the authors themselves in their own laboratories assisted by their pupils. As an aid to microscopists in their investigations it will be found invaluable—in fact, the very best with which we are acquainted.—*Cincinnati Med. News*, February, 1880.

FENWICK (SAMUEL), M.D.,
Assistant Physician to the London Hospital.

THE STUDENT'S GUIDE TO MEDICAL DIAGNOSIS. From the Third Revised and Enlarged English Edition. With eighty-four illustrations on wood. In one very handsome volume, royal 12mo., cloth, \$2 25. (*Just Issued.*)

GREEN (T. HENRY), M.D.,

Lecturer on Pathology and Morbid Anatomy at Charing-Cross Hospital Medical School, &c.

PATHOLOGY AND MORBID ANATOMY. Third American, from the Fourth and Enlarged and Revised English Edition. In one very handsome octavo volume of 332 pages, with 132 illustrations; cloth, \$2 25. (*Now Ready.*)

This is unquestionably one of the best manuals on the subject of pathology and morbid anatomy that can be placed in the student's hands, and we are glad to see it kept up to the times by new editions. Each edition is carefully revised by the author, with the view of making it include the most recent advances in pathology, and of omitting whatever may have become obsolete.—*N. Y. Med. Jour.*, Feb. 1879.

The treatise of Dr. Green is compact, clearly expressed, up to the times, and popular as a text-book, both in England and America. The cuts are suffi-

ciently numerous, and usually well made. In the present edition, such new matter has been added as was necessary to embrace the later results in pathological research. No doubt it will continue to enjoy the favor it has received at the hands of the profession.—*Med and Surg. Reporter*, Feb. 1, 1879.

For practical, ordinary daily use, this is undoubtedly the best treatise that is offered to students of pathology and morbid anatomy.—*Cincinnati Lancet and Clinic*, Feb. 8, 1879.

GLUGE'S ATLAS OF PATHOLOGICAL HISTOLOGY Translated, with Notes and Additions, by JOSEPH LEIDY, M. D. In one volume, very large imperial quarto, with 320 copper-plate figures, plain and colored, cloth. \$4 00.

PAVY'S TREATISE ON THE FUNCTION OF DIGESTION: its Disorders and their Treatment. From the second London edition. In one handsome volume, small octavo, cloth, \$2 00.

LA ROCHE ON YELLOW FEVER, considered in its Historical, Pathological, Etiological, and Therapeutical Relations. In two large and handsome octavo volumes of nearly 1600 pp., cloth. \$7 00.

HOLLAND'S MEDICAL NOTES AND REFLECTIONS. 1 vol. 8vo., pp. 500, cloth. \$3 50.

BARLOW'S MANUAL OF THE PRACTICE OF MEDICINE. With Additions by D. F. COMBIE, M. D. 1 vol. 8vo., pp. 600, cloth. \$2 50.

TODD'S CLINICAL LECTURES ON CERTAIN ACUTE DISEASES. In one neat octavo volume, of 320 pp., cloth. \$2 50.

STOKES'S LECTURES ON FEVER. Edited by JOHN WILLIAM MOORE, M. D., Assistant Physician to the Cork Street Fever Hospital. In one neat 8vo. volume cloth, \$2 00.

FLINT (AUSTIN), M.D.,

Professor of the Principles and Practice of Medicine in Bellevue Med. College, N. Y.

A TREATISE ON THE PRINCIPLES AND PRACTICE OF MEDICINE; designed for the use of Students and Practitioners of Medicine. Fourth edition, revised and enlarged. In one large and closely printed octavo volume of about 1100 pp.; cloth, \$6 00; or strongly bound in leather, with raised bands, \$7 00. (*Lately Issued.*)

This excellent treatise on medicine has acquired for itself in the United States a reputation similar to that enjoyed in England by the admirable lectures of Sir Thomas Watson. We have referred to many of the most important chapters, and find the revision spoken of in the preface is a genuine one, and that the author has very fairly brought up his matter to the level of the knowledge of the present day. The work has this great recommendation, that it is in one volume, and therefore will not be so terrifying to the student as the bulky volumes which several of our English text-books of medicine have developed into. —*British and Foreign Med.-Chir. Rev.*, Jan. 1876.

It is of course unnecessary to introduce or eulogize this now standard treatise. The present edition has been enlarged and revised to bring it up to the author's present level of experience and reading.

His own clinical studies and the latest contributions to medical literature both in this country and in Europe, have received careful attention, so that some portions have been entirely rewritten, and about seventy pages of new matter have been added. —*Chicago Med. Jour.*, June, 1873.

Has never been surpassed as a text-book for students and a book of ready reference for practitioners. The force of its logic, its simple and practical teachings, have left it without a rival in the field. *N. Y.—Med. Record*, Sept. 15, 1874.

It is given to very few men to tread in the steps of Austin Flint, whose single volume on medicine, though here and there defective, is a masterpiece of lucid condensation and of general grasp of an enormously wide subject. —*Lond. Practitioner*, Dec.'73.

BY THE SAME AUTHOR.

CLINICAL MEDICINE; a Systematic Treatise on the Diagnosis and Treatment of Diseases. Designed for Students and Practitioners of Medicine. In one large and handsome octavo volume of 795 pages; cloth, \$4 50; leather, \$5 50; half Russia, \$6. (*Now Ready.*)

It is here that the skill and learning of the great clinician are displayed. He has given us a storehouse of medical knowledge, excellent for the student, convenient for the practitioner, the result of a long life of the most faithful clinical work, collected by an energy as vigilant and systematic as untiring, and weighed by a judgment no less clear than his observation is close. —*Archives of Medicine*, Dec. 1879.

The author of the above work has anticipated a want long felt by those for whom it was especially written—the clinical student during his pupillage, and the busy practitioner. He has given to the medical profession a very necessary and useful work, complete in detail, accurate in observation, brief in statement. —*St. Louis Courier of Med.*, Oct. 1879.

There is every reason to believe that this book will be well received. The active practitioner is frequently in need of some work that will enable him to obtain information in the diagnosis and treatment of cases with comparatively little labor. Dr. Flint has the faculty of expressing himself

clearly, and at the same time so concisely as to enable the searcher to traverse the entire ground of his search, and at the same time obtain all that is essential, without plodding through an interminable space. —*N. Y. Med. Jour.*, Nov. 1879.

The eminent teacher who has written the volume under consideration has recognized the needs of the American profession, and the result is all that we could wish. The style in which it is written is peculiarly the author's; it is clear and forcible, and marked by those characteristics which have rendered him one of the best writers and teachers this country has ever produced. We have not space for so full a consideration of this remarkable work as we would desire. —*S. Louis Clin. Record*, Oct. 1879.

It is venturing little to say that there are few men so well fitted as Dr. Flint to impart information on these last mentioned subjects, and the present work is a timely one as relates both to the author's capacity to undertake it and the need for it as an accompaniment to the multitude now issued, in which the subject of treatment is but little considered. —*New Remedies*, Nov. 1879.

BY THE SAME AUTHOR.

ESSAYS ON CONSERVATIVE MEDICINE AND KINDRED TOPICS. In one very handsome royal 12mo. volume. Cloth, \$1 38. (*Just Issued.*)

WATSON (THOMAS), M.D., &c.

LECTURES ON THE PRINCIPLES AND PRACTICE OF PHYSIC. Delivered at King's College, London. A new American, from the Fifth revised and enlarged English edition. Edited, with additions, and several hundred illustrations, by HENRY HARTSHORNE, M.D., Professor of Hygiene in the University of Pennsylvania. In two large and handsome 8vo. vols. Cloth, \$9 00; leather, \$11 00. (*Lately Published.*)

HARTSHORNE (HENRY), M.D.,

Professor of Hygiene in the University of Pennsylvania

ESSENTIALS OF THE PRINCIPLES AND PRACTICE OF MEDICINE. A handy-book for Students and Practitioners. Fourth edition, revised and improved. With about one hundred illustrations. In one handsome royal 12mo. volume, of about 550 pages, cloth, \$2 63; half bound, \$2 88. (*Lately Issued.*)

DAVIS'S CLINICAL LECTURES ON VARIOUS IMPORTANT DISEASES; being a collection of the Clinical Lectures delivered in the Medical Wards of Mercy Hospital, Chicago. Edited by FRANK H. DAVIS, M.D. Second edition, enlarged. In one handsome royal 12mo. volume. Cloth, \$1 75.

THE CYCLOPEDIA OF PRACTICAL MEDICINE: comprising Treatises on the Nature and Treatment of Diseases, Materia Medica and Therapeutics, Dis-

eases of Women and Children, Medical Jurisprudence, etc. etc. By DUNGLISON, FORBES, TWEEDE, and CONOLLY. In four large super-royal octavo volumes, of 3264 double-columned pages, strongly and handsomely bound in leather, \$15; cloth, \$11.

STURGES'S INTRODUCTION TO THE STUDY OF CLINICAL MEDICINE. Being a Guide to the Investigation of Disease. In one handsome 12mo. volume, cloth, \$1 25. (*Lately Issued.*)

B **BRISTOWE (JOHN SYER), M.D., F.R.C.P.,**
Physician and Joint Lecturer on Medicine, St. Thomas's Hospital.

A TREATISE ON THE PRACTICE OF MEDICINE. Second American edition, revised by the Author. Edited, with Additions, by JAMES H. HUTCHINSON, M.D., Physician to the Penna. Hospital. In one handsome octavo volume of nearly 1200 pages. With illustrations. Cloth, \$5 00; leather, \$6 00; half Russia, \$6 50. (*Now Ready.*)

In reprinting this work from the recent thoroughly revised second English edition, the author has made such corrections as seemed advisable, and has added a chapter on Insanity. The Editor has likewise revised his additions in the light of the latest experience, and the work is presented as reflecting in every way the most modern aspect of medical science, and as fully entitled to maintain the distinguished position accorded to it on both sides of the Atlantic as an authoritative guide for the student, and a complete though concise book of reference for the practitioner. Notwithstanding the author's earnest effort at compression, the additions have amounted to about one-tenth of the previous edition; but by the use of an enlarged page these have been accommodated without increasing the size of the volume, while a reduction in the price renders it one of the cheapest works accessible to the profession.

The second edition of this excellent work, like the first, has received the benefit of Dr. Hutchinson's annotations, by which the phases of disease which are peculiar to this country are indicated, and thus a treatise which was intended for British practitioners and students is made more practically useful on this side of the water. We see no reason to modify the high opinion previously expressed with regard to Dr. Bristowe's work, except by adding our appreciation of the careful labors of the author in following the lateral growth of medical science. The chapter on diseases of the skin and of the nervous system, with a new one on insanity compiled from the best sources outside of the author's own long experience, and the valuable portion relating to general pathology, aid greatly in completing an exceptionally good book for purposes of reference and instruction. — *Boston Medical and Surgical Journal*, February, 1880.

The popularity of the work depends, no doubt, upon the clear and incisive way in which it is written, and the attention to details likely to occur in practice, rather than the discussion of questions of theory. — *New Remedies*, Jan. 1880.

What we said of the first edition, we can, with increased emphasis, repeat concerning this: "Every page is characterized by the utterance of a thoughtful man. What has been said, has been well said, and the book is a fair reflex of all that is certainly known on the subjects considered." — *Ohio Med. Recorder*, Jan. 7, 1880.

This is not only one of the latest and most comprehensive works out on the general subject of Theory and Practice of Medicine, but it is unquestionably one of the best. — *So Med. Practitioner*, Jan. 1880.

R **RICHARDSON (BENJ. W.), M.D., F.R.S., M.A., LL.D., F.S.A.,**
Fellow of the Royal College of Physicians, London.

PREVENTATIVE MEDICINE. In one octavo volume of over 400 pages. (*In Press.*)

W **WOODBURY (FRANK), M.D.,**

Physician to the German Hospital, Philadelphia, late Chief Assist. to Med. Clinic, Jeff. College Hospital, etc.

A HANDBOOK OF THE PRINCIPLES AND PRACTICE OF
Medicine; for the use of Students and Practitioners. Based upon Husband's Handbook of Practice. In one neat volume, royal 12mo. (*Preparing.*)

H **HABERSHON (S. O.), M.D.**

Senior Physician to and late Lecturer on the Principles and Practice of Medicine at Guy's Hospital, etc.

ON THE DISEASES OF THE ABDOMEN, COMPRISING THOSE
of the Stomach, and other parts of the Alimentary Canal, Oesophagus, Cæcum, Intestines, and Peritoneum. Second American, from the third enlarged and revised English edition. With illustrations. In one handsome octavo volume of over 500 pages. Cloth, \$3 50. (*Now Ready.*)

This valuable treatise on diseases of the stomach and abdomen has been out of print for several years, and is therefore not so well known to the profession as it deserves to be. It will be found a cyclopædia of information, systematically arranged, on all diseases of the alimentary tract, from the mouth to the rectum. A fair proportion of each chapter is devoted to symptoms, pathology, and therapeutics. The present edition is fuller than former ones in many particulars, and has been thoroughly revised and

amended by the author. Several new chapters have been added, bringing the work fully up to the times, and making it a volume of interest to the practitioner in every field of medicine and surgery. Perverted nutrition is in some form associated with all diseases we have to combat, and we need all the light that can be obtained on a subject so broad and general. Dr. Habershon's work is one that every practitioner should read and study for himself. — *N. Y. Med. Journ.*, April, 1879.

F **FOTHERGILL (J. MILNER), M.D. Edin., M.R.C.P. Lond.,**
Asst. Phys. to the West Lond. Hosp.; Asst. Phys. to the City of Lond. Hosp., etc.

THE PRACTITIONER'S HANDBOOK OF TREATMENT; Or, the Principles of Therapeutics. Second edition, revised and enlarged. In one very neat octavo volume of about 700 pages. (*In Press.*)

B **BY THE SAME AUTHOR.**

THE ANTAGONISM OF THERAPEUTIC AGENTS, AND WHAT IT TEACHES. Being the Fothergillian Prize Essay for 1878. In one neat volume, royal 12mo. of 156 pages; cloth, \$1 00. (*Just Ready.*)

REYNOLDS (J. RUSSELL), M.D.,

Prof. of the Principles and Practices of Medicine in Univ. College, London.

A SYSTEM OF MEDICINE WITH NOTES AND ADDITIONS BY HENRY HARTSHORNE, M.D., late Professor of Hygiene in the University of Penna. In three large and handsome octavo volumes, containing 3052 closely printed double-columned pages, with numerous illustrations. Sold only by subscription. Price per vol., in cloth, \$5 00; in sheep, \$6.00; half Russia, raised bands, \$6.50. Per set in cloth, \$15; sheep, \$18; half Russia, \$19.50

VOLUME I. (just ready) contains GENERAL DISEASES and DISEASES OF THE NERVOUS SYSTEM.

VOLUME II. (just ready) contains DISEASES OF RESPIRATORY and CIRCULATORY SYSTEMS.

VOLUME III. (just ready) contains DISEASES OF THE DIGESTIVE and BLOOD GLANDULAR SYSTEMS, OF THE URINARY ORGANS, OF THE FEMALE REPRODUCTIVE SYSTEM, and OF THE CUTANEOUS SYSTEM.

Reynolds's SYSTEM OF MEDICINE, recently completed, has acquired, since the first appearance of the first volume, the well-deserved reputation of being the work in which modern British medicine is presented in its fullest and most practical form. This could scarce be otherwise in view of the fact that it is the result of the collaboration of the leading minds of the profession, each subject being treated by some gentleman who is regarded as its highest authority—as for instance, Diseases of the Bladder by Sir HENRY THOMPSON, Malpositions of the Uterus by GRAILY HEWITT, Insanity by HENRY MAUDSLEY, Consumption by J. HUGHES BENNET, Diseases of the Spine by CHARLES BLAND RADCLIFFE, Pericarditis by FRANCIS SIBSON, Alcoholism by FRANCIS E. ANSTIE, Renal Affections by WILLIAM ROBERTS, Asthma by HYDE SALTER, Cerebral Affections by H. CHARLTON BASTIAN, Gout and Rheumatism by ALFRED BARING GARROD, Constitutional Syphilis by JONATHAN HUTCHINSON, Diseases of the Stomach by WILSON FOX, Diseases of the Skin by BALMANNO SQUIRE, Affections of the Larynx by MORELL MACFENZIE, Diseases of the Rectum by BLIZARD CURLING, Diabetes by LAUDER BRUNTON, Intestinal Diseases by JOHN SYER BRISTOWE, Catalepsy and Somnambulism by THOMAS KING CHAMBERS, Apoplexy by J. HUGHLINGS JACKSON, Angina Pectoris by Professor GAIRDNER, Emphysema of the Lungs by Sir WILLIAM JENNER, etc. etc. All the leading schools in Great Britain have contributed their best men in generous rivalry, to build up this monument of medical science. St. Bartholomew's, Guy's, St. Thomas's, University College, St. Mary's in London, while the Edinburgh, Glasgow, and Manchester schools are equally well represented, the Army Medical School at Netley, the military and naval services, and the public health boards. That a work conceived in such a spirit, and carried out under such auspices should prove an indispensable treasury of facts and experience, suited to the daily wants of the practitioner, was inevitable, and the success which it has enjoyed in England, and the reputation which it has acquired on this side of the Atlantic, have sealed it with the approbation of the two pre-eminently practical nations.

Its large size and high price having kept it beyond the reach of many practitioners in this country who desire to possess it, a demand has arisen for an edition at a price which shall render it accessible to all. To meet this demand the present edition has been undertaken. The five volumes and five thousand pages of the original have by the use of a smaller type and double columns, been compressed into three volumes of over three thousand pages, clearly and handsomely printed, and offered at a price which renders it one of the cheapest works ever presented to the American profession.

But not only is the American edition more convenient and lower priced than the English; it is also better and more complete. Some years having elapsed since the appearance of a portion of the work, additions are required to bring up the subjects to the existing condition of science. Some diseases, also, which are comparatively unimportant in England, require more elaborate treatment to adapt the articles devoted to them to the wants of the American physician; and there are points on which the received practice in this country differs from that adopted abroad. The supplying of these deficiencies has been undertaken by HENRY HARTSHORNE, M.D., late Professor of Hygiene in the University of Pennsylvania, who has endeavored to render the work fully up to the day, and as useful to the American physician as it has proved to be to his English brethren. The number of illustrations has also been largely increased, and no effort spared to render the typographical execution unexceptionable in every respect.

Really too much praise can scarcely be given to this noble book. It is a cyclopædia of medicine written by some of the best men of Europe. It is full of useful information such as one finds frequent need of in one's daily work; for no one head can possibly carry all the knowledge one needs in general practice, and one must refer sometimes to one's library. As a book of reference it is invaluable. It is up with the times. It is clear and concentrated in style, and its form is worthy of its famous publisher.—*Louisville Med. News*, Jan. 31, 1880.

"Reynolds' System of Medicine" is justly considered the most popular work on the principles and practice of medicine in the English language. The contributors to this work are gentlemen of well-known reputation on both sides of the Atlantic. Each gentleman has striven to make his part of the work as practical as possible, and the information contained is such as is needed by the busy practitioner.—*St. Louis Med. and Surg. Journ.*, Jan. 20, 1880.

Dr. Hartshorne has made ample additions and revisions, all of which give increased value to the volume, and render it more useful to the American practitioner. There is no volume in English medical literature more valuable, and every purchaser will, on becoming familiar with it, congratulate himself on the possession of this vast store-

house of information, in regard to so many of the subjects with which he should be familiar.—*Gaillard's Med. Journ.*, Feb. 1880.

There is no medical work which we have in times past more frequently and fully consulted when perplexed by doubts as to treatment, or by having unusual or apparently inexplicable symptoms presented to us than "Reynolds' System of Medicine." Among its contributors are gentlemen who are as well known by reputation upon this side of the Atlantic as in Great Britain, and whose right to speak with authority upon the subjects about which they have written, is recognized the world over. They have evidently striven to make their essays as practical as possible, and while these are sufficiently full to entitle them to the name of monographs, they are not loaded down with such an amount of detail as to render them wearisome to the general reader. In a word, they contain just that kind of information which the busy practitioner frequently finds himself in need of. In order that any deficiencies may be supplied, the publishers have committed the preparation of the book for the press to Dr. Henry Hartshorne, whose judicious notes distributed throughout the volume afford abundant evidence of the thoroughness of the revision to which he has subjected it.—*Am. Jour. Med. Sciences*, Jan. 1880.

BARTHOLOW (ROBERTS), A.M., M.D., LL.D.

Prof. of Materia Medica and General Therapeutics in the Juss. Med. Coll. of Phila., &c.

A PRACTICAL TREATISE ON ELECTRICITY IN ITS APPLICATION TO MEDICINE. In one very handsome octavo volume of about 450 pages, with illustrations. (*Preparing.*)

FINLAYSON (JAMES), M.D.,

Physician and Lecturer on Clinical Medicine in the Glasgow Western Infirmary, &c.

CLINICAL DIAGNOSIS; A Handbook for Students and Practitioners of Medicine. In one handsome 12mo. volume, of 546 pages, with 85 illustrations. Cloth, \$2 63. (*Just Issued.*)

The book is an excellent one, clear, concise, convenient, practical. It is replete with the very knowledge the student needs when he quits the lecture-room and the laboratory for the ward and sick-room, and does not lack in information that will meet the wants of experienced and older men.—*Phila. Med. Times*, Jan. 4, 1879.

This is one of the really useful books. It is attractive

before preface to the final page, and ought to be given a place on every office table, because it contains in a condensed form all that is valuable in semeiology and diagnostics to be found in bulkier volumes, and because in its arrangement and complete index, it is unusually convenient for quick reference in any emergency that may come upon the busy practitioner.—*N. C. Med. Journ.*, Jan. 1879.

HAMILTON (ALLAN McLANE), M.D.,

Attending Physician at the Hospital for Epileptics and Paralytics, Blackwell's Island, N.Y., and at the Out-Patients' Department of the New York Hospital.

NERVOUS DISEASES; THEIR DESCRIPTION AND TREATMENT.

In one handsome octavo volume of 512 pages, with 53 illus.; cloth, \$3 50. (*Just Issued.*)

This is unquestionably the best and most complete text-book of nervous diseases that has yet appeared, and were international jealousy in scientific affairs at all possible, we might be excused for a feeling of chagrin that it should be of American parentage. This work, however, has been performed in New York, and has been so well performed that no room is left for anything but commendation. With great skill, Dr. Hamilton has presented to his

readers a succinct and lucid survey of all that is known of the pathology of the nervous system, viewed in the light of the most recent researches. From the preliminary description of the methods of examination and study, and of the instruments of precision employed in the investigation of nervous diseases, up till the final collection of formulæ, the book is eminently practical.—*Brain*, London, Oct. 1878.

CHARCOT (J. M.),

Professor to the Faculty of Med. Paris, Phys. to La Salpêtrière, &c.

LECTURES ON DISEASES OF THE NERVOUS SYSTEM. Translated from the Second Edition by GEORGE SIEGERSON, M.D., M.Ch., Lecturer on Biology, &c., Cath. Univ. of Ireland. With illustrations. 1 vol. 8vo. of 288 pages. Cloth, \$1 75. (*Just Ready.*)

CLINICAL OBSERVATIONS ON FUNCTIONAL NERVOUS DISORDERS By C. HANDFIELD JONES, M.D., Physician to St. Mary's Hospital, &c. See

and American Edition. In one handsome octavo volume of 348 pages, cloth, \$3 25.

MORRIS (MALCOLM), M.D.,

Joint Lecturer on Dermatology, St. Mary's Hospital Med. School.

SKIN DISEASES, Including their Definitions, Symptoms, Diagnosis, Prognosis, Morbid Anatomy, and Treatment. A Manual for Students and Practitioners. In one 12mo. volume of over 300 pages. With illustrations. Cloth, \$1 75. (*Now Ready.*)

To physicians who would like to know something about skin diseases, so that when a patient presents himself for relief they can make a correct diagnosis and prescribe a rational treatment, we unhesitatingly recommend this little book of Dr. Morris. The affections of the skin are described in a terse, lucid manner, and their several characteristics so plainly set forth that diagnosis will be easy. The treatment in each case is such as the experience of the most eminent dermatologists advise.—*Cincinnati Medical News*, April, 1880.

This is emphatically a learner's book; for we can safely say, so far as our judgment goes, that in the whole range of medical literature of a like scope there is no book which for clearness of expression, and methodical arrangement is better adapted to promote a rational conception of dermatology, a branch confessedly difficult and perplexing to the beginner.—*St. Louis Courier of Medicine*, April, 1880.

The author of this manual has evidently a full and intimate acquaintance with the literature of dermatology, and with the most recent developments and

appliances of cutaneous medicine. He has produced a plain, practical book, by aid of which, who chooses may train his eye to the recognition of light but significant differences. The descriptions are neither too vague nor over-refined; the directions for treatment are clear and succinct.—*London Brain*, April, 1880.

The author has handled his subject in a clear and concise manner, and as a text-book to students his manual will be found useful.—*Medical and Surgical Reporter*, March 27, 1880.

The author's task has been well done and has produced one of the best recent works upon the difficult subject of which it treats; there is no work published which gives a better view of the elementary facts and principles of dermatology.—*New Orleans Medical and Surgical Journal*, April, 1880.

This excellent little book is the first work of a distinguished pupil of Jonathan Hutchinson; it recommends itself above all by its clearness, method, and precision.—*Paris Annales de Dermatologie et de Syphiligraphie*, 25 April, 1880.

FOX (TILBURY), M.D., F.R.C.P., and T. C. FOX, B.A., M.R.C.S.,

Physician to the Department for Skin Diseases, University College Hospital.

EPITOME OF SKIN DISEASES. WITH FORMULÆ. FOR STUDENTS AND PRACTITIONERS. Second edition, thoroughly revised and greatly enlarged. In one very handsome 12mo. volume of 216 pages. Cloth, \$1 38. (*Just Issued.*)

FLINT (AUSTIN), M.D.,

Professor of the Principles and Practice of Medicine in Bellevue Hospital Med. College, N. Y.

A MANUAL OF PERCUSSION AND AUSCULTATION; of the Physical Diagnosis of Diseases of the Lunge and Heart, and of Thoracic Aneurism. Second edition. In one handsome royal 12mo. volume: cloth, \$1 63. (*Just Ready.*)

Prof. Flint is so well known as a medical teacher and writer that it seems superfluous to state that the subject has been treated in a thorough and systematic manner. In revising it for a second edition the author has confined himself to such additions as seem likely to render it more useful, not only to students engaged in the practical study of the subject, but also to practitioners as a hand book for ready reference, and we do not hesitate in saying that it would prove a valuable addition to every

physician's library.—*Med. and Surg. Reporter*, March 18, 1880.

The little work before us has already become a standard one, and has become extensively adopted as a text-book. There is certainly none better. It contains the substance of the lessons which the author has for many years given, in connection with practical instruction in auscultation and percussion, to private classes, composed of medical students and practitioners.—*Cincinnati Med. News*, Feb. 1880.

BY THE SAME AUTHOR.

PHTHISIS: ITS MORBID ANATOMY, ETIOLOGY, SYMPTOMATIC EVENTS AND COMPLICATIONS, FATALITY AND PROGNOSIS, TREATMENT, AND PHYSICAL DIAGNOSIS; in a series of Clinical Studies. By AUSTIN FLINT, M.D., Prof. of the Principles and Practice of Medicine in Bellevue Hospital Med. College, New York. In one handsome octavo volume: \$3 50. (*Lately Issued.*)

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON THE DIAGNOSIS, PATHOLOGY, AND TREATMENT OF DISEASES OF THE HEART. Second revised and enlarged edition. In one octavo volume of 550 pages, with a plate, cloth, \$4.

Dr. Flint chose a difficult subject for his researches, and has shown remarkable powers of observation and reflection, as well as great industry, in his treatment of it. His book must be considered the fullest

and clearest practical treatise on those subjects, and should be in the hands of all practitioners and students. It is a credit to American medical literature.—*Amer. Journ. of the Med. Sciences*, July, 1860.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON THE PHYSICAL EXPLORATION OF THE CHEST AND THE DIAGNOSIS OF DISEASES AFFECTING THE RESPIRATORY ORGANS. Second and revised edition. In one handsome octavo volume of 595 pages, cloth, \$4 50.

BROWN (LENNOX), F.R.C.S. Ed.,

Senior Surgeon to the Central London Throat and Ear Hospital, &c.

THE THROAT AND ITS DISEASES. With one hundred Typical Illustrations in colors, and fifty wood engravings, designed and executed by the author. In one very handsome imperial octavo volume of 351 pages; cloth, \$5 00. (*Just Ready.*)

SEILER (CARL), M.D.,

Lecturer on Laryngoscopy at the Univ. of Penna., Chief of the Throat Dispensary at the Univ. Hospital, Phila., &c.

HANDBOOK OF DIAGNOSIS AND TREATMENT OF DISEASES OF THE THROAT AND NASAL CAVITIES. In one handsome royal 12mo. volume, of 156 pages, with 35 illustrations; cloth, \$1. (*Just Ready.*)

We most heartily commend this book as showing sound judgment in practice, and perfect familiarity with the literature of the specialty it so ably epitomizes.—*Phila. Med. Times*, July 5, 1879.

A convenient little handbook, clear, concise, and accurate in its method, and admirably fulfilling its purpose of bringing the subject of which it treats within the comprehension of the general practitioner.—*N. C. Med. Jour.*, June, 1879.

WILLIAMS'S PULMONARY CONSUMPTION; its Nature, Varieties, and Treatment. With an Analysis of One Thousand Cases to exemplify its duration. In one neat octavo volume of about 350 pages; cloth, \$2 50.

SLADE ON DIPHTHERIA; its Nature and Treatment, with an account of the History of its Prevalence in various Countries. Second and revised edition. In one neat royal 12mo. volume, cloth, \$1 25.

WALSHE ON THE DISEASES OF THE HEART AND GREAT VESSELS. Third American Edition. In 1 vol. 8vo., 420 pp., cloth, \$3 00.

CHAMBERS'S MANUAL OF DIET AND REGIMEN IN HEALTH AND SICKNESS. In one handsome octavo volume. Cloth, \$2 75.

LA ROCHE ON PNEUMONIA. 1 vol. 8vo., cloth, of 500 pages. Price, \$3 00.

WILSON'S STUDENT'S BOOK OF CUTANEOUS MEDICINE AND DISEASES OF THE SKIN. In one very handsome royal 12mo. volume. \$3 50.

FULLER ON DISEASES OF THE LUNGS AND AIR-PASSAGES. Their Pathology, Physical Diagnosis, Symptoms, and Treatment. From the second and revised English edition. In one handsome octavo volume of about 500 pages; cloth, \$3 50.

SMITH ON CONSUMPTION; ITS EARLY AND REMEDIABLE STAGES. 1 vol. 8vo., pp. 254. \$2 25.

BASHAM ON RENAL DISEASES: a Clinical Guide to their Diagnosis and Treatment. With Illustrations. In one 12mo. vol. of 304 pages, cloth, \$2 00.

LECTURES ON THE STUDY OF FEVER. By A. HUNSON, M.D., M.R.I.A., Physician to the Meath Hospital. In one vol. 8vo., cloth, \$2 50.

A TREATISE ON FEVER. By ROBERT D. LYONS, K.C.C. In one octavo volume of 362 pages, cloth \$2 25.

HILLIER'S HANDBOOK OF SKIN DISEASES, for Students and Practitioners. Second Am Ed. In one royal 12mo. vol. of 358 pp. With illustrations. Cloth, \$2 25.

BUMSTEAD (FREEMAN J.), M.D., LL.D.,

Professor of Venereal Diseases at the Col. of Phys. and Surg., New York, &c.

THE PATHOLOGY AND TREATMENT OF VENEREAL DISEASES. Including the results of recent investigations upon the subject. Fourth edition, revised and largely rewritten with the co-operation of R. W. Taylor, M.D., of New York, Prof. of Dermatology in the Univ. of Vt. In one large and handsome octavo volume of 835 pages, with 138 illustrations. Cloth, \$4 75; leather, \$5 75; half Russia, \$6 25. (*Just Ready.*)

This work, on its first appearance, immediately took the position of a standard authority on its subject wherever the language is spoken, and the success of an Italian translation shows that it is regarded with equal favor on the Continent of Europe. In repeated editions the author labored sedulously to render it more worthy of its reputation, and in the present revision no pains have been spared to perfect it as far as possible. Several years having elapsed since the publication of the third edition, much material has been accumulated during the interval by the industry of syphilologists, and new views have been enunciated. All this so far as confirmed by observation and experience, has been incorporated; many portions of the volume been rewritten, the series of illustrations has been enlarged and improved, and the whole may be regarded rather as a new work than as a new edition. It is confidently presented as fully on a level with the most advanced condition of syphilology, and as a work to which the practitioner may refer with the certainty of finding clearly and succinctly set forth whatever falls within the scope of such a treatise.

We have to congratulate our countrymen upon the truly valuable addition which they have made to American literature. The careful estimate of the value of the volume, which we have made, justifies us in declaring that this is the best treatise on venereal diseases in the English language, and, we might add, if there is a better in any other tongue we cannot name it; there are certainly no books in which the student or the general practitioner can find such an excellent *résumé* of the literature of any topic, and such practical suggestions regarding the treatment of the various complications of every venereal disease. We take pleasure in repeating that we believe this to be the best treatise on venereal disease in the English language, and we congratulate the authors upon their brilliant addition to American medical literature.—*Chicago Med. Journal and Examiner*, February, 1880.

It is, without exception, the most valuable single work on all branches of the subject of which it treats in any language. The pathology is sound, the work is, at the same time, in the highest degree practical, and the hints that he will get from it for the management of any one case, at all obscure or obstinate,

will more than repay him for the outlay.—*Archives of Medicine*, April, 1880.

This now classical work on venereal disease comes to us in its fourth edition rewritten, enlarged, and materially improved in every way. Dr. Taylor, as we had every reason to expect, has performed this part of his work with unusual excellence. We feel that what has been written has done but scanty justice to the merits of this truly great treatise.—*St. Louis Courier of Medicine*, Feb. 1880.

We find that we have here practically a new book—that the statement of the title-page, as to the fact that it has been largely rewritten, is a sufficiently modest announcement for the important changes in the text. After a thorough examination of the present edition, we can assert confidently that the enormous labor we have described has been here most faithfully and conscientiously performed.—*Amer. Journ. Med. Sci.*, Jan. 1880.

It is one of the best general treatises on venereal diseases with which we are acquainted, and is especially to be recommended as a guide to the treatment of syphils.—*London Practitioner*, March, 1880.

CULLERIER (A.), and

Surgeon to the Hôpital du Midi.

BUMSTEAD (FREEMAN J.),

Professor of Venereal Diseases in the College of Physicians and Surgeons, N. Y.

AN ATLAS OF VENEREAL DISEASES. Translated and Edited by FREEMAN J. BUMSTEAD. In one large imperial 4to. volume of 328 pages, double-columns, with 26 plates, containing about 150 figures, beautifully colored, many of them the size of life; strongly bound in cloth, \$17 00; also, in five parts, stout wrappers, at \$3 per part.

Anticipating a very large sale for this work, it is offered at the very low price of THREE DOLLARS a Part, thus placing it within the reach of all who are interested in this department of practice. Gentlemen desiring early impressions of the plates would do well to order it without delay. A specimen of the plates and text sent free by mail, on receipt of 25 cents.

LEE'S LECTURES ON SYPHILIS AND SOME FORMS OF LOCAL DISEASE AFFECTING PRINCIPALLY THE ORGANS OF GENERATION. In one handsome octavo volume; cloth, \$2 25.

HILL ON SYPHILIS AND LOCAL CONTAGIOUS DISORDERS. In one handsome octavo volume; cloth, \$3 25.

WEST (CHARLES), M.D.,

Physician to the Hospital for Sick Children, London, &c.

LECTURES ON THE DISEASES OF INFANCY AND CHILDHOOD. Fifth American from the sixth revised and enlarged English edition. In one large and handsome octavo volume of 678 pages. Cloth, \$4 50; leather, \$5 50. (*Lately Issued.*)

BY THE SAME AUTHOR. (*Lately Issued.*)

ON SOME DISORDERS OF THE NERVOUS SYSTEM IN CHILDHOOD; being the Lumleian Lectures delivered at the Royal College of Physicians of London, in March, 1871. In one volume small 12mo., cloth, \$1 00.

BY THE SAME AUTHOR.

LECTURES ON THE DISEASES OF WOMEN. Third American, from the Third London edition. In one neat octavo volume of about 550 pages, cloth, \$3 75; leather, \$4 75.

SMITH (J. LEWIS), M.D.,

Clinical Professor of Diseases of Children in the Bellevue Hospital Med. College, N. Y.

A COMPLETE PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Fourth Edition, revised and enlarged. In one handsome octavo volume of about 750 pages, with illustrations. Cloth, \$4 50; leather, \$5 50; half Russia, \$6. (*Now Ready.*)

The very marked favor with which this work has been received wherever the English language is spoken, has stimulated the author, in the preparation of the Fourth Edition, to spare no pains in the endeavor to render it worthy in every respect of a continuance of professional confidence. Many portions of the volume have been rewritten, and much new matter introduced, but by an earnest effort at condensation, the size of the work has not been materially increased.

In the period which has elapsed since the third edition of the work, so extensive have been the advances that whole chapters required to be rewritten, and hardly a page could pass without some material correction or addition. This labor has occupied the writer closely, and he has performed it conscientiously, so that the book may be considered a faithful portrait of an exceptionally wide clinical experience in infantile diseases, erected by a careful study of the recent literature of the subject.—*Med. and Surg. Reporter*, April 5, 1879.

It is scarcely necessary for us to say the work before us is a standard work upon diseases of children, and that no work has a higher standing than it upon those affections. In consequence of its thorough revision, the work has been made of more value than ever, and may be regarded as fully abreast of the times. We cordially commend it to students and physicians. There is no better work in the language on diseases of children.—*Cincinnati Med. News*, March, 1879.

The author has evidently determined that it shall not lose ground in the esteem of the profession for want of the latest knowledge on that important department of medicine. He has accordingly incorporated in the present edition the useful and practical results of the latest study and experience, both American and foreign, especially those bearing on therapeutics. Altogether the book has been greatly improved, while it has not been greatly increased in size.—*New York Medical Journal*, June, 1879.

This excellent work is so well known that an extended notice at this time would be superfluous. The author has taken advantage of the demand for another new edit on to revise in a most careful manner the entire book; and the numerous corrections and additions evince a determination on his part to keep fully abreast with the rapid progress that is being made in the knowledge and treatment of children's diseases. By the adoption of a somewhat closer type, an increase in size of only thirty pages has been necessitated by the new subject matter introduced.—*Boston Med. and Surg. Jour.*, May 29, 1879.

Probably no other work ever published in this country upon a medical subject has reached such a height of popularity as has this well-known treatise. As a text and reference-book it is pre-eminently the authority upon diseases of children. It stands deservedly higher in the estimation of the profession than any other work upon the same subject.—*Nashville Journ. of Med. and Surg.*, May, 1879.

The author of this work has acquired an immense experience as physician to three of the large charities of New York in which children are treated. These asylums afford unsurpassed opportunities for observing the effects of different plans of treatment, and the results as embodied in this volume may be accepted with faith, and should be in the possession of all practitioners *now*. In view of the approaching season when the diseases of children always increase.—*Nat. Med. Review*, April, 1879.

SWAYNE (JOSEPH GRIFFITHS), M.D.,

Physician-Accoucheur to the British General Hospital, &c.

OBSTETRIC APHORISMS FOR THE USE OF STUDENTS COMMENCING MIDWIFERY PRACTICE Second American, from the Fifth and Revised London Edition with Additions by E. R. HUTCHINS, M.D. With illustrations. In one neat 12mo. volume. Cloth, \$1 25. (*Lately Issued.*)

* See p. 3 of this Catalogue for the terms on which this work is offered as a premium to subscribers to the "AMERICAN JOURNAL OF THE MEDICAL SCIENCES."

CHURCHILL ON THE PUERPERAL FEVER AND OTHER DISEASES PECULIAR TO WOMEN. 1 vol. 8vo., pp. 450, cloth. \$2 50.

DEWEES'S TREATISE ON THE DISEASES OF FEMALES. With illustrations. Eleventh Edition, with the Author's last improvements and corrections. In one octavo volume of 536 pages, with plates, cloth. \$3 00.

MEIGS ON THE NATURE, SIGNS, AND TREATMENT OF CHILD BED FEVER. 1 vol. 8vo., pp. 365, cloth. \$2 00.

ASHWELL'S PRACTICAL TREATISE ON THE DISEASES PECULIAR TO WOMEN. Third American, from the Third and revised London edition. 1 vol. 8vo., pp. 528, cloth. \$3 50.

HODGE (HUGH L.), M.D.,

Emeritus Professor of Obstetrics, &c., in the University of Pennsylvania.

ON DISEASES PECULIAR TO WOMEN; including Displacements of the Uterus. With original illustrations. Second edition, revised and enlarged. In one beautifully printed octavo volume of 531 pages, cloth, \$4 50.

CHURCHILL (FLEETWOOD), M.D., M.R.I.A.

ON THE THEORY AND PRACTICE OF MIDWIFERY. A new American from the fourth revised and enlarged London edition. With notes and additions by D. FRANCIS CONDIM, M.D., author of a "Practical Treatise on the Diseases of Children," &c. With one hundred and ninety four illustrations. In one very handsome octavo volume of nearly 700 large pages. Cloth, \$4 00; leather, \$5 00.

MONTGOMERY'S EXPOSITION OF THE SIGNS AND SYMPTOMS OF PREGNANCY. With two exquisitely colored plates, and numerous wood-cuts. In 1 vol. 8vo., of nearly 600 pp., cloth, \$3 75.

CONDIE'S PRACTICAL TREATISE ON THE DISEASES OF CHILDREN. Sixth edition, revised and augmented. In one large octavo volume of nearly 800 closely-printed pages, cloth, \$5 25; leather \$6 25.

RIGBY'S SYSTEM OF MIDWIFERY. With notes and Additional illustrations. Second American edition. One volume octavo, cloth 422 pages, \$2 50.

SMITH'S PRACTICAL TREATISE ON THE WAITING DISEASES OF INFANCY AND CHILDHOOD. Second American, from the second revised and enlarged English edition. In one handsome octavo volume, cloth, \$2 50.

THOMAS (T. GAILLARD), M.D.,*Professor of Obstetrics, &c., in the College of Physicians and Surgeons, N. Y., &c***A PRACTICAL TREATISE ON THE DISEASES OF WOMEN. Fourth edition, enlarged and thoroughly revised. In one large and handsome octavo volume of 800 pages, with 191 illustrations. Cloth, \$5 00; leather, \$6 00. (Just Issued.)**

A work which has reached a fourth edition, and that, too, in the short space of five years, has achieved a reputation which places it almost beyond the reach of criticism, and the favorable opinions which we have already expressed of the former editions seem to require that we should do little more than announce this new issue. We cannot refrain from saying that, as a practical work, this is second to none in the English, or, indeed, in any other language. The arrangement of the contents, the admirably clear manner in which the subject of the differential diagnosis of several of the diseases is handled, leave nothing to be desired by the practitioner who wants a thoroughly clinical work, one to which he can refer in difficult cases of doubtful diagnosis with the certainty of gaining light and instruction. Dr. Thomas is a man with a very clear head and decided views, and there seems to be nothing which he so much dislikes as hazy notions of diagnosis and blind routine and unreasonable therapeutics. The student who will thoroughly study this book and test its principles by clinical observation, will certainly not be guilty of these faults.—*London Lancet*, Feb. 13, 1875.

Reluctantly we are obliged to close this unsatisfactory notice of so excellent a work, and in conclusion would remark that, as a teacher of gynecology, both didactic and clinical, Prof. Thomas has certainly taken the lead far ahead of his *confères*, and as an author he certainly has met with unusual and merited success.—*Am. Journ. of Obstetrics*, Nov. 1874.

This volume of Prof. Thomas in its revised form

is classical without being pedantic, full in the details of anatomy and pathology, without ponderous translation of pages of German literature, describes distinctly the details and difficulties of each operation, without wearying and useless minutiae, and is in all respects a work worthy of confidence, justifying the high regard in which its distinguished author is held by the profession.—*Am. Supplement, Obstet. Journ.*, Oct. 1874.

Professor Thomas fairly took the Profession of the United States by storm when his book first made its appearance early in 1868. Its reception was simply enthusiastic, notwithstanding a few adverse criticisms from our transatlantic brethren, the first large edition was rapidly exhausted, and in six months a second one was issued, and in two years a third one was announced and published, and we are now promised the fourth. The popularity of this work was not ephemeral, and its success was unprecedented in the annals of American medical literature. Six years is a long period in medical scientific research, but Thomas's work on "Diseases of Women" is still the leading native production of the United States. The order, the matter, the absence of theoretical disputativeness, the fairness of statement, and the elegance of diction, preserved throughout the entire range of the book, indicate that Professor Thomas did not overestimate his powers when he conceived the idea and executed the work of producing a new treatise upon diseases of women.—*PROF. PALLER, in Louisville Med. Journal*, Sept. 1874.

BARNES (ROBERT), M.D., F.R.C.P.,*Obstetric Physician to St. Thomas's Hospital, &c.***A CLINICAL EXPOSITION OF THE MEDICAL AND SURGICAL DISEASES OF WOMEN. Second American, from the Second Enlarged and Revised English Edition. In one handsome octavo volume, of 784 pages, with 181 illustrations. Cloth, \$4 50; leather, \$5 50; half Russia, \$6. (Just Ready.)**

The call for a new edition of Dr. Barnes's work on the Diseases of Females has encouraged the author to make it even more worthy of the favor of the profession than before. By a rearrangement and careful pruning space has been found for a new chapter on the Gynecological Relations of the Bladder and Bowel Disorders, without increasing the size of the book, while many new illustrations have been introduced where experience has shown them to be needed. It is therefore hoped that the volume will be found to reflect thoroughly and accurately the present condition of gynecological science.

Dr. Barnes stands at the head of his profession in the old country, and it requires but scant scrutiny of his book to show that it has been sketched by a master. It is plain, practical common sense; shows very deep research without being pedantic; is eminently calculated to inspire enthusiasm without inculcating rashness; points out the dangers to be avoided as well as the success to be achieved in the various operations connected with this branch of medicine; and will do much to smooth the rugged path of the young gynecologist and relieve the perplexity of the man of mature years.—*Canadian Journ. of Med. Science*, Nov. 1878.

We pity the doctor who, having any considerable practice in diseases of women, has no copy of "Barnes" for daily consultation and instruction. It is at once a book of great learning, research, and individual experience, and at the same time eminently practical. That it has been appreciated by the profession, both in Great Britain and in this country, is shown by the second edition following so soon upon the first.—*Am. Practitioner*, Nov. 1878.

Dr. Barnes's work is one of a practical character, largely illustrated from cases in his own experience, but by no means confined to such, as will be learned from the fact that he quotes from no less than 628 medical authors in numerous countries. Coming from such an author, it is not necessary to say that

the work is a valuable one, and should be largely consulted by the profession.—*Am. Supp. Obstetrical Journ. Gt. Britain and Ireland*, Oct. 1878.

No other gynecological work holds a higher position, having become an authority everywhere in diseases of women. The work has been brought fully abreast of present knowledge. Every practitioner of medicine should have it upon the shelves of his library, and the student will find it a superior text-book.—*Cincinnati Med. News*, Oct. 1878.

This second revised edition, of course, deserves all the commendation given to its predecessor, with the additional one that it appears to include all or nearly all the additions to our knowledge of its subject that have been made since the appearance of the first edition. The American references are, for an English work, especially full and appreciative, and we can cordially recommend the volume to American readers.—*Journ. of Nervous and Mental Disease*, Oct. 1878.

This second edition of Dr. Barnes's great work comes to us containing many additions and improvements which bring it up to date in every feature. The excellences of the work are too well known to require enumeration, and we hazard the prophecy that they will for many years maintain its high position as a standard text-book and guide-book for students and practitioners.—*N. C. Med. Journ.*, Oct. 1878.

CHADWICK (JAMES R.), A.M., M.D.**A MANUAL OF THE DISEASES PECULIAR TO WOMEN. In one neat volume, royal 12mo., with illustrations. (Preparing.)**

EMMET (THOMAS ADDIS), M.D.
Surgeon to the Woman's Hospital, New York, etc.

THE PRINCIPLES AND PRACTICE OF GYNÆCOLOGY, for the use of Students and Practitioners of Medicine. Second Edition. Thoroughly Revised. In one large and very handsome octavo volume of 875 pages, with 133 illustrations. Cloth, \$5; leather, \$6; half Russia, \$6 50. (*Just Ready*.)

PREFACE TO THE SECOND EDITION.

The unusually rapid exhaustion of a large edition of this work, while flattering to the author as an evidence that his labors have proved acceptable, has in a great measure heightened his sense of responsibility. He has therefore endeavored to take full advantage of the opportunity afforded to him for its revision. Every page has received his earnest scrutiny; the criticisms of his reviewers have been carefully weighed; and while no marked increase has been made in the size of the volume, several portions have been rewritten, and much new matter has been added. In this minute and thorough revision, the labor involved has been much greater than is perhaps apparent in the results, but it has been cheerfully expended in the hope of rendering the work more worthy of the favor which has been accorded to it by the profession.

In no country of the world has gynecology received more attention than in America. It is, then, with a feeling of pleasure that we welcome a work on diseases of women from so eminent a gynecologist as Dr. Emmet, and the work is essentially clinical, and leaves a strong impress of the author's individuality. To criticize, with the care it merits, the book throughout, would demand far more space than is at our command. In parting, we can say that the work teems with original ideas, fresh and valuable methods of practice, and is written in a clear and elegant style, worthy of the literary reputation of the country of Longfellow and Oliver Wendell Holmes.—*Brit. Med. Journ.*, Feb. 21, 1880.

No gynecological treatise has appeared which contains an equal amount of original and useful matter; nor does the medical and surgical history of America include a book more novel and useful. The tabular and statistical information which it contains is marvellous, both in quantity and accuracy, and cannot be otherwise than invaluable to future investigators. It is a work which demands

not careless reading but profound study. Its value as a contribution to gynecology is, perhaps, greater than that of all previous literature on the subject combined.—*Chicago Med. Gas.*, April 6, 1880.

The wide reputation of the author makes its publication an event in the gynecological world; and a glance through its pages shows that it is a work to be studied with care. . . . It must always be a work to be carefully studied and frequently consulted by those who practise this branch of our profession.—*London Med. Times and Gas.*, Jan. 10, 1880.

The character of the work is too well known to require extended notice—suffice it to say that no recent work upon any subject has attained such great popularity so rapidly. As a work of general reference upon the subject of Diseases of Women it is invaluable. As a record of the largest clinical experience and observation it has no equal. No physician who pretends to keep up with the advances of this department of medicine can afford to be without it.—*Nashville Journ. of Medicine and Surgery*, May, 1880.

DUNCAN (J. MATTHEWS), M.D., LL.D., F.R.S.E., etc.

CLINICAL LECTURES ON THE DISEASES OF WOMEN, Delivered in Saint Bartholomew's Hospital. In one very neat octavo volume of 173 pages. Cloth, \$1 50. (*Just Ready*.)

They are in every way worthy of their author; indeed, we look upon them as among the most valuable of his contributions. They are all upon matters of great interest to the general practitioner. Some of them deal with subjects that are not, as a rule, adequately handled in the text-books; others of them, while bearing upon topics that are usually treated of at length in such works, yet bear such a stamp of individuality that, if widely read, as they certainly deserve to be, they cannot fail to exert a wholesome restraint upon the undue eagerness with which many young physicians seem bent upon following the wild teachings which so infect the gynecology of the present day.—*N. Y. Med. Journ.*, March, 1880.

The author is a remarkably clear lecturer, and his discussion of symptoms and treatment is full and suggestive. It will be a work which will not fail to be read with benefit by practitioners as well as by students.—*Phila. Med. and Surg. Reporter*, Feb. 7, 1880.

We have read this book with a great deal of pleasure. It is full of good things. The hints on pathology and treatment scattered through the book are sound, trustworthy, and of great value. A healthy scepticism, a large experience, and a clear judgment are everywhere manifest. Instead of bristling with advice of doubtful value and unsound character, the book is in every respect a safe guide.—*The London Lancet*, Jan. 21, 1880.

RAMSBOTHAM (FRANCIS H.), M.D.

THE PRINCIPLES AND PRACTICE OF OBSTETRIC MEDICINE AND SURGERY, in reference to the Process of Parturition. A new and enlarged edition, thoroughly revised by the author. With additions by W. V. KEATINGE, M.D., Professor of Obstetrics, &c., in the Jefferson Medical College, Philadelphia. In one large and handsome imperial octavo volume of 650 pages, strongly bound in leather, with raised bands; with sixty-four beautiful plates, and numerous wood-cuts in the text, containing in all nearly 200 large and beautiful figures. \$7 00.

WINCKEL (F.),

Professor and Director of the Gynecological Clinic in the University of Rostock.

A COMPLETE TREATISE ON THE PATHOLOGY AND TREATMENT OF CHILD-BED, for Students and Practitioners. Translated, with the consent of the author, from the Second German Edition, by JAMES READ CHADWICK, M.D. In one octavo volume. Cloth, \$4 00. (*Lately Issued*.)

TANNER (THOMAS H.), M.D.

ON THE SIGNS AND DISEASES OF PREGNANCY. First American from the Second and Enlarged English Edition. With four colored plates and illustrations on wood. In one handsome octavo volume of about 500 pages, cloth, \$4 25.

LEISHMAN (WILLIAM), M.D.,

Regius Professor of Midwifery in the University of Glasgow, &c.

A SYSTEM OF MIDWIFERY, INCLUDING THE DISEASES OF PREGNANCY AND THE PUERPERAL STATE. Third American edition, revised by the Author, with additions by JOHN S. PARRY, M.D., Obstetrician to the Philadelphia Hospital, &c. In one large and very handsome octavo volume, of 733 pages, with over two hundred illustrations. Cloth, \$4 50; leather, \$5 50; half Russia, \$6. (*Just Ready.*)

Few works on this subject have met with as great a demand as this one appears to have. To judge by the frequency with which its author's views are quoted, and its statements referred to in obstetrical literature, one would judge that there are few physicians devoting much attention to obstetrics who are without it. The author is evidently a man of ripe experience and conservative views, and in no branch of medicine are these more valuable than in this.—*New Remedies*, Jan. 1880.

We gladly welcome the new edition of this excellent text-book of midwifery. The former editions have been most favorably received by the profession on both sides of the Atlantic. In the preparation of the present edition the author has made such alterations as the progress of obstetrical science seems to require, and we cannot but admire the ability with which the task has been performed. We consider it an admirable text-book for students during their attendance upon lectures, and have great pleasure in recommending it. As an exponent of the midwifery of the present day it has no superior in the English language.—*Canada Lancet*, Jan. 1880.

The book is greatly improved, and as such will be welcomed by those who are trying to keep posted in the rapid advances which are being made in the study of obstetrics.—*Boston Med. and Surg. Journ.*, Nov. 27, 1879.

To the American student the work before us must prove admirably adapted, complete in all its parts, essentially modern in its teachings and with demonstrations noted for clearness and precision, it will gain in favor and be recognized as a work of standard merit. The work cannot fail to be popular, and is cordially recommended.—*N. O. Med. and Surg. Journ.*, March, 1880.

Leishman's is certainly one of the best systematic works on midwifery. It is very complete in all the parts essential for such a treatise. To practitioners and students it is to be strongly recommended as a safe and reliable guide to the modern practice of midwifery.—*Canada Med. and Surg. Journal*, March, 1880.

PARRY (JOHN S.), M.D.,

Obstetrician to the Philadelphia Hospital, Vice-Prest. of the Obstet. Society of Philadelphia.

EXTRA-UTERINE PREGNANCY: ITS CLINICAL HISTORY, DIAGNOSIS, PROGNOSIS, AND TREATMENT. In one handsome octavo volume. Cloth, \$2 50. (*Lately Issued.*)

HODGE (HUGH L.), M.D.,

Emeritus Professor of Midwifery, &c., in the University of Pennsylvania, &c.

THE PRINCIPLES AND PRACTICE OF OBSTETRICS. Illustrated with large lithographic plates containing one hundred and fifty-nine figures from original photographs, and with numerous wood-cuts. In one large and beautifully printed quarto volume of 550 double-columned pages, strongly bound in cloth, \$14.

The work of Dr. Hodge is something more than a simple presentation of his particular views in the department of Obstetrics; it is something more than an ordinary treatise on midwifery; it is, in fact, a cyclopædia of midwifery. He has aimed to embody in a single volume the whole science and art of Obstetrics. An elaborate text is combined with accurate and varied pictorial illustrations, so that no fact or principle is left unstated or unexplained.—*Am. Med. Times*, Sept. 3, 1864.

body in a single volume the whole science and art of Obstetrics. An elaborate text is combined with accurate and varied pictorial illustrations, so that no fact or principle is left unstated or unexplained.—*Am. Med. Times*, Sept. 3, 1864.

*** Specimens of the plates and letter-press will be forwarded to any address, free by mail, on receipt of six cents in postage stamps.

STIMSON (LEWIS A.), A.M., M.D.,

Surgeon to the Presbyterian Hospital.

A MANUAL OF OPERATIVE SURGERY. In one very handsome royal 12mo. volume of about 500 pages, with 332 illustrations; cloth, \$2 50. (*Just Issued.*)

The work before us is a well printed, profusely illustrated manual of over four hundred and seventy pages. The novice, by a perusal of the work, will gain a good idea of the general domain of operative surgery, while the practical surgeon has presented to him within a very concise and intelligible form the latest and most approved selections of operative procedure. The precision and conciseness with which the different operations are described enable the author to compress an immense amount of practical information in a very small compass.—*N. Y. Medical Record*, Aug. 3, 1878.

This volume is devoted entirely to operative surgery, and is intended to familiarize the student with the details of operations and the different modes of

performing them. The work is handsomely illustrated, and the descriptions are clear and well drawn. It is a clever and useful volume; every student should possess one. The preparation of this work does away with the necessity of pondering over larger works on surgery for descriptions of operations, as it presents in a nut-shell just what is wanted by the surgeon without an elaborate search to find it.—*Med. Med. Journal*, Aug. 1878.

The author's conciseness and the repleteness of the work with valuable illustrations entitle it to be classed with the text-books for students of operative surgery, and as one of reference to the practitioner.—*Cincinnati Lancet and Clinic*, July 27, 1878.

SKRY'S OPERATIVE SURGERY. In 1 vol. 8vo. cl., of 660 pages; with about 100 wood-cuts. \$3 25.

COOPER'S LECTURES ON THE PRINCIPLES AND PRACTICE OF SURGERY. In 1 vol. 8vo. cl'h, 760 p. \$2.

GIBSON'S INSTITUTES AND PRACTICE OF SURGERY. Eighth edit'n, improved and altered. With thirty-four plates. In two handsome octavo volumes, about 1000 pp., leather, raised band. \$6 60.

THE PRINCIPLES AND PRACTICE OF SURGERY. By WILLIAM PIRRIE, F.R.S.E., Professor of Surgery in the University of Aberdeen. Edited by JOHN

NEILL, M.D., Professor of Surgery in the Penna. Medical College, Surg'n to the Pennsylvania Hospital, &c. In one very handsome octavo vol. of 780 pages, with 316 illustrations, cloth, \$3 75.

MILLER'S PRINCIPLES OF SURGERY. Fourth American, from the Third Edinburgh Edition. In one large 8vo. vol. of 700 pages, with 340 illustrations, cloth, \$3 75.

MILLER'S PRACTICE OF SURGERY. Fourth American, from the last Edinburgh Edition. Revised by the American editor. In one large 8vo. vol. of nearly 700 pages, with 364 illustrations: cloth, \$3 75.

GROSS (SAMUEL D.), M.D.,

Professor of Surgery in the Jefferson Medical College of Philadelphia.

SYSTEM OF SURGERY: Pathological, Diagnostic, Therapeutic, and Operative. Illustrated by upwards of Fourteen Hundred Engravings. Fifth edition, carefully revised, and improved. In two large and beautifully printed imperial octavo volumes of about 2300 pp., strongly bound in leather, with raised bands, \$15; half Russia, raised bands, \$16.

The continued favor, shown by the exhaustion of successive large editions of this great work, proves that it has successfully supplied a want felt by American practitioners and students. In the present revision no pains have been spared by the author to bring it in every respect fully up to the day. To effect this a large part of the work has been rewritten, and the whole enlarged by nearly one-fourth, notwithstanding which the price has been kept at its former very moderate rate. By the use of a close, though very legible type, an unusually large amount of matter is condensed in its pages, the two volumes containing as much as four or five ordinary octavos. This, combined with the most careful mechanical execution, and its very durable binding renders it one of the cheapest works accessible to the profession. Every subject properly belonging to the domain of surgery is treated in detail, so that the student who possesses this work may be said to have in it a surgical library.

We have now brought our task to a conclusion, and have seldom read a work with the practical value of which we have been more impressed. Every chapter is so concisely put together, that the busy practitioner, when in difficulty, can at once find the information he requires. His work, on the contrary, is cosmopolitan, the surgery of the world being fully represented in it. The work, in fact, is so historically unprejudiced, and so eminently practical, that it is almost a false compliment to say that we believe it to be destined to occupy a foremost place as a work of reference, while a system of surgery like the present system of surgery is the practice of surgeons. The printing and binding of the work is unexceptionable; indeed, it contrasts, in the latter respect, remarkably with English medical and surgical cloth-bound publications, which are generally so wretchedly stitched as to require re-binding before they are any time in use.—*Dub. Journ. of Med. Sci.*, March, 1874.

Dr. Gross's Surgery, a great work, has become still greater, both in size and merit, in its most recent form. The difference in actual number of pages is not more than 130, but the size of the page having been increased to what we believe is technically termed "elephant," there has been room for considerable additions, which, together with the alterations, are improvements.—*London Lancet*, Nov. 16, 1872.

It combines, as perfectly as possible, the qualities of a text-book and work of reference. We think this last

edition of Gross's "Surgery," will confirm his title of "*Primus inter Pares*." It is learned, scholar-like, methodical, precise, and exhaustive. We scarcely think any living man could write so complete and faultless a treatise, or comprehend more solid, instructive matter in the given number of pages. The labor must have been immense, and the work gives evidence of great powers of mind, and the highest order of intellectual discipline and methodical disposition, and arrangement of acquired knowledge and personal experience.—*N.Y. Med. Journ.*, Feb. 1873.

As a whole, we regard the work as the representative "System of Surgery" in the English language.—*St. Louis Medical and Surg. Journ.*, Oct. 1872.

The two magnificent volumes before us afford a very complete view of the surgical knowledge of the day. Some years ago we had the pleasure of presenting the first edition of Gross's Surgery to the profession as a work of unrivalled excellence; and now we have the result of years of experience, labor, and study, all condensed upon the great work before us. And to students or practitioners desirous of enriching their library with a treasure of reference, we can simply commend the purchase of these two volumes of immense research.—*Cincinnati Lancet and Observer*, Sept. 1872.

A complete system of surgery—not a mere text-book of operations, but a scientific account of surgical theory and practice in all its departments.—*Brit. and For. Med. Chir. Rev.*, Jan. 1873.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON THE DISEASES, INJURIES, and Malformations of the Urinary Bladder, the Prostate Gland, and the Urethra. Third Edition, thoroughly Revised and Condensed, by SAMUEL W. GROSS, M.D., Surgeon to the Philadelphia Hospital. In one handsome octavo volume of 574 pages, with 170 illustrations: cloth, \$4 50. (Just Issued.)

For reference and general information, the physician or surgeon can find no work that meets their necessities more thoroughly than this, a revised edition of an excellent treatise, and no medical library should be without it. Replete with handsome illustrations and good ideas, it has the unusual advantage of being easily comprehended, by the reasonable and practical manner in which the various subjects are systematized and arranged. We heartily recommend it to the profession as a valuable addition to the important literature of dis-

eases of the urinary organs.—*Atlanta Med. Journ.*, Oct. 1876.

It is with pleasure we now again take up this old work in a decidedly new dress. Indeed, it must be regarded as a new book in very many of its parts. The chapters on "Diseases of the Bladder," "Prostate Body," and "Lithotomy," are splendid specimens of descriptive writing; while the chapter on "Stricture" is one of the most concise and clear that we have ever read.—*New York Med. Journ.*, Nov. 1876.

BY THE SAME AUTHOR.

A PRACTICAL TREATISE ON FOREIGN BODIES IN THE AIR-PASSAGES. In 1 vol. 8vo., with illustrations, pp. 468, cloth, \$2 75.

DRUITT (ROBERT), M.R.C.S., &c.

THE PRINCIPLES AND PRACTICE OF MODERN SURGERY.

A new and revised American, from the eighth enlarged and improved London edition. Illustrated with four hundred and thirty-two wood engravings. In one very handsome octavo volume, of nearly 700 large and closely printed pages, cloth, \$4 00; leather, \$5 00.

All that the surgical student or practitioner could desire.—*Dublin Quarterly Journal*.

It is a most admirable book. We do not know when we have examined one with more pleasure.—*Boston Med. and Surg. Journal*.

In Mr. Drutt's book, though containing only some seven hundred pages, both the principles and the

practice of surgery are treated, and so clearly and perspicuously, as to elucidate every important topic. We have examined the book most thoroughly, and can say that this success is well merited. His book moreover, possesses the inestimable advantages of having the subjects perfectly well arranged and classified and of being written in a style at once clear and succinct.—*Am. Journal of Med. Sciences*.

ASHHURST (JOHN, Jr.), M.D.,*Prof. of Clinical Surgery, Univ. of Pa., Surgeon to the Episcopal Hospital, Philadelphia.***THE PRINCIPLES AND PRACTICE OF SURGERY. Second edition, enlarged and revised. In one very large and handsome octavo volume of over 1000 pages, with 542 illustrations. Cloth, \$6; leather, \$7; half Russia, \$7.50. (Just Ready.)**

Conscientiousness and thoroughness are two very marked traits of character in the author of this book. Out of these traits largely has grown the success of his mental fruit in the past, and the present offer seems in no wise an exception to what has gone before. The general arrangement of the volume is the same as in the first edition, but every part has been carefully revised, and much new matter added.—*Phila. Med. Times*, Feb. 1, 1879.

We have previously spoken of Dr. Ashhurst's work in terms of praise. We wish to reiterate those terms here, and to add that no more satisfactory representation of modern surgery has yet fallen from the press. In point of judicial fairness, of power of condensation, of accuracy and conciseness of expression and thoroughly good English, Prof. Ashhurst has no superior among the surgical writers in America.—*Am. Practitioner*, Jan. 1879.

The attempt to embrace in a volume of 1000 pages the whole field of surgery, general and special, would be a hopeless task unless through the most tireless industry in collating and arranging, and the wisest judgment in condensing and excluding. These facilities have been abundantly employed by the author, and he has given us a most excellent treatise, brought up by the revision for the second edition to the latest date. Of course this book is not designed for specialists, but as a course of general surgical knowledge and for general practitioners, and as a text-book for students it is not surpassed by any that has yet appeared, whether of home or foreign authorship.—*N. Carolina Med. Journal*, Jan. 1879.

Ashhurst's Surgery is too well known in this country to require special commendation from us. This, its second edition, enlarged and thoroughly revised, brings it nearer our idea of a model text-book than any recently published treatise. Though numerous additions have been made, the size of the work is not materially increased. The main trouble of text-books of modern times is that they are too cumbersome. The student needs a book which will furnish him the most information in the shortest time. In every respect this work of Ashhurst is the model text-book—full, comprehensive and compact.—*Nashville Jour. of Med. and Surg.*, Jan. '79.

The favorable reception of the first edition is a guarantee of the popularity of this edition, which is fresh from the editor's hands with many enlargements and improvements. The author of this work is deservedly popular as an editor and writer, and his contributions to the literature of surgery have gained for him wide reputation. The volume now offered the profession will add new laurels to those already won by previous contributions. We can only add that the work is well arranged, filled with practical matter, and contains in brief and clear language all that is necessary to be learned by the student of surgery whilst in attendance upon lectures, or the general practitioner in his daily routine practice.—*Med. Med. Journal*, Jan. 1879.

The fact that this work has reached a second edition so very soon after the publication of the first one, speaks more highly of its merits than anything we might say in the way of commendation. It seems to have immediately gained the favor of students and physicians.—*Cincinnati Med. News*, Jan. '79.

BRYANT (THOMAS), F.R.C.S.,*Surgeon to Guy's Hospital.***THE PRACTICE OF SURGERY. Second American, from the Second and Revised English Edition. With Six Hundred and Seventy-two Engravings on Wood. In one large and very handsome imperial octavo volume of over 1000 large and closely printed pages. Cloth, \$6; leather, \$7. (Just Ready.)**

This work has enjoyed the advantage of two thorough revisions at the hand of the author since the appearance of the first American edition, resulting in a very notable enlargement of size and improvement of matter. In England this has led to the division of the work into two volumes, which are here comprised in one, the size being increased to a large imperial octavo, printed on a condensed but clear type. The series of illustrations has undergone a like revision, and will be found correspondingly improved.

The marked success of the work on both sides of the Atlantic shows that the author has succeeded in the effort to give to student and practitioner a sound and trustworthy guide in the practice of Surgery; while the simultaneous appearance of the present edition in England and in this country affords to the American reader the benefit of the most recent advances made abroad in surgical science.

There are so many text-books of surgery, so many written by skilled and distinguished hands, that to obtain the honor of a third edition in England is no light praise. Mr. Bryant merits this, by clearness of style, and good judgment in selecting the operations he recommends, in his new editions he goes carefully over the old grounds, in light of later research. On these and many allied points, Mr. Bryant is a calm and unpartisan observer, and his book throughout has the great merit of maintaining the true scientific, judicial tone of mind.—*Med. and Surg. Reporter*, March 22, 1879.

The work before us is the American reprint of the last London edition, and has the advantage over the latter in being of more convenient size, and in being compressed into one volume. The author has rewritten the greater part of the work, and has succeeded, in the amount of new matter added, in making it markedly distinctive from previous editions. A few extra pages have been added, and also a few new illustrations introduced. The publishers have presented the work in a creditable style. As a concise and practical manual of British surgery it is perhaps without an equal, and will doubtless always be a favorite text-book with the student and practitioner.—*N. Y. Med. Record*, March 22, 1879.

Another edition of this manual having been called for, the author has availed himself of the opportunity to make no few alterations in the substance as well as in the arrangement of the work, and, with a view to its improvement, has recast the materials and revised the whole. We ourselves are of the opinion that there is no better work on surgery extant.—*Cincinnati Med. News*, March, 1879.

Bryant's Surgery has been favorably received from the first, and evidently grows in the esteem of the profession with each succeeding edition. In glancing over the volume before us we find proof in almost every chapter of the thorough revision which the work has undergone, many parts having been cut out and replaced by matter entirely fresh.—*N. Y. Med. Journ.*, April, 1879.

Welcome as the new edition is, and as much as it is entitled to commendation, yet its appearance at this time is, in a certain sense, a matter of regret, as it will be in competition with another work, lately issued from the same press. But, the difficult task of forming a judgment as to the relative merits of Bryant and Ashhurst we will not attempt, but predict that, considering the high excellence of both, many others will likewise be forced to hesitate long in making choice between them.—*Cincinnati Lancet and Clinic*, March 22, 1879.

ERICHSEN (JOHN E.),*Professor of Surgery in University College, London, etc.*

THE SCIENCE AND ART OF SURGERY; being a Treatise on Surgical Injuries, Diseases, and Operations. Carefully revised by the author from the Seventh and enlarged English Edition. Illustrated by eight hundred and sixty two engravings on wood. In two large and beautiful octavo volumes of nearly 2000 pages: cloth, \$8 50; leather, \$10 50; half Russia, \$11 50. (*New Ready.*)

In revising this standard work the author has spared no pains to render it worthy of a continuance of the very marked favor which it has so long enjoyed, by bringing it thoroughly on a level with the advance in the science and art of surgery made since the appearance of the last edition. To accomplish this has required the addition of about two hundred pages of text, while the illustrations have undergone a marked improvement. A hundred and fifty additional wood-cuts have been inserted, while about fifty other new ones have been substituted for figures which were not deemed satisfactory. In its enlarged and improved form it is therefore presented with the confident anticipation that it will maintain its position in the front rank of text-books for the student, and of works of reference for the practitioner, while its exceedingly moderate price places it within the reach of all.

The seventh edition is before the world as the last work of surgical science. There may be monographs which excel it upon certain points, but as a compendium upon surgical principles and practice it is unrivalled. It will well reward practitioners to read it, for it has been a peculiar province of Mr. Erichsen to demonstrate the absolute interdependence of medical and surgical science. We need scarcely add, in conclusion, that we heartily commend the work to students that they may be grounded in a sound faith, and to practitioners as an invaluable guide at the bedside.—*Am Practitioner*, April, 1878.

It is no little compliment to say that this is the best edition Mr. Erichsen has ever produced of his well-known book. Besides inheriting the virtues of its predecessors, it possesses excellences quite its own. Having stated that Mr. Erichsen has incorporated into this edition every recent improvement in the science and art of surgery, it would be a supererogation to give a detailed criticism. In short, we unhesitatingly aver that we know of no other single work where the student and practitioner can gain at once so clear an insight into the principles of surgery, and so complete a knowledge of the exigencies of surgical practice.—*London Lancet*, Feb. 14, 1878.

For the past twenty years Erichsen's Surgery has maintained its place as the leading text-book, not only in this country, but in Great Britain. That it is able to hold its ground, is abundantly proven by the thoroughness with which the present edition has been revised, and by the large amount of valuable material that has been added. Aside from this, one hundred and fifty new illustrations have been inserted, including quite a number of microscopical appearances of pathological processes. So marked is this change for the better, that the work almost appears as an entirely new one.—*Med. Record*, Feb. 23, 1878.

Of the many treatises on Surgery which it has been our task to study, or our pleasure to read, there is none which in all points has satisfied us so well as the classic treatise of Erichsen. His polished, clear style, his freedom from prejudice and hobbies, his unsurpassed grasp of his subject, and vast clinical experience, qualify him admirably to write a model text-book. When we wish, at the least cost of time, to learn the most of a topic in surgery, we turn, by preference, to his work. It is a pleasure, therefore, to see that the appreciation of it is general, and has led to the appearance of another edition.—*Med. and Surg. Reporter*, Feb. 2, 1878.

Notwithstanding the increase in size, we observe that much old matter has been omitted. The entire work has been thoroughly written up, and not merely amended by a few extra chapters. A great improvement has been made in the illustrations. One hundred and fifty new ones have been added, and many of the old ones have been redrawn. The author highly appreciates the favor with which his work has been received by American surgeons, and has endeavored to render his latest edition more than ever worthy of their approval. That he has succeeded admirably, must, we think, be the general opinion. We heartily recommend the book to both student and practitioner.—*N. Y. Med. Journal*, Feb. 1878.

Erichsen has stood so prominently forward for years as a writer on Surgery, that his reputation is world wide, and his name is as familiar to the medical student as to the accomplished and experienced surgeon. The work is not a reprint of former editions, but has in many places been entirely rewritten. Recent improvements in surgery have not escaped his notice, various new operations have been thoroughly analyzed, and their merits thoroughly discussed. One hundred and fifty new wood-cuts add to the value of this work.—*N. O. Med. and Surg. Journal*, March, 1878.

HOLMES (TIMOTHY), M.D.,*Surgeon to St. George's Hospital, London.*

SURGERY, ITS PRINCIPLES AND PRACTICE. In one handsome octavo volume of nearly 1000 pages, with 411 illustrations. Cloth, \$6; leather, \$7; half Russia, \$7 50. (*Just Issued.*)

This is a work which has been looked for on both sides of the Atlantic with much interest. Mr. Holmes is a surgeon of large and varied experience, and one of the best known, and perhaps the most brilliant writer upon surgical subjects in England. It is a book for students—and an admirable one—and for the busy general practitioner. It will give a student all the knowledge needed to pass a rigid examination. The book fairly justifies the high expectations that were formed of it. Its style is clear and forcible, even brilliant at times, and the conciseness needed to bring it within its proper limits has not impaired

its force and distinctness.—*N. Y. Med. Record*, April 14, 1878.

It will be found a most excellent epitome of surgery by the general practitioner who has not the time to give attention to more minute and extended works and to the medical student. In fact, we know of no one we can more cordially recommend. The author has succeeded well in giving a plain and practical account of each surgical injury and disease, and of the treatment which is most commonly advisable. It will no doubt become a popular work in the profession, and especially as a textbook.—*Cincinnati Med. News*, April, 1876.

ASHTON ON THE DISEASES, INJURIES, AND MALFORMATIONS OF THE RECTUM AND ANUS: with remarks on Habitual Constipation. Second American, from the fourth and enlarged London Edition. With illustrations. In one 8vo. vol. of 287 pages, cloth, \$3 25.

SARGENT ON BANDAGING AND OTHER OPERATIONS OF MINOR SURGERY. New edition, with an additional chapter on Military Surgery. One 12mo. vol. of 383 pages with 184 wood-cuts. Cloth, \$1 75.

HAMILTON (FRANK H.), M.D.,
Professor of Fractures and Dislocations, &c., in Bellevue Hosp. Med. College, New York.

A PRACTICAL TREATISE ON FRACTURES AND DISLOCATIONS. Fifth edition, revised and improved. In one large and handsome octavo volume of nearly 800 pages, with 344 illustrations. Cloth, \$5 75; leather, \$6 75. (*Lately Issued.*)

There is no better work on the subject in existence than that of Dr. Hamilton. It should be in the possession of every general practitioner and surgeon.—*The Am. Journ. of Obstetrics*, Feb. 1876.

The value of a work like this to the practical physician and surgeon can hardly be over-estimated, and the necessity of having such a book revised to the latest dates, not merely on account of the practical importance

of its teachings, but also by reason of the medico-legal bearings of the cases of which it treats, and which have recently been the subject of useful papers by Dr. Hamilton and others, is sufficiently obvious to every one. The present volume seems to amply fill all the requisites. We can safely recommend it as the best of its kind in the English language, and not excelled in any other.—*Journ. of Nervous and Mental Disease*, Jan. 1876.

NETTLESHIP (EDWARD), F.R.C.S.,
Ophthalmic Surg. and Lect. on Ophth. Surg. at St. Thomas' Hospital, London.

MANUAL OF OPHTHALMIC MEDICINE. In one royal 12mo. volume of over 350 pages, with 89 illustrations. Cloth, \$2. (*Just Ready.*)

The book is written in a careful and logical manner, and though extremely concise, we have failed to notice any evidence of ambiguity. It is rendered more compact and homogeneous by frequent references, by page number, to other portions of the work; repetitions are thus avoided, and we have been surprised to find how much information our author has succeeded in conveying in so small a space. A careful study of the book will well repay the general practitioner, even though it should serve only as a *monitor*. It is particularly useful in the latter regard, as the subject of treatment is presented in a thoroughly conservative manner.—*N. Y. Med. Record*, March 6, 1880.

The author has succeeded in touching upon about all the points, operations, diseases of the eye in relation to general diseases, and has prepared a very acceptable book.—*Cincinnati Lancet and Clinic*, Feb. 7, 1880.

It is *multum in parvo*, containing all the leading points to be remembered in the pathology, description, and treatment of diseases of the eye. It will be found especially valuable in preparing for examinations. Practitioners will find it convenient as a work of reference, when they wish to refresh their

memories in respect to the features of some affections.—*Cincinnati Med. News*, Jan. 1880.

The author is to be congratulated upon the very successful manner in which he has accomplished his task; he has succeeded in being concise without sacrificing clearness, and, including the whole ground covered by more voluminous text-books, has given an excellent *résumé* of all the practical information they contain. We do not hesitate to pronounce Mr. Nettleship's book the best manual on ophthalmic surgery for the use of students' and "busy practitioners," with which we are acquainted.—*Am. Jour. Med. Sciences*, April, 1880.

A careful examination has convinced us that it is the best work of its class that has come to our notice. While all matter is condensed to the utmost, there are few points that are obscured thereby, and all are rendered really attractive to the average student. We can only hope that every medical student shall be compelled to master the entire volume ere receiving his degree. General practitioners who take care of eye cases would do well to refresh their memories by its careful study.—*Detroit Lancet*, April, 1880.

CARTER (R. BRUDENELL), F.R.C.S.,
Ophthalmic Surgeon to St. George's Hospital, etc.

A PRACTICAL TREATISE ON DISEASES OF THE EYE. Edited, with test-types and Additions, by JOHN GREEN, M.D. (of St. Louis, Mo.). In one handsome octavo volume of about 500 pages, and 124 illustrations. Cloth, \$3 75. (*Just Issued.*)

It is with great pleasure that we can endorse the work as a most valuable contribution to practical ophthalmology. Mr. Carter never deviates from the end he has in view, and presents the subject in a clear and concise manner, easy of comprehension, and hence the more valuable. We would especially commend, however, as worthy of high praise, the manner in which the therapeutics of disease of the eye is elaborated, for here the author is particularly clear and practical, where other writers are unfortunately too often deficient. The final

chapter is devoted to a discussion of the uses and selection of spectacles, and is admirably compact, plain, and useful, especially the paragraphs on the treatment of presbyopia and myopia. In conclusion, our thanks are due the author for many useful hints in the great subject of ophthalmic surgery and therapeutics, a field where of late years we glean but a few grains of sound wheat from a mass of chaff.—*New York Medical Record*, Oct. 23, 1875.

WELLS (J. SOELBERG),
Professor of Ophthalmology in King's College Hospital, &c.

A TREATISE ON DISEASES OF THE EYE. Third American, from the Third London Edition. Thoroughly revised, with copious additions, by Chas. S. Bull, M.D., Surgeon and Pathologist to the New York Eye and Ear Infirmary. Illustrated with about 250 engravings on wood, and six colored plates. Together with selections from the Test-types of Jaeger and Snellen. In one large and very handsome octavo volume of 900 pages. (*In Press.*)

BROWNE (EDGAR A.),
Surgeon to the Liverpool Eye and Ear Infirmary, and to the Dispensary for Skin Diseases.

HOW TO USE THE OPHTHALMOSCOPE. Being Elementary Instructions in Ophthalmoscopy, arranged for the Use of Students. With thirty-five illustrations. In one small volume royal 12mo. of 120 pages: cloth, \$1. (*Now Ready.*)

LAURENCE'S HANDY-BOOK OF OPHTHALMIC SURGERY, for the use of Practitioners. Second edition, revised and enlarged. With numerous illustrations. In one very handsome octavo volume, cloth, \$2 75.

LAWSON'S INJURIES TO THE EYE, ORBIT, AND EYELIDS: their immediate and Remote Effects. With about one hundred illustrations. In one very handsome octavo volume, cloth, \$3 50.

BURNETT (CHARLES H.), M.A., M.D.,

Aural Surg. to the Presb. Hosp., Surgeon-in-charge of the Infr. for Dis. of the Ear, Phila.

THE EAR, ITS ANATOMY, PHYSIOLOGY, AND DISEASES.

A Practical Treatise for the Use of Medical Students and Practitioners. In one handsome octavo volume of 615 pages, with eighty-seven illustrations: cloth, \$4 50; leather, \$5 50; half Russia, \$6 00. (Now Ready.)

Recent progress in the investigation of the structures of the ear, and advances made in the modes of treating its diseases, would seem to render desirable a new work in which all the resources of the most advanced science should be placed at the disposal of the practitioner. This it has been the aim of Dr. Burnett to accomplish, and the advantages which he has enjoyed in the special study of the subject are a guarantee that the result of his labors will prove of service to the profession at large, as well as to the specialist in this department.

Foremost among the numerous recent contributions to aural literature will be ranked this work of Dr. Burnett. It is impossible to do justice to this volume of over 600 pages in a necessarily brief notice. It must suffice to add that the book is profusely and accurately illustrated, the references are conscientiously acknowledged, while the result has been to produce a treatise which will henceforth rank with the classic writings of Wilde and Von Trösch.—*The Lond. Practitioner*, May, 1879.

On account of the great advances which have been made of late years in otology, and of the increased interest manifested in it, the medical profession will welcome this new work, which presents clearly and concisely its present aspect, whilst clearly indicating the direction in which further researches can be most profitably carried on. Dr. Burnett from his own matured experience, and availing himself of the observations and discoveries of others, has produced a work, which as a text-book, stands *factis princeps* in our language. We had marked several passages as well worthy of quotation and the attention of the general practitioner, but their number and the space at our command forbid. Perhaps it is better, as the book ought to be in the hands of every

medical student, and its study will well repay the busy practitioner in the pleasure he will derive from the agreeable style in which many otherwise dry and mostly unknown subjects are treated. To the specialist the work is of the highest value, and his sense of gratitude to Dr. Burnett will, we hope, be proportionate to the amount of benefit he can obtain from the careful study of the book, and a constant reference to its trustworthy pages.—*Edinburgh Med. Jour.*, Aug. 1878.

The book is designed especially for the use of students and general practitioners, and places at their disposal much valuable material. Such a book as the present one, we think, has long been needed, and we may congratulate the author on his success in filling the gap. Both student and practitioner can study the work with a great deal of benefit. It is profusely and beautifully illustrated.—*N. Y. Hospital Gazette*, Oct. 16, 1877.

Dr. Burnett is to be commended for having written the best book on the subject in the English language, and especially for the care and attention he has given to the scientific side of the subject.—*N. Y. Med. Journ.*, Dec. 1877.

TAYLOR (ALFRED S.), M.D.,

Lecturer on Med. Jurisp. and Chemistry in Guy's Hospital.

POISONS IN RELATION TO MEDICAL JURISPRUDENCE AND

MEDICINE. Third American, from the Third and Revised English Edition. In one large octavo volume of 850 pages; cloth, \$5 50; leather, \$6 50. (Just Issued.)

The present is based upon the two previous editions; "but the complete revision rendered necessary by time has converted it into a new work." This statement from the preface contains all that it is desired to know in reference to the new edition. The works of this author are already in the library of every physician who is liable to be called upon for medico-legal testimony (and what one is not), so that all that is required to be known about the present book is that the author has kept it abreast with the times. What makes it now, as always, especially valuable to the practitioner is its conciseness and practical character, only those poisonous substances

being described which give rise to legal investigations.—*The Clinic*, Nov. 6, 1876.

Dr. Taylor has brought to bear on the compilation of this volume, stores of learning, experience, and practical acquaintance with his subject, probably far beyond what any other living authority on toxicology could have amassed or utilized. He has fully sustained his reputation by the consummate skill and legal acumen he has displayed in the arrangement of the subject-matter, and the result is a work on Poisons which will be indispensable to every student or practitioner in law and medicine.—*The Dublin Journ. of Med. Sci.*, Oct. 1876.

BY THE SAME AUTHOR.

MEDICAL JURISPRUDENCE. Seventh American Edition. Edited

by JOHN J. REESE, M.D., Prof. of Med. Jurisp. in the Univ. of Penn. In one large octavo volume of nearly 900 pages. Cloth, \$5 00; leather, \$6 00. (Lately Issued.)

To the members of the legal and medical profession, it is unnecessary to say anything commendatory of Taylor's Medical Jurisprudence. We might as well undertake to speak of the merit of Chitty's Pleadings.—*Chicago Legal News*, Oct. 16, 1873.

It is beyond question the most attractive as well as most reliable manual of medical jurisprudence published in the English language.—*Am. Journal of Syphillography*, Oct. 1873.

It is altogether superfluous for us to offer anything in behalf of a work on medical jurisprudence by an author who is almost universally esteemed to be the

best authority on this specialty in our language. On this point, however, we will say that we consider Dr. Taylor to be the safest medico-legal authority to follow, in general, with which we are acquainted in any language.—*Va. Clin. Record*, Nov. 1873.

This latest edition of the Manual is probably the best of all, as it contains more material and is worked up to the latest views of the author as expressed in the last edition of the Principles. Dr. Reese, the editor of the Manual, has done everything to make his work acceptable to his medical countrymen.—*N. Y. Med. Record*, Jan. 16, 1874.

BY THE SAME AUTHOR.

THE PRINCIPLES AND PRACTICE OF MEDICAL JURISPRU-

DENCE. Second Edition, Revised, with numerous Illustrations. In two large octavo volumes, cloth, \$10 00; leather, \$12 00

This great work is now recognised in England as the fullest and most authoritative treatise on every department of its important subject. In laying it, in its improved form, before the American profession, the publishers trust that it will assume the same position in this country.

ROBERTS (WILLIAM), M.D.,
Lecturer on Medicine in the Manchester School of Medicine, &c.

A PRACTICAL TREATISE ON URINARY AND RENAL DISEASES, including Urinary Deposits. Illustrated by numerous cases and engravings. Third American, from the Third Revised and Enlarged London Edition. In one large and handsome octavo volume of over 600 pages. Cloth, \$4. (*Just Ready.*)

THOMPSON (SIR HENRY),
Surgeon and Professor of Clinical Surgery to University College Hospital.

LECTURES ON DISEASES OF THE URINARY ORGANS. With illustrations on wood. Second American from the Third English Edition. In one neat octavo volume. Cloth, \$2 25. (*Just Issued.*)

BY THE SAME AUTHOR.

ON THE PATHOLOGY AND TREATMENT OF STRICTURE OF THE URETHRA AND URINARY FISTULÆ. With plates and wood-cuts. From the third and revised English edition. In one very handsome octavo volume, cloth, \$3 50. (*Lately Published.*)

TUKE (DANIEL HACK), M.D.,
Joint author of "The Manual of Psychological Medicine," &c.

ILLUSTRATIONS OF THE INFLUENCE OF THE MIND UPON THE BODY IN HEALTH AND DISEASE. Designed to illustrate the Action of the Imagination. In one handsome octavo volume of 416 pages, cloth, \$3 25. (*Lately Issued.*)

BLANDFORD (G. FIELDING), M.D., F.R.C.P.,
Lecturer on Psychological Medicine at the School of St. George's Hospital, &c.

INSANITY AND ITS TREATMENT: Lectures on the Treatment, Medical and Legal, of Insane Patients. With a Summary of the Laws in force in the United States on the Confinement of the Insane. By ISAAC RAY, M. D. In one very handsome octavo volume of 471 pages; cloth, \$3 25.

It satisfies a want which must have been sorely felt by the busy general practitioners of this country. It takes the form of a manual of clinical description of the various forms of insanity, with a description of the mode of examining persons suspected of insanity. We call particular attention to this feature of the book, as giving it a unique value to the general practitioner. If we pass from theoretical considerations to descriptions of the varieties of insanity as

actually seen in practice and the appropriate treatment for them, we find in Dr. Blandford's work a considerable advance over previous writings on the subject. His pictures of the various forms of mental disease are so clear and good that no reader can fail to be struck with their superiority to those given in ordinary manuals in the English language or (so far as our own reading extends) in any other.—*London Practitioner*, Feb. 1871.

LEA (HENRY C.).

SUPERSTITION AND FORCE: ESSAYS ON THE WAGER OF LAW, THE WAGER OF BATTLE, THE ORDEAL, AND TORTURE. Third Revised and Enlarged Edition. In one handsome royal 12mo. volume of 552 pages. Cloth, \$2 50. (*Just Ready.*)

This valuable work is in reality a history of civilization as interpreted by the progress of jurisprudence. . . . In "Superstition and Force" we have a philosophic survey of the long period intervening between primitive barbarity and civilized enlightenment. There is not a chapter in the work that should not be most carefully studied, and however well versed the reader may be in the science of jurisprudence, he will find much in Mr. Lea's volume of which he was previously ignorant. The book is a valuable addition to the literature of social science.—*Westminster Review*, Jan. 1880.

The appearance of a new edition of Mr. Henry C. Lea's "Superstition and Force" is a sign that our highest scholarship is not without honor in its native country. Mr. Lea has met every fresh demand for his work with a careful revision of it, and the present edition is not only fuller and, if possible,

more accurate than either of the preceding, but, from the thorough elaboration, is more like a harmonious concert and less like a batch of studies.—*The Nation*, Aug. 1, 1878.

Many will be tempted to say that this, like the "Decline and Fall," is one of the uncriticizable books. Its facts are innumerable, its deductions simple and inevitable, and its *chevaux-de-frise* of references bristling and dense enough to make the keenest, stoutest, and best equipped assailant think twice before advancing. Nor is there anything controversial in it to provoke assault. The author is no polemic. Though he obviously feels and thinks strongly, he succeeds in attaining impartiality. Whether looked on as a picture or a mirror, a work such as this has a lasting value.—*Lippincott's Magazine*, Oct. 1878.

BY THE SAME AUTHOR. (*Lately Published.*)

STUDIES IN CHURCH HISTORY—THE RISE OF THE TEMPORAL POWER—BENEFIT OF CLERGY—EXCOMMUNICATION. In one large royal 12mo. volume of 516 pp.; cloth, \$2 75.

The story was never told more calmly or with greater learning or wiser thought. We doubt, indeed, if any other study of this field can be compared with this for clearness, accuracy, and power.—*Chicago Examiner*, Dec. 1870.

Mr. Lea's latest work, "Studies in Church History," fully sustains the promise of the first. It deals with three subjects—the Temporal Power, Benefit of Clergy, and Excommunication, the record of which


has a peculiar importance for the English student, and is a chapter on Ancient Law likely to be regarded as final. We can hardly pass from our mention of such works as these—with which that on "Sacredotal Celibacy" should be included—without noting the literary phenomenon that the head of one of the first American houses is also the writer of some of its most original books.—*London Athenæum*, Jan. 7, 1871.

INDEX TO CATALOGUE.

	PAGE		PAGE
American Journal of the Medical Sciences	1	Hodge on Women	21
Allen's Anatomy	7	Hodge's Obstetrics	25
Anatomical Atlas, by Smith and Horner	7	Holland's Medical Notes and Reflections	14
Ashton on the Rectum and Anus	28	*Holmes's Surgery	28
Attfield's Chemistry	10	Holden's Landmarks	6
Ashwell on Diseases of Females	21	Horner's Anatomy and Histology	7
*Ashhurst's Surgery	27	Hudson on Fever	19
Browne on Ophthalmoscope	29	Hill on Venereal Diseases	20
Browne on the Throat	19	Hillier's Handbook of Skin Diseases	19
*Burnett on the Ear	30	Jooss (C. Handfield) on Nervous Disorders	18
*Barnes on Diseases of Women	22	Knapp's Chemical Technology	31
Barnes' Midwifery	24	Lea's Superstition and Force	31
Bellamy's Surgical Anatomy	7	Lea's Studies in Church History	20
Bryant's Practice of Surgery	27	Lee on Syphilis	25
Bloxam's Chemistry	31	*Leishman's Midwifery	14
Blandford on Insanity	11	La Roche on Yellow Fever	19
Basham on Renal Diseases	19	La Roche on Pneumonia, &c.	29
Bartholow on Electricity	18	Laurence and Moon's Ophthalmic Surgery	29
Barlow's Practice of Medicine	14	Lawson on the Eye	9
Bowman's (John E.) Practical Chemistry	18	Lehmann's Physiological Chemistry, 2 vols.	9
*Bristowe's Practice	20	Lehmann's Chemical Physiology	9
*Bumstead on Venereal	20	Ludlow's Manual of Examinations	19
Bumstead and Cuillerier's Atlas of Venereal	20	Lyons on Fever	2
*Carpenter's Human Physiology	12	Medical News and Abstract	18
Carpenter on the Use and Abuse of Alcohol	14	Morris on Skin Diseases	21
*Cornill and Ranvier	29	Meigs on Puerperal Fever	25
Carter on the Eye	7	Miller's Practice of Surgery	25
Cleland's Dissector	9	Miller's Principles of Surgery	21
Classen's Chemistry	11	Montgomery on Pregnancy	29
Clowes' Chemistry	5	Nettleship's Ophthalmic Medicine	6
Century of American Medicine	22	Neill and Smith's Compendium of Med. Science	24
Chadwick on Diseases of Women	18	Obstetrical Journal	25
Charcot on the Nervous System	19	Parry on Extra-Uterine Pregnancy	14
Chambers on Diet and Regimen	12	Pavy on Digestion	11
Christison and Griffith's Dispensatory	21	*Parrish's Practical Pharmacy	25
Churchill's System of Midwifery	21	Pirrie's System of Surgery	24
Churchill on Puerperal Fever	21	*Playfair's Midwifery	7
Condie on Diseases of Children	25	Quain and Sharpey's Anatomy, by Leidy	17
Cooper's (B. E.) Lectures on Surgery	20	*Reynolds' System of Medicine	16
Cuillerier's Atlas of Venereal Diseases	15	Richardson's Preventative Medicine	31
Cyclopaedia of Practical Medicine	23	Roberts on Urinary Diseases	28
Duncan on Diseases of Women	9	Ramsbotham on Parturition	9
*Dalton's Human Physiology	15	Remsen's Principles of Chemistry	21
Davis's Clinical Lectures	21	Rigby's Midwifery	4
Dawson on Diseases of Females	26	Rodwell's Dictionary of Science	25
*Dewitt's Modern Surgery	4	Simson's Operative Surgery	21
*Dunngilson's Medical Dictionary	7	Swayne's Obstetric Aphorisms	19
Ellis's Demonstrations in Anatomy	28	Seller on the Throat	28
*Erichsen's System of Surgery	23	Sargent's Minor Surgery	7
*Emmet on Diseases of Women	12	Sharpey and Quain's Anatomy, by Leidy	26
Farquharson's Therapeutics	8	Skay's Operative Surgery	19
Foster's Physiology	14	Slade on Diphtheria	7
Fenwick's Diagnosis	18	Schäfer's Histology	21
Finlayson's Clinical Diagnosis	19	*Smith (J. L.) on Children	7
Flint on Respiratory Organs	19	Smith (H. H.) and Horner's Anatomical Atlas	19
Flint on the Heart	16	Smith (Edward) on Consumption	21
Flint's Practice of Medicine	15	Smith on Wasting Diseases in Children	12
Flint's Essays	15	*Stillé's Therapeutics	13
*Flint's Clinical Medicine	19	*Stillé & Malsch's Dispensatory	16
Flint on Phthisis	19	Sturges on Clinical Medicine	14
Flint on Percussion	16	Stokes on Fever	5
Fothergill's Handbook of Treatment	16	Tanner's Manual of Clinical Medicine	28
Fothergill's Antagonism of Therapeutic Agents	10	Tanner on Pregnancy	30
Fownes' Elementary Chemistry	18	Taylor's Medical Jurisprudence	30
Fox on Diseases of the Skin	19	Taylor's Principles and Practice of Med. Jurisp	31
Fuller on the Lungs, &c.	9	Taylor on Poisons	22
Green's Pathology and Morbid Anatomy	25	Tuke on the Influence of the Mind	31
Greene's Medical Chemistry	14	*Thomas on Diseases of Females	31
Gibson's Surgery	9	Thompson on Urinary Organs	14
Gluge's Pathological Histology, by Leidy	16	Thompson on Stricture	16
*Gray's Anatomy	12	Todd on Acute Diseases	19
Galloway's Analysis	26	Woodbury's Practice	15
Griffith's (R. E.) Universal Formulary	26	Walshe on the Heart	20
Gross on Urinary Organs	18	Watson's Practice of Physic	20
Gross on Foreign Bodies in Air-Passages	26	Wells on the Eye	19
*Gross's System of Surgery	26	West on Diseases of Females	20
Habershon on the Abdomen	18	West on Diseases of Children	20
Hamilton on Dislocations and Fractures	29	West on Nervous Disorders of Children	19
Hartshorne's Essentials of Medicine	15	Williams on Consumption	7
Hartshorne's Conspectus of the Medical Sciences	6	Wilson's Human Anatomy	19
Hartshorne's Anatomy and Physiology	8	Wilson's Handbook of Cutaneous Medicine	9
Hamilton on Nervous Diseases	18	Wöhler's Organic Chemistry	28
Heath's Practical Anatomy	6	Winckel on Childbed	
Hoblyn's Medical Dictionary	4		

Books marked * are also bound in half Russia.

HENRY C. LEA'S SON & CO.—Philadelphia.



LANE MEDICAL LIBRARY

To avoid fine, this book should be returned
on or before the date last stamped below.

--	--	--

J111 Green, T.H. 13132
G79 Introduction to
1878 pathology and morbid
anatomy. 3d American
ed.

